



Antibodies for HIV prevention in young women

Salim S. Abdool Karim^{a,b}, Quarraisha Abdool Karim^{a,b}, and Cheryl Baxter^a

Purpose of review

Young women in sub-Saharan Africa bear a disproportionate HIV burden. They urgently require new HIV prevention approaches that they can use. This review provides an overview of the use of antiretrovirals for HIV preexposure prophylaxis (PrEP), highlighting some of the challenges with this technology and explores the potential role of mAbs for HIV prevention in women.

Recent findings

Recent findings on the initial steps in viral entry and establishment of a productive local infectious nidus in the vaginal epithelium has provided important clues for HIV prevention in the female genital tract. Topical and oral formulations of antiretroviral drugs have been shown to prevent HIV infection in women with varying levels of success, depending principally on adherence. Further, several new broad and potent mAbs have been isolated over the last 5 years. Nonhuman primate studies demonstrate that broadly neutralizing HIV mAbs can protect rhesus macaques from simian immunodeficiency virus–HIV chimera (SHIV) infection. These findings have created newfound enthusiasm for passive immunization as a potential prevention strategy for women.

Summary

If potent broadly neutralizing mAbs are effective in preventing HIV infection in women, this outcome could fill an important gap in HIV prevention technologies for young women, especially in Africa.

Keywords

HIV prevention, mAb, passive immunization, young women

INTRODUCTION

There has been a substantial decline in new HIV infections globally over the past decade. Estimates from the Joint United Nations Programme on HIV/AIDS (UNAIDS) indicate a 38% drop in the number of new HIV infections annually from 3.4 million in 2001 to 2.1 million in 2013 [1]. Despite these encouraging trends, HIV remains a substantial global health challenge with an estimated 35 million people living with HIV in 2013 [1]. In most countries, even those with a declining overall prevalence, HIV continues to spread in certain key populations. About a quarter of all new HIV infections occurring globally are in young women.

HIV IN YOUNG WOMEN IN SUB-SAHARAN AFRICA

In sub-Saharan Africa, where just over 70% of all new HIV infections occur, young women bear a disproportionate burden of HIV infection. In this region, not only do young women aged 15–24 years have HIV rates up to eight-fold higher than their male peers [2], but they also acquire HIV infection at least 5–7 years earlier than their male peers [3,4].

Although HIV prevalence in other sub-Saharan African countries do not reach the same levels as those observed in South Africa, similar trends of higher HIV prevalence among young women than young men occur throughout eastern and southern Africa (Table 1) [5].

Many countries in southern Africa have substantial HIV burdens. For example, South Africa, which is home to less than 1% of the global population, accounts for about 17% of the global burden of HIV infection. The HIV prevalence in the South African general population is estimated to be about 12% [6]. HIV continues to spread rapidly in South Africa and

^aCAPRISA - Centre of AIDS Programme of Research in South Africa, Nelson R Mandela School of Medicine, University of KwaZulu-Natal, KwaZulu-Natal, South Africa and ^bDepartment of Epidemiology, Columbia University, New York, New York, USA

Correspondence to Salim S. Abdool Karim, CAPRISA, 2nd Floor, Doris Duke Medical Research Institute, Nelson R Mandela School of Medicine, University of KwaZulu-Natal, Private Bag X7, Congella, 4013, Durban, South Africa. Tel: +27 31 260 4550; fax: +27 31 260 4549; e-mail: Salim.Abdoolkarim@caprisa.org

Curr Opin HIV AIDS 2015, 10:183–189

DOI:10.1097/COH.0000000000000147

KEY POINTS

- New HIV prevention approaches that women can use are urgently required.
- Topical and oral formulations of antiretroviral drugs have been shown to prevent HIV infection in women with varying levels of success, depending principally on adherence.
- Several new broad and potent mAbs have been isolated over the last 5 years.
- Nonhuman primate studies demonstrate that some of these mAbs can protect rhesus macaques from SHIV infection.
- If potent broadly neutralizing mAbs are effective in preventing HIV infection in women, it could fill an important gap in HIV prevention technologies for young women, especially in Africa.

many other countries in southern Africa resulting in 'generalized' epidemics, as described by UNAIDS. One of the key drivers of the continued spread of HIV infection in this region is the age–sex difference in HIV acquisition between young boys and girls [5]. Temporal trends of the evolving HIV epidemic in this region have been monitored mainly through annual seroprevalence surveys in pregnant women. In South Africa, HIV prevalence among pregnant women has increased from 0.8% in 1990 to 29.5% in 2012 [7]. Despite the rapid scale-up of antiretroviral therapy provision in South Africa from less than 50 000 in 2004 to about 2.6 million in 2012, HIV prevalence among pregnant women utilizing public

sector facilities has remained stable at about 29.5% for the past 7 years. In the context of having the largest AIDS treatment programme in the world, South Africa has not yet witnessed an increase in HIV prevalence reflecting the survival benefits of antiretroviral therapy because of the continued high mortality rates, especially in HIV-tuberculosis (TB) coinfecting patients. In 2012, HIV prevalence rates among pregnant women exceeded 40% in five sub-districts within South Africa [7]. In one of these high-burden subdistricts, annual cross-sectional surveys of antenatal clinic attendees demonstrate a concerning rise of HIV infection among young women below the age of 20 years, increasing from 13.0% in 2007 to 15.1% in 2010 and 22.1% in 2013. The survey in 2013 revealed that HIV prevalence was 39.7% among pregnant women aged 20–24 years and 63.1% among those aged 25–29 years.

Several cohort studies conducted in South Africa between 2002 and 2010 [8–15] demonstrate high HIV incidence rates (Table 2). In the CAPRISA 004 trial, wherein intensive monthly risk reduction counselling was provided, the HIV incidence rate was 9.1 per 100 women-years among 18-to 40-year-old women in the placebo arm [16].

WHAT MAKES YOUNG WOMEN MORE VULNERABLE TO HIV?

A complex interplay of biology, sex-power disparities, social, political and economic factors contribute to the excess vulnerability of young women to HIV infection compared with men. The per-act probability of acquiring HIV for women is estimated to be 1 per 1000 sexual encounters compared with 1

Table 1. HIV prevalence (%) among people 15–24 years old, by sex in selected African countries, 2008–2011

Country	HIV prevalence (%)		Fold difference
	Females	Males	
South Africa	14.1	4.0	3.5
Lesotho	13.9	4.9	2.8
Mozambique	13.5	10	1.4
Botswana	11.5	5.5	2.1
Zambia	9.4	4.9	1.9
Zimbabwe	8.1	4.0	2.0
Malawi	5.9	2.0	3.0
Kenya	5.1	1.3	3.9
Central African Republic	4.8	0.9	5.3
Tanzania	4.1	1.2	3.4
Congo	2.8	0.9	3.1
Rwanda	1.8	0.4	4.5

Data adapted from [5].

Table 2. HIV incidence rates from cohort studies that have been conducted in South Africa between 2002 and 2010

Years of study conduct	Population (age range in years)	Location	HIV incidence rate (per 100 person-years) [95% CI]
2002–2005	2523 HIV-negative women (18–40)	KwaZulu-Natal, South Africa	6.6 overall 10.0 in women ≤ 24 years old 6.0 in women 25–34 years old 3.6 in women ≥ 35 years old
2003–2004	958 HIV-uninfected women (18–35)	KwaZulu-Natal, South Africa	3.8 [2.6–5.2] overall 5.3 [2.7–9.2] in Durban 6.2 [3.4–10.5] in Hlabisa
2004–2007	594 HIV-uninfected women (14–30)	KwaZulu-Natal, South Africa	4.7 [1.5–10.9] in < 18 years 6.9 [4.8–9.6] in ≥ 18 years
2004–2007	594 HIV-uninfected women (14–30)	KwaZulu-Natal, South Africa	6.5 [4.4–9.2] in rural women 6.4 [2.6–13.2] in urban women 17.2 [2.1–62.2] in urban women < 20 years
2005–2009	1048 HIV-uninfected women (≥ 18)	2 sites in KwaZulu-Natal, South Africa	4.9 [2.8–8.1] in Durban 7.7 [4.1–13.2] in Hlabisa
2007–2008	598 sexually active HIV-negative women (18–35)	North-West Province and Western Cape, South Africa	6.0 [3.0–9.0] in North-West 4.5 [1.8–7.1] in Western Cape
2007–2010	444 HIV-negative women (18–40)	KwaZulu-Natal, South Africa	9.1 [6.9–11.7]
2008–2009	624 high-risk HIV-negative women (18–35)	Bloemfontein and Rustenburg, South Africa	5.5 [2.5–10.4] in Bloemfontein 3.0 [0.4–10.8] in Rustenburg

CI, confidence interval.

per 2000 encounters for the male partner in peno-vaginal sex [17]. Thus, women appear to be biologically more likely to become infected than men. Factors such as intimate partner violence [18] and early sexual debut [19] have also been shown to be associated with an increased risk of acquiring HIV in women. Sexual debut also marks the initial exposure to a number of sexually transmitted pathogens, including viruses such as herpes simplex virus type-2 and human papilloma virus that also increase the risk of HIV acquisition [20–23]. Teenage pregnancy rates are also high in sub-Saharan Africa, leading to girls dropping out from school, which itself is a risk factor for HIV acquisition [24]. Although controversial, some studies have suggested that the long-acting injectable hormonal contraceptive, Depo-Provera (DMPA), may be associated with an increased HIV risk [25], and this effect may be further amplified in younger women. Data from several African countries have shown that young women who have sexual partners who are 5–10 years older than them are at an increased risk for acquiring HIV [26–30]. In some cases, young women, particularly those from impoverished backgrounds, form relationships with older men for financial and social

security [31]. Young people are also more likely to be inexperienced in sexual risk-taking and they may not be able to negotiate condom use with older partners.

EXISTING HIV PREVENTION TECHNOLOGIES

Several effective prevention strategies are already available but are often not being implemented at the necessary scale and magnitude to those most in need. For example, although condoms are highly effective in preventing HIV, UNAIDS estimates that in sub-Saharan Africa, each sexually active individual only has access to about eight male condoms per year [1].

Further, traditional behaviour change prevention programmes, based on the ABCC (Abstinence, Be faithful, Condoms and Circumcision) approach, have had a little, if any, impact in women in sub-Saharan Africa. Programmes promoting abstinence have been unsuccessful in delaying the age of sexual debut. Data from a national HIV prevalence, incidence and behaviour survey in South Africa show that rates of reported early sexual debut, that is, before the age of 15 years, has remained almost

stable at 10% between 2002 and 2012 [6]. Further, abstinence is an inappropriate prevention strategy for women in relationships or who are married. Sexual inequality and the threat of sex-based violence also limits a woman's ability to convince her male partner to use condoms, to remain faithful or have an HIV test. Medical male circumcision, which is a proven HIV prevention strategy for men [32], has a little immediate direct benefit to the woman [33]. Although male circumcision could potentially impact on HIV incidence rates in women in the long-term through diminished exposure to HIV because of lower HIV levels among circumcised men, this prevention option is not controlled by women.

The development of women initiated HIV prevention strategies could make a potentially important impact in this 'key population', as it offers women who are unable to negotiate safer sex with their partner an option to remain HIV uninfected. Two promising HIV prevention options that could be of direct benefit to women are topical (microbicides) and oral antiretroviral drugs administered as preexposure prophylaxis (PrEP).

Microbicides, which are chemical products designed to prevent the sexual acquisition of HIV, have been in development since the early 1990s. Although no microbicides are licensed or available yet, the CAPRISA 004 tenofovir gel trial [16] provided proof-of-concept that topical antiretroviral microbicides, when used before and after sex, can reduce sexual transmission of HIV and herpes simplex type-2 (HSV-2). The overall protective effect of tenofovir gel against HIV was a modest 39%, with the most adherent women achieving 54% protection and those with genital tenofovir concentrations in excess of 1000 ng/ml demonstrating 74% protection [34]. A confirmatory study of the effectiveness of tenofovir gel, the FACTS 001 [35] trial, is currently underway and the results are anticipated in the first quarter of 2015. A successful outcome from the FACTS 001 trial could lead to licensure of tenofovir gel as the first microbicide for HIV prevention.

Oral antiretrovirals as PrEP have also been shown in clinical trials to reduce the risk of HIV acquisition in HIV-negative partners in discordant couples [36], MSM [37], at-risk men and women [38], and people who inject drugs (PWID) [39,40] between 44 and 75%. The initial studies tested daily oral antiretrovirals for PrEP. In late 2014, a trial in the UK and one in France testing coital dosing of oral PrEP announced that their Data Safety Monitoring Boards had recommended the trials stop as participants in the active arms had shown significant benefits that the placebo groups could no

longer be justified. Despite the potential impact that PrEP could have on the epidemic, only the USA has approved the oral antiretroviral, Truvada, as an HIV prevention option [41]. Scale-up of this option has, however, been suboptimal.

Apart from the unavailability of PrEP in most countries, several other challenges could hinder the rapid scale-up of PrEP. The most significant challenge in using antiretrovirals either as microbicides or as oral PrEP is inadequate adherence to the prescribed dosing regimen. Several microbicide and PrEP trials, testing the same products, have produced vastly different findings. The failure to demonstrate a protective effect in some of the trials has been ascribed to suboptimal adherence. For example, in the FEM-PrEP trial [42], only 24% of the women allocated to the daily oral Truvada [a combination of tenofovir disoproxil fumarate (TDF) and emtricitabine (FTC)] group had detectable drug levels. Similarly, in the VOICE trial, only 25, 29 and 30% of women allocated to the daily tenofovir gel, daily oral TDF and daily oral TDF/FTC groups, respectively, had detectable drug levels [43]. Even in successful trials, the degree of adherence directly impacts on the effectiveness of the intervention. In the CAPRISA 004 trial for example, the effectiveness of tenofovir gel increased to 54% when women used the gel according to the dosing strategy in more than 80% of all sexual encounters but was only 28% when the gel was used less than 50% of the time [16]. A case-control study of the iPrEX trial showed that effectiveness was increased to 90% [95% confidence interval (95% CI) 71–98, $P < 0.001$] in those with a detectable drug [44].

In addition to impacting on the effectiveness, suboptimal adherence could also lead to increased resistance. Thus far, the concerns about resistance have not been demonstrated empirically, but are likely to occur in programmatic scale-up. An additional concern relates to therapeutic success rates in patients with breakthrough infections following PrEP. Data from seroconverters in CAPRISA 004 demonstrate no impact of topical antiretrovirals used as PrEP on disease progression [45] or therapeutic success rates. A separate concern about resistance is the use of the same drugs (e.g. tenofovir) in therapy and prevention and some consideration about setting aside a class (or classes) of antiretrovirals for use in prevention only is warranted.

A wider array of HIV prevention options and ones that overcome the adherence challenges are required. Long-acting, slow release products that are less dependent on user compliance compared with oral or gel formulations are already in development. Examples of such products include the monthly intravaginal ring containing the antiretroviral drug,

dapivirine, and the long-acting injectable antiretroviral agents such as rilpivirine and cabotegravir, which are administered every 2–3 months. The monthly dapivirine ring studies underway by MTN and IPM are in advanced stages of clinical trial testing and offer another potential HIV prevention option for women who are unable to use products that require daily or intermittent dosing. Results from these studies are anticipated in the second half of 2015.

THE POTENTIAL OF BROADLY NEUTRALIZING MABS FOR HIV PREVENTION

The use of mAbs to treat human diseases, particularly cancer, is a rapid expanding therapeutic strategy with substantial new commercial potential. To date, only one mAb, Palivizumab (trade name, Synagis), has been licensed for an infectious disease and is used to prevent lung disease caused by respiratory syncytial virus. Given the difficulties in designing an HIV vaccine able to stimulate protective antibodies, the mass-production of mAbs opens up the possibility of testing them as passive immunotherapy. mAbs are also being tested as a drug for treating HIV infection.

Large numbers of highly potent and broadly cross-reactive neutralizing antibodies (bNAbs) have been isolated from several HIV-infected donors over the past 5 years [46²²]. Collectively, these mAbs target five to six conserved neutralization-sensitive epitopes on the HIV-1 envelope including sites on gp120, the CD4⁺ binding site (CD4bs), glycans on the V1, V2 and V3 loops, the membrane proximal external region (MPER) of gp41 and epitopes that span gp120 and gp41 [47²³]. Those that target V3 and V1V2 glycans show the highest potency (IC₅₀ of 0.01–1 µg/ml) while those targeting the MPER and CD4bs are generally broader (neutralize >90% of global isolates).

Several challenge studies have shown that broadly neutralizing mAbs can both protect rhesus macaques from simian immunodeficiency virus (SIV)–HIV chimera (SHIV) infections as well as treat, and possibly cure, macaques with SHIV infections. Studies using the SHIV model in macaques have shown that these mAbs can provide sterilizing immunity against both high-dose and repeated low-dose challenge [48,49]. In particular, PGT121, an antibody that targets glycans in and around the V3 region, demonstrated protection from a single high-dose vaginal challenge with SHIV SF162P3 [50]. All animals that received 1 mg/kg and more than half that received 0.2 mg/kg were protected, consistent with the high potency of PGT121 [50].

Other studies have highlighted the potential for using bNAbs in passive immunization strategies, with immunotherapy in some cases resulting in durable viral control [51²⁴,52²⁵]. These findings provide a strong rationale for human passive protection and therapeutic trials using mAbs and a number of such studies are planned. Several of the mAbs are currently being cloned and produced for preclinical and early clinical testing.

WHY MABS COULD BE A SUITABLE HIV PREVENTION OPTION FOR WOMEN

If mAbs are shown to be well tolerated and efficacious in humans, they could be long-acting HIV-specific prevention strategies as an alternative to current antiretroviral strategies that require daily or intermittent administration. As mAbs may provide protection for a few months after a single administration, this could be a more effective option, as it will not depend on adherence to the same extent as daily pill-taking or coital gel applications. This strategy would be cost effective and patients would only need to visit the clinic every few months to receive their treatment/prophylaxis. Another advantage that a long-acting antibody product offers is that its administration would be unrelated to sex. Further, an mAb could potentially be formulated for systemic or topical application. A topical formulation would produce a product with fewer side effects and could result in much higher levels of mAbs at the site of infection. An injectable product would potentially be administered less frequently but would require a trained healthcare worker to administer the product. Intramuscular injection of mAbs leads to detectable antibodies in the genital tract at a level sufficient in monkeys to protect even against high-dose genital challenge.

A further advantage of mAbs for HIV prevention is that regardless of antiretroviral drugs being used for treatment, these mAbs could be effective against all viruses, including antiretroviral-resistant viruses. To reduce the potential development of resistance to mAbs, combinations of antibodies would be preferable and different antibodies could be used for prevention and treatment.

CONCLUSION

Given the high burden of HIV infection among young women in sub-Saharan Africa and the limited prevention options available, the development of novel HIV prevention technologies such as mAbs, which could be administered as a once-monthly topical application or a three-monthly injection,

could be a game changer in the HIV epidemic in Africa.

Acknowledgements

None.

Financial support and sponsorship

All authors are employees of the Centre for the AIDS Programme of Research in South Africa, which was created by the U.S. National Institutes for Health's Comprehensive International Program of Research on AIDS (CIPRA grant #AI51794).

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. UNAIDS. The gap report. http://www.unaids.org/en/media/unaids/contentassets/documents/unaidspublication/2014/UNAIDS_Gap_report_en.pdf [Accessed 16 October 2014]. Geneva, Switzerland: Joint United Nations Programme on HIV/AIDS (UNAIDS); 2014.
2. UNAIDS. UNAIDS report on the global AIDS Epidemic 2010. <http://www.unaids.org/globalreport/> [Accessed 11 February 2014]. Geneva, Switzerland: Joint United Nations Programme on HIV/AIDS; 2010.
3. Abdool Karim Q, Abdool Karim SS, Singh B, et al. Seroprevalence of HIV infection in rural South Africa. *AIDS* 1992; 6:1535–1539.
4. Shisana O, Rehle T, Simbayi LC, et al. South African National HIV Prevalence, Incidence, Behaviour and Communication Survey 2008: a turning tide among teenagers?. Cape Town: HSRC Press; 2009.
5. Abdool Karim S, Abdool Karim Q. The evolving HIV epidemic in South Africa. *Int J Epidemiol* 2002; 31:37–40.
6. Shisana O, Rehle T, Simbayi LC, et al. South African National HIV Prevalence, Incidence, and Behaviour Survey 2012. Cape Town: HSRC Press; 2014.
7. Department of Health. The 2012 National Antenatal Sentinel HIV & Herpes Simplex Type-2 Prevalence Survey in South Africa. Pretoria: Department of Health; 2014.
8. Ramjee G, Kapiga S, Weiss S, et al. The value of site preparedness studies for future implementation of phase 2/III HIV prevention trials: experience from the HPTN 055 study. *J Acquir Immune Defic Syndr* 2008; 47:93–100.
9. Nel A, Louw C, Hellstrom E, et al. HIV prevalence and incidence among sexually active females in two districts of South Africa to determine microbicide trial feasibility. *PLoS One* 2011; 6:e21528.
10. Nel A, Mabude Z, Smit J, et al. HIV incidence remains high in KwaZulu-Natal, South Africa: evidence from three districts. *PLoS One* 2012; 7:e35278.
11. Feldblum PJ, Latka MH, Lombaard J, et al. HIV incidence and prevalence among cohorts of women with higher risk behaviour in Bloemfontein and Rustenburg, South Africa: a prospective study. *BMJ open* 2012; 2:e000626.
12. Ramjee G, Wand H, Whitaker C, et al. HIV incidence among nonpregnant women living in selected rural, semi-rural and urban areas in KwaZulu-Natal, South Africa. *AIDS Behav* 2012; 16:2062–2071.
13. Abdool Karim Q, Kharsany AB, Frohlich JA, et al. Stabilizing HIV prevalence masks high HIV incidence rates amongst rural and urban women in KwaZulu-Natal, South Africa. *Int J Epidemiol* 2011; 40:922–930.
14. Abdool Karim Q, Kharsany AB, Frohlich JA, et al. HIV incidence in young girls in KwaZulu-Natal, South Africa – public health imperative for their inclusion in HIV biomedical intervention trials. *AIDS Behav* 2012; 16:1870–1876.
15. Abdool Karim SS, Richardson B, Ramjee G, et al. Safety and effectiveness of BufferGel and 0.5% PRO2000 gel for the prevention of HIV infection in women. *AIDS* 2010; 25:957–966.
16. Abdool Karim Q, Abdool Karim SS, Frohlich JA, et al. Effectiveness and safety of tenofovir gel, an antiretroviral microbicide, for the prevention of HIV infection in women. *Science* 2010; 329:1168–1174.
17. Centers for Diseases Control and Prevention. HIV transmission risk. 2013 [cited 9 February 2015]. <http://www.cdc.gov/hiv/policies/law/risk.html>
18. Jewkes RK, Dunkle K, Nduna M, Shai N. Intimate partner violence, relationship power inequity, and incidence of HIV infection in young women in South Africa: a cohort study. *Lancet* 2010; 376:41–48.
19. Pettifor A, O'Brien K, Macphail C, et al. Early coital debut and associated HIV risk factors among young women and men in South Africa. *Int Perspect Sex Reprod Health* 2009; 35:82–90.
20. Wald A, Link K. Risk of human immunodeficiency virus infection in herpes simplex virus type 2-seropositive persons: a meta-analysis. *J Infect Dis* 2002; 185:45–52.
21. Van de Perre P, Segondy M, Foulongne V, et al. Herpes simplex virus and HIV-1: deciphering viral synergy. *Lancet Infect Dis* 2008; 8:490–497.
22. Glynn JR, Biraro S, Weiss HA. Herpes simplex virus type 2: a key role in HIV incidence. *AIDS* 2009; 23:1595–1598.
23. Freeman EE, Weiss HA, Glynn JR, et al. Herpes simplex virus 2 infection increases HIV acquisition in men and women: systematic review and meta-analysis of longitudinal studies. *AIDS* 2006; 20:73–83.
24. Hallfors D, Cho H, Rusakaniko S, et al. Supporting adolescent orphan girls to stay in school as HIV risk prevention: evidence from a randomized controlled trial in Zimbabwe. *Am J Public Health* 2011; 101:1082–1088.
25. Heffron R, Donnell D, Rees H, et al. Use of hormonal contraceptives and risk of HIV-1 transmission: a prospective cohort study. *Lancet Infect Dis* 2012; 12:19–26.
26. Gregson S, Nyamukapa CA, Garnett GP, et al. Sexual mixing patterns and sex-differentials in teenage exposure to HIV infection in rural Zimbabwe. *Lancet* 2002; 359:1896–1903.
27. Kelly RJ, Gray RH, Sewankambo NK, et al. Age differences in sexual partners and risk of HIV-1 infection in rural Uganda. *JAIDS* 2003; 32:446–451.
28. MacPhail C, Williams BG, Campbell C. Relative risk of HIV infection among young men and women in a South African township. *Int J STD AIDS* 2002; 13:331–342.
29. Pettifor AE, Rees HV, Kleinschmidt I, et al. Young people's sexual health in South Africa: HIV prevalence and sexual behaviors from a nationally representative household survey. *AIDS* 2005; 19:1525–1534.
30. Houlihan CF, Larke NL, Watson-Jones D, et al. HPV infection and increased risk of HIV acquisition. A systematic review and meta-analysis. *AIDS* 2012; 26:2211–2222.
31. Abdool Karim Q. Barriers to preventing human immunodeficiency virus in women: experiences from KwaZulu-Natal, South Africa. *J Am Med Womens Assoc* 2001; 56:193–196.
32. Siegfried N, Muller M, Deeks JJ, Volmink J. Male circumcision for prevention of heterosexual acquisition of HIV in men. *Cochrane Database Syst Rev* 2009; CD003362.
33. Weiss HA, Hankins CA, Dickson K. Male circumcision and risk of HIV infection in women: a systematic review and meta-analysis. *Lancet Infect Dis* 2009; 9:669–677.
34. Abdool Karim SS, Kashuba A, Werner L, Abdool Karim Q. Drug concentrations following topical and oral antiretroviral preexposure prophylaxis: implications for HIV prevention in women. *Lancet* 2011; 378:279–281.
35. CONRAD. FACTS 001: safety and effectiveness of tenofovir gel in the prevention of human immunodeficiency virus (HIV-1) infection in young women and the effects of tenofovir gel on the incidence of herpes simplex virus (HSV-2) infection. <http://clinicaltrials.gov/ct2/show/NCT01386294> [Accessed 20 September 2013].
36. Baeten JM, Donnell D, Ndase P, et al. Antiretroviral prophylaxis for HIV prevention in heterosexual men and women. *N Engl J Med* 2012; 367:399–410.
37. Grant RM, Lama JR, Anderson PL, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med* 2010; 363:2587–2599.
38. Thigpen MC, Kebaabetswe PM, Paxton LA, et al. Antiretroviral preexposure prophylaxis for heterosexual HIV transmission in Botswana. *N Engl J Med* 2012; 367:423–434.
39. Abdool Karim SS. HIV preexposure prophylaxis in injecting drug users. *Lancet* 2013; 381:2060–2062.
40. Choopanya K, Martin M, Suntharasamai P, et al. Antiretroviral prophylaxis for HIV infection in injecting drug users in Bangkok, Thailand (the Bangkok Tenofovir Study): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet* 2013; 381:2083–2090.

This study provides evidence for PreP in a population of PWID.

41. U.S. Food and Drug Administration. FDA approves first drug for reducing the risk of sexually acquired HIV infection. 2012 [cited 9 February 2015]. <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm312210.htm>
42. Van Damme L, Corneli A, Ahmed K, et al. The FEM-PreP Trial of Emtricitabine/Tenofovir Disoproxil Fumarate (Truvada) among African Women [Abstract 32LB]. Conference of Retroviruses and Opportunistic Infections; 5–8 March 2012; Seattle, WA.
43. Marrazzo JM, Ramjee G, Richardson BA, et al., for the VOICE Study Team. Tenofovir-based preexposure prophylaxis for HIV infection among African women. *New Engl J Med* 2015; 372:509–518.
44. Anderson PL, Lama JR, Buchbinder S, et al. Expanded case-control analysis of drug detection in the global iPrEx trial [abstractMOLBPE034]. 6th IAS Conference on HIV Pathogenesis, Treatment and Prevention; 17–20 July 2011; Rome, Italy.
45. Garrett NJ, Werner L, Naicker N, et al. HIV disease progression in seroconvertors from the CAPRISA 004 Tenofovir Gel Preexposure Prophylaxis Trial. *J AIDS* 2015; 68:55–61.

This study shows that prior exposure to tenofovir gel has no impact on postinfection CD4⁺ cell counts or the rate of CD4⁺ cell count decline.

46. Klein F, Mouquet H, Dosenovic P, *et al.* Antibodies in HIV-1 vaccine development and therapy. *Science* 2013; 341:1199–1204.

This study explores passive immunotherapy in animal models and suggests that bNAbs may become a valuable addition to the armamentarium of drugs that work against HIV-1.

47. Falkowska E, Le KM, Ramos A, *et al.* Broadly neutralizing HIV antibodies define a glycan-dependent epitope on the prefusion conformation of gp41 on cleaved envelope trimers. *Immunity* 2014; 40:657–668.

This study describes a set of human mAbs that recognize a glycan-dependent epitope on the prefusion conformation of gp41.

48. Mascola JR, Lewis MG, Stiegler G, *et al.* Protection of Macaques against pathogenic simian/human immunodeficiency virus 89.6PD by passive transfer of neutralizing antibodies. *J Virol* 1999; 73:4009–4018.

49. Parren PW, Marx PA, Hessel AJ, *et al.* Antibody protects macaques against vaginal challenge with a pathogenic R5 simian/human immunodeficiency virus at serum levels giving complete neutralization in vitro. *J Virol* 2001; 75:8340–8347.

50. Moldt B, Rakasz EG, Schultz N, *et al.* Highly potent HIV-specific antibody neutralization in vitro translates into effective protection against mucosal SHIV challenge in vivo. *Proc Natl Acad Sci U S A* 2012; 109:18921–18925.

51. Shingai M, Nishimura Y, Klein F, *et al.* Antibody-mediated immunotherapy of macaques chronically infected with SHIV suppresses viraemia. *Nature* 2013; 503:277–280.

This study shows that immunotherapy or a combination of immunotherapy as well as conventional antiretroviral drugs might be useful as a treatment for chronically HIV-1 infected individuals experiencing immune dysfunction.

52. Barouch DH, Whitney JB, Moldt B, *et al.* Therapeutic efficacy of potent neutralizing HIV-1-specific monoclonal antibodies in SHIV-infected rhesus monkeys. *Nature* 2013; 503:224–228.

This study demonstrated a profound therapeutic effect of potent neutralizing HIV-1 specific mAbs in SHIV-infected rhesus monkeys as well as an impact on host immune responses.