

# FEASABILITY OF FOCUSED PARATHYROIDECTOMY IN DEVELOPING COUNTRIES – A Scoping Review

By

KAPIL RUGNATH

Submitted in partial fulfillment of the academic requirements

for the degree of MMed

in the Department of General Surgery

School of Clinical Medicine

College of Health Sciences

University of KwaZulu-Natal

Durban

2023

As the candidate's supervisor I have approved this thesis for submission.

Signed: \_\_\_\_\_  \_\_\_\_\_ Name: S Mewa Kinoo Date: 09 August 2023

## Declaration

I.....Kapil Rugnath.....declare that

(i) The research reported in this dissertation, except where otherwise indicated, is my original work. (ii) This dissertation has not been submitted for any degree or examination at any other university.

(iii) This dissertation does not contain other persons' data, pictures, graphs or other information, unless specifically acknowledged as being sourced from other persons.

(iv) This dissertation does not contain other persons' writing, unless specifically acknowledged as being sourced from other researchers. Where other written sources have been quoted, then:

a) their words have been re-written but the general information attributed to them has been referenced;

b) where their exact words have been used, their writing has been placed inside quotation marks, and referenced.

(v) Where I have reproduced a publication of which I am an author, co-author or editor, I have indicated in detail which part of the publication was actually written by myself alone and have fully referenced such publications.

(vi) This dissertation does not contain text, graphics or tables copied and pasted from the Internet, unless specifically acknowledged, and the source being detailed in the dissertation and in the References sections.

Signed: \_\_\_\_\_  


Date: \_\_\_\_\_

## Overview of The Thesis

Hyperparathyroidism (HPT) is characterized by pathologically excessive parathyroid hormone production. This may be as a consequence of pathology within the parathyroid gland (primary HPT) or as a sequela of pathology outside the parathyroid gland (secondary & tertiary HPT). The gold standard surgical management of HPT was bilateral neck exploration (BNE) but with improvement in localising methods, focused parathyroidectomy (FP) has made the gold standard contentious.

BNE, entailing a large neck incision and 4 gland exploration, demonstrates excellent cure rates of 95%. FP, whereby a single targeted miniature incision is made over the offending gland, is possible in the category of primary HPT as in up to 80% of cases the aetiology is a single gland adenoma. The prerequisites to embarking on this minimally invasive operative technique are accurate pre-operative localisation of the exact site of the offending gland adenoma and assistance of intra-operative adjuncts to confirm successful excision.

Considering the prerequisites required for FP, the initial concern of cost arose. Additionally, concerns of potential inferior success rates were initially entertained. Subsequently, numerous studies revealed this technique to be superior in terms of operative time; cost; convalescence; cosmesis; success rates and the ability to be performed under local anaesthetic, as a day case. As such, the current standard of care for primary HPT where a single gland adenoma is localised, is the FP.

The majority of the studies and trials demonstrating the superiority of FP emanate from developed high-income countries where there is a relative abundance of resources, and as such, the questioning of applicability or relevance of these studies to developing middle-income countries is valid. With the natural hesitancy to adopt these recommendations uncritically in developing countries, where the lack of resources might impede this technique, this concern of applicability, and therefore feasibility, needed to be addressed.

To address this, a scoping review of the literature was conducted, looking specifically at FP, pre-operative localisation and intra-operative adjuncts, in developing countries (upper and lower-middle income). The aim of this review was to ascertain if FP is feasible in these countries by assessing the availability and accuracy of pre-operative localisation, success of FP judged by cure rates and the availability and utility of intra-operative adjuncts.

This review will benefit surgeons in developing countries by demonstrating that FP is not an esoteric procedure described in ivory towers to be entertained in reverie only. Dispelling the myth of non-applicability due to resource constraints, by establishing the technique to be feasible in developing countries, more local surgeons can now, for localised single parathyroid gland adenomas, diverge away from knee-jerk routine bilateral neck explorations. Reassured with support of relevant local literature, they may now confidently embark on learning and performing the technique of FP resulting in satisfied patients who enjoy the superior outcomes associated with this technique.

## **Ethics Declaration**

This study has been approved by the University of Kwa-Zulu Natal Biomedical Research Ethics Committee (BREC).

BREC reference: BREC/00005172/2023

## **Dedication**

I would like to dedicate this dissertation to my parents, Dr Robin Rugnath and Mrs Ravina Rugnath. Words truly cannot express enough the immense gratitude I have for all that you have done for me.

## **Acknowledgements**

I would like to acknowledge my partner, Dr Kaileigh Veeran, for the unwavering support and love given to me throughout this journey. I would not be who I am today and would not have achieved even a fraction of what I have without you.

I would additionally like to acknowledge and thank my supervisor Mr Mewa-Kinoo for the constant support and assistance with reviewing all articles contained within library database used for this study.

## Table of Contents

<b>Declaration .....</b>	<b>2</b>
<b>Overview Of The Thesis .....</b>	<b>3</b>
<b>Ethics Declaration.....</b>	<b>4</b>
<b>Dedication .....</b>	<b>5</b>
<b>Acknowledgements .....</b>	<b>6</b>
<b>Table Of Contents.....</b>	<b>7</b>
<b>Part 1: The Review Of Literature .....</b>	<b>8</b>
<b>Part 2 Submission Ready Manuscript .....</b>	<b>26</b>
<b>Appendices.....</b>	<b>43</b>
<b>APPENDIX 1: THE FINAL STUDY PROTOCOL.....</b>	<b>43</b>
<b>APPENDIX 2: GUIDELINES FOR AUTHORSHIP FOR THE JOURNAL SELECTED FOR SUBMISSION OF THE MANUSCRIPT .....</b>	<b>57</b>
<b>APPENDIX 3: ETHICAL APPROVALS.....</b>	<b>62</b>
<b>APPENDIX 4: DATA COLLECTION TOOLS .....</b>	<b>63</b>
<b>APPENDIX 5: RAW DATA – ARTICLES ENCOUNTERED UPON SEARCH OF THE LITERATURE .....</b>	<b>63</b>

## Part 1: The Review of Literature

### Anatomy of Parathyroid Glands

The parathyroid glands (PG), originally described in 1852 by Richard Owen, and subsequently described in humans for the first time in 1880 by Ivar Sandstrom, are endocrine glands located in the neck, posterior to the thyroid gland (1,2). Usually, there are 4 glands - 2 superior and 2 inferior, however they can vary in location (intrathyroidal or ectopic) and number (reduced in number or supernumerary) due to embryological variations. Intra-thyroidal PG's are relatively rare, with a reported rate of 2%, while ectopic glands, which may be located anywhere along a line from the angle of the mandible superiorly, down to the mediastinum inferiorly, vary in prevalence, with rates ranging between 2% - 43% (3,4). Variability in number, both reduction and additional supernumerary glands, are seen in about 19% of cases (5).

### Physiology of Parathyroid glands

The PG are responsible for maintaining calcium equilibrium within the body. This is achieved through the formation and release of parathyroid hormone (PTH), which maintains eucalcaemia by responding to a series of negative feedback loops. When serum calcium levels are low, via a negative feedback loop, PTH is released from the PG which triggers: resorption of calcium from bone; resorption of calcium, in exchange for phosphate, at the kidneys and release of  $\alpha$ -1-hydroxylase from the kidneys, which activates vitamin D causing an increase in calcium absorption from the gastrointestinal tract. Additionally, PTH release is stimulated by elevated persists unabated and is futile as the kidneys have reduced functional capacity in CKD, and as such, are unable to appropriately respond to PTH. As a result, all 4 PG are culprits in the cause of HPT in this scenario, as they each respond continuously to the inappropriate levels of calcium and phosphate. Treatment may involve pharmacotherapeutics such as calcimimetics, or operative management by parathyroidectomy (7).

Tertiary HPT represents a disease progression from secondary HPT, whereby the PG undergo hyperplasia from incessant stimulation, resulting in autonomous function. Even after the CKD may be completely addressed with a renal transplant, HPT in this scenario continues unrelenting, as a consequence of the now autonomous PG's. Similarly, the culprits here are all 4 PG, that have each undergone hyperplasia. Treatment involves parathyroidectomy (7).

### Operative Options for Management For Parathyroid Gland Pathology

The operative management of HPT depends on the type of HPT, with secondary and tertiary HPT always requiring a 4-gland exploration and excision as the standard of care, known as a bilateral neck exploration (BNE). The operative management of primary HPT is contentious, as the aetiology in around 80% of cases is due only to a single gland adenoma, allowing for a more focused approach and thus obviating the need for 4 gland BNE. However, in cases of primary HPT where multi-gland adenomas are identified, a 4-gland BNE may be needed (7).

### The History of Parathyroidectomy Approaches

The discovery of the PG in humans by *Ivor Sandstrom (1880)* was followed by nearly 3 decades of uncertainty regarding their function, with reports of inexplicable tetany following thyroidectomy, which was averted when these glands were preserved (2,9). *Professor*

*Halsted*, who is generally regarded as the most innovative and influential surgeon from the United States of America, with an astounding number of contributions to surgery, in 1907 reinforced the importance of the PG and their preservation when he stated, “*the fact that these little organs perform some highly important function is sufficient reason for the endeavor to preserve all of them*” (10,11). In 1909, *MacCallum et al* described hypocalcaemia as the cause of tetany in these patients after the PG were excised, and discovered the utility of calcium injections in reversing the condition (12). In the subsequent years, a clearer comprehensible understanding of the PG physiology and pathophysiology followed, culminating in the first successful parathyroidectomy by *Felix Mandl* in 1925 (13).

As a result of the prolific work across multiple years by *Cope et al* and *Churchill et al*, HPT was divided into 3 distinct classifications with the discovery of adenomatous or carcinomatous aetiology in cases of primary disease and diffuse hyperplasia in cases of secondary and tertiary disease (14-21). Additionally, the former was seen mostly in a single gland and rarely in 2 glands, with the latter, diffuse hyperplasia, seen in all 4 glands. These findings lead to *Cope*'s suggestion of at least 2 gland exploration for cases of primary HPT and 4 gland exploration for secondary and tertiary HPT (14). This formed the basis for the myriad of evolutionary changes parathyroidectomy for primary HPT has undergone since the first parathyroidectomy 98 years ago.

Parathyroidectomy by cervical exploration and 4 gland exploration (BNE) was the unchallenged standard of care for the half century that followed *Mandl*'s first successful procedure, with numerous publications released in that period supporting its effectiveness for all aetiology of HPT.

levels of phosphate, via a negative feedback loop, causing renal excretion of phosphate with retention of calcium in exchange, resulting from PTH's stimulatory effect on the kidneys (6).

### Pathology of Parathyroid Glands

Hyperparathyroidism (HPT) is a pathological state in which excessive amounts of PTH is produced, with varying aetiologies, resulting in excessive levels of calcium termed hypercalcaemia. It may be broadly classified into primary, secondary and tertiary, with the majority of patients overall mostly asymptomatic (70% - 80% of cases) (7). The remaining cases of symptomatic presentation usually involve complaints grouped as ‘stones, bones, groans and psychic moans’, derived from renal stones, osteoclastic bone pain and/or fractures, abdominal moans from constipation or peptic ulceration pain and psychic groans from hypercalcaemic induced psychoses/depression (8).

Primary HPT is the most common form of this pathological state, and is the 3<sup>rd</sup> most common endocrine disorder globally (7). The aetiology is most often (80% of cases) a single gland adenoma, whilst the remaining causes include: multi-gland adenoma, parathyroid hyperplasia and parathyroid carcinoma. Primary HPT is managed operatively when the patient is symptomatic or asymptomatic and meets the international workshop of HPT criteria.

Secondary HPT most often occurs in patients with chronic kidney disease (CKD), where there is inadequate resorption of calcium and inadequate excretion of phosphate by the poorly functioning kidneys. This results in reduced calcium levels (hypocalcaemia) and elevated phosphate level (hyperphosphataemia), which stimulates the PG to produce PTH, in an effort to both raise the calcium levels and reduce the phosphate levels to normal. This process

*Norris et al (1947)* described their experience with 322 cases of parathyroid adenoma with successful operative management by a standard cervical exploration and 4 gland assessment (BNE) (22).

*Black et al* described their experience with surgical management of HPT in 63 patients in 1948, in 112 patients in 1953 and in 207 patients in 1956 (23-25). The authors recommended parathyroidectomy by way of cervical exploration with 4 gland assessment (BNE) as a standard approach in keeping with descriptions from colleagues. In the event that a 4<sup>th</sup> gland is not identified, the authors advised a complete cervical exploration including all known possible ectopic locations and thyroid incisions to identify intra-thyroidal PG. A full mediastinal exploration was only advised for follow up re-operative procedures.

*Rienhoff et al (1950)* reported experience with 27 cases of hyperparathyroidism with a specific focus on the surgical management (26). His approach mirrors the previous authors with a recommendation of a primary systematic approach by cervical exploration and 4 gland assessment (BNE), with continued search of the superior mediastinum via the cervical incision and a secondary approach via sternotomy for a full mediastinal exploration, in the event of an elusive offending gland. Interestingly the authors had stated, after an extensive literature search, that only a mere 597 total cases of hyperparathyroidism had been reported globally, since 1903 until circa 1950 (26).

*McGeown et al (1959)* recommended 4 gland exploration via the standard cervical exploration approach (BNE), coupled with a superior mediastinal exploration if an offending gland was not identified (27). A full mediastinal exploration via sternotomy was only advised as a re-operative strategy if no offending gland was identified upon initial exploration and the patient remained hyperparathyroid.

### The Search for A Preoperative Localisation Study

Despite its effectiveness, cases requiring re-operation were still encountered, with the offending ectopic gland only discovered upon extensive search in known other possible ectopic locations (28). As a result, an endeavor to correctly identify these elusive glands prior to embarking on operative therapy was made, marking the birth of preoperative localisation. Originally, cine-oesophagography, selective venous sampling and arteriography were used from 1950 onwards with mostly poor results (29).

*Doppman et al (1969)* noted the inadequate and varied localisation rates with arteriography (0% to 66%) to be due to staining of both parathyroid tissue and overlying thyroid tissue as well as poor anatomic localisation in cases with no staining of any tissues (30). With some modification to the standard technique by combining selective inferior thyroid artery injections, anteroposterior and oblique projections, prolonged filming and routine subtraction, the best results achieved were localisations accurate in only 4 out of 9 patients.

Pre-operative <sup>75</sup>Se-selenomethionine scanning and preoperative injections of toluidine blue was assessed subsequently, with unimpressive results in the former and adverse cardiac effects in the latter (31,32). The use of ultrasonography (US) and computed tomography (CT) in preoperative localisation was subsequently described in the early 1970s, however in keeping with prior modalities, the sensitivities were poor (33,34).

The first rung on the evolutionary ladder of parathyroidectomy was described by *Roth et al* in 1975 and involved a unilateral neck exploration, whereby a single side of the neck was explored facilitating assessment of the 2 ipsilateral PG only (35). This technique was based on the rationale that in 80% of cases the cause of HPT is a single gland adenoma. The decision of the side to be explored was determined by either palpation of the neck, an oesophagogram, vascular sampling (venography or angiography) results or a guess.

This conservative parathyroidectomy revolution was supported by *Edis et al (1977)* and improved upon by *Tibblin et al (1982)* (36,37). The improvement entailed a modification of the conservative unilateral approach to involve an intra-operative saline float test & oil-red-O staining to discern between parathyroid adenoma and hyperplasia, the latter of which would then mandate further contra-lateral exploration. Although excellent cure rates were realised in the group with unilateral exploration, unfortunately far more patients who were intended for this technique upfront were subject to contra-lateral exploration, negating the intended benefit of a more conservative, strictly unilateral exploration. Therefore, a solution to address the difficulty in identifying pathological gland site/s was desperately needed.

The answer came in the form of drastic preoperative localisation advancements. Firstly, improvements in US technology and resultant superior accuracy rates were realised, as demonstrated by *Sample et al (1978)* and reaffirmed by numerous subsequent publications (38).

*Edis et al (1979)* demonstrated the use of high resolution US for parathyroid adenoma localisation in 10 patients (39). With the use of high resolution US, all parathyroid adenomas were accurately located. Despite the impressive results, the limitation of this study was the patient selection, as only patients with suspected large adenomas were included.

*Duffy et al (1980)* further demonstrated the immense benefit of high resolution US in preoperative localisation (40). The authors were able to correctly identify enlarged PG of greater than 5mm, from both primary and secondary HPT, in 19 of 22 patients translating to a accuracy of 83%.

*Moreau et al (1981) & Simeone et al (1981)* bolstered the effectiveness of high resolution US for preoperative localisation by demonstrating sensitivity rates for parathyroid adenomas of 93% and 87,5% respectively (41,42).

The benefit from these impressive improvements in US technology and accuracy rates were demonstrated by *Vogel et al (1998)* who assessed unilateral neck exploration compared to BNE between 1989 & 1996, with accurate US (90% accuracy) as a preoperative localisation method (43). They demonstrated equivalent cure rates, reduced operative time, reduced morbidity and improved cosmesis with the unilateral neck exploration, compared to BNE.

The second additional imaging advancement in parathyroid preoperative localisation was high resolution CT. This improvement in CT scanning was assessed in 1983 and demonstrated to be accurate, having achieved improved localisation rates of 76%, over the inferior 45% rates previously accomplished with conventional CT (44).

The most important step in parathyroidectomy innovation came with profound improvements in preoperative scintigraphic localisation. Parathyroid scintigraphy entails intravenous administration of a radio-labelled tracer coupled with the use of a gamma camera that reveals

the anatomical location of the gland/structure in which the radio-labelled tracer concentrates. Originally in the early 1980s, Thallium 201 ( $^{201}\text{Tl}$ ) was used as a radiotracer for parathyroid scintigraphy, with poor results due to the rapid uptake and concentration of this tracer by the thyroid, in addition to the intended parathyroid tissue (45).

Additional radiotracers used include Iodine 123 ( $\text{I}^{123}$ ) and Technetium 99m ( $^{99\text{m}}\text{Tc}$ ) Pertechnate with mediocre results due to relatively modest uptake & washout and were demonstrated to be limited by the sizes and location of parathyroid adenomas (46).

The major breakthrough came by *Coakley et al (1989)* who described the Technetium 99m Sestamibi ( $^{99\text{m}}\text{Tc}$  Sesta-methoxyisobutylisonitrile) scan as a new form of preoperative localisation (47). The  $^{99\text{m}}\text{Tc}$  Sestamibi (Sestamibi) showed increased parathyroid and thyroid uptake with slower washout, thus resulting in better delineation of the glands. Additionally, compared to  $^{201}\text{Tl}$ , it was demonstrated that this radiotracer had superior physical characteristics for the gamma camera imaging, with better image quality consequently.

*Taillefer et al (1992)* assessed Sestamibi in 23 patients with HPT, with subsequent cervical exploration revealing adenomas in 21 patients and parathyroid hyperplasia in 2 (48). Upon correlation, Sestamibi had correctly identified 19 of the 21 adenomas, therefore demonstrated an accuracy rate of 90% in preoperative localisation for patients with primary HPT.

*Kwan et al (1993)* compared  $^{201}\text{Tl}$ /pertechnate with Sestamibi. There were 27 cases performed with the  $^{201}\text{Tl}$ /pertechnate and 24 cases performed with Sestamibi (49). The accuracy rates were 77% and 95% respectively, demonstrating the superiority of Sestamibi.

This advancement allowed accurate preoperative localisation such that the single offending gland was identified allowing for a more targeted parathyroidectomy approach. A single miniature incision over the site of the offending gland, performed under a general or local anaesthetic, with deep dissection, followed by identification and excision of the pathological gland was possible, and so minimally invasive parathyroidectomy, also known as focused parathyroidectomy (FP), was born (50). This was revolutionary, as in theory, numerous benefits would be achieved over the standard of care, which was BNE, a well tried and tested method as demonstrated by the following authors.

*Van Heerden et al (1991)* reported their experience at the Mayo Clinic with 379 patients with primary HPT undergoing BNE (51). They demonstrated a cure rate of 99,5% with a morbidity and mortality of each less than 1%, advising a more liberal approach to cervical exploration.

*Uden et al (1992)* described their experience with BNE in 250 patients with primary HPT undergoing BNE (52). They demonstrated an overall cure rate of 98,8%, with no difference in the clinical and metabolic benefits from BNE in either older or younger patient groups.

*Chen et al (1996)* demonstrated significant improvements in cure rate and outcomes from BNE performed by an endocrine surgeon in a high-volume centre (53). They demonstrated a cure rate of 97%, with low morbidity, zero mortality and overall reduction in length of stay.

## The Resistance To Change From The Gold Standard

Some authors, as demonstrated, continued to focus on their work with BNE, and not engage with the newly touted FP, however, there were other authors who were firmly against the notion of straying from the gold standard of BNE to the more attractive FP.

*Delbridge et al (1998)* assessed BNE in 733 patients with primary hyperparathyroidism (54). They achieved a cure rate of 99%, solidifying the success of this procedure. Remarkably, the authors stated the interest in ‘minimal access’ or unilateral parathyroid surgery was passing and misguided. They recommended avoiding these minimally invasive techniques as they believed there would be an inevitable increase in failure rates due to multi-gland disease (MGD), in exchange for a minimal cosmetic benefit. Additionally, the authors advised against routine preoperative localisation.

## The Cost Issue Of FP

Some authors claimed the expense of additional preoperative localisation required to perform FP significantly adds to the overall costs, thus making this technique unfeasible.

*Aarum et al (2007)* compared 23 patients undergoing FP with preoperative localisation against 26 patients undergoing BNE, looking specifically at feasibility in terms of cost and cure (55). Cure rates were similar at 96% and 94% respectively, however the authors demonstrated FP to be 21% more costly due to the requisite preoperative localisation studies

*Slepavicius et al (2008)* in their prospective randomised blinded trial compared 23 patients undergoing BNE to 24 patients undergoing FP (56). The authors revealed both techniques to be safe and effective, with FP showing numerous advantages, however they also demonstrated greater costs associated with FP.

*Baliski et al (2008)* compared the ‘gold standard’ BNE to minimally invasive parathyroidectomy in terms of cost (57). They demonstrated BNE to have the lowest cost associated, with excellent cure rates in return.

Additionally, some authors claimed that FP was associated with inferior cure rates and increased recurrence rates.

*Delbridge et al (2000)* reported on their experience with FP in 50 patients (58). Only 42 patients of the 50 undergoing FP were cured, translating to a cure rate of 84%. Of the remaining 8 patients, 7 required conversion to BNE and 1 was a failed procure. The authors concluded that although FP may have been feasible, there were concerns associated.

*Norman et al (2012)* reported their experience with 15000 parathyroidectomies, comparing BNE with FP (59). They demonstrated FP to be associated with a higher recurrence rate, with a 1 year recurrence rate of 3% – 5% and a 10 year recurrence rate of 4% - 6%.

Eventually, through the next 2 decades, articles subsequently demonstrated FP to be equally effective as BNE in terms of cure, with reduced overall cost, mostly stemming from markedly reduced operative time and hospital stay and reduced morbidity, confirming this technique to be safe and feasible (60).

*Udelsman et al (2000)* revealed the immense benefits of FP for primary HPT (61). They revealed their experience with 100 cases of minimally invasive parathyroidectomy, with a cure rate of 100%, reduced hospital costs and no long term complications.

*Burkey et al (2003)* reported on their experience with 100 parathyroidectomies, 70 cases of FP and 30 cases of BNE (62). Both methods of parathyroidectomy resulted in an overall 100% cure, however FP was demonstrated to be superior to BNE, with reduced operative time, reduced length of hospital stay, reduced morbidity and resulted in rapid convalescence.

*Pang et al (2007)* described their experience with 500 cases of consecutive FP compared to 601 cases of BNE over a 5 year period (63). They demonstrated FP to be safe and effective, with a cure rate of 97,4% with no evidence of any increase in morbidity, compared to BNE (3% vs 3,99%).

*Udelsman et al (2002)* subsequently compared 255 cases of minimally invasive parathyroidectomy to 401 cases of BNE (13). The authors demonstrated equivalent cure rates, with FP associated with reduced morbidity (1,2% vs 3% respectively), 50% reduced operative time, a seven-fold reduction in length of hospital stay and a nearly 50% reduction in cost, compared to BNE.

*Norlen et al (2015)* assessed outcomes of parathyroidectomies in 4569 patients, of which 2531 underwent FP and 2038 underwent BNE (64). The authors demonstrated FP to have an equivalent long term recurrence rate to BNE (0,6% vs 0,4% respectively), with significantly reduced morbidity (3,6% vs 7,6%).

*Udelsman et al (2011)* ultimately described their final cumulative experience with minimally invasive parathyroidectomy (65). Overall they compared 1037 cases of FP to 613 cases of BNE. The authors demonstrated a higher cure rate with FP (99,4%) compared to BNE (97,1%), with reduced morbidity, length of stay and cost, concluding FP to be the superior technique.

*Wong et al (2011)* discussed their experience with 100 parathyroidectomy cases. 93 underwent FP and 7 underwent BNE (conversions from FP) (66). The operative time for FP was significantly lower than for BNE, with an excellent overall cure rate of 96%.

### Sestamibi Scans

The improvements in FP results were mostly due to the marked improvements in Sestamibi. Sestamibi originally made use of planar 2D images, however, further advancements were realised with the advent of the single photon emission computed tomography (SPECT) which, when used in combination with the Sestamibi (Sestamibi SPECT), produced detailed 3D images resulting in improved accuracy rates. The combination of CT to Sestamibi SPECT (Sestamibi SPECT/CT) produced 3D images with anatomical detail and with the addition of intravenous contrast during the CT (termed 4D CT), exquisitely detailed anatomical images were produced (67,68).

*Wong et al (2015)* in their meta-analysis of 24 studies over 10 years assessed Sestamibi, Sestamibi SPECT and Sestamibi SPECT/CT as preoperative localisation modalities with a focus on their sensitivities (69). They demonstrated an incremental improvement in

sensitivities from the original Sestamibi, to Sestamibi SPECT and finally the best sensitivity with Sestamibi SPECT/CT.

*Yeh et al (2019)* assessed 400 patients with 4D CT, Sestamibi SPECT/CT and both modalities combined as forms of preoperative localisation (70). They demonstrated 4D CT to have the highest sensitivity, superior to Sestamibi SPECT/CT. Additionally, the combination of both did not improve the sensitivity.

### Multiple Gland Disease (MGD)

Despite excellent preoperative localisation accuracy and exceptional cure rates realised with FP, the hindrance of MGD and missed double adenomas remained a challenge. The incidence of MGD is estimated to be between 10% - 15% (71).

*Lee et al (2002)* demonstrated their experience with MGD (72). The authors assessed 214 consecutive cases of BNE and compared this to 2166 cases of BNE and 2095 cases of FP reported in the literature. They found a 20,6% incidence of MGD in their cohort, a 19,3% incidence of MGD in the literature from BNE and a 5,3% incidence of MGD reported with FP in the literature. They concluded the concern of missed cases of MGD with FP was evident from the low rate of MGD reported with FP in the literature.

*Gough et al (2006)* assessed 50 patients requiring re-operative parathyroid surgery (73). The authors demonstrated 62% of cases of persistent or recurrent primary HPT requiring re-operation was due to MGD. Additionally, they also revealed the poor sensitivity of preoperative localisation for MGD.

*Siperstein et al (2008)* performed BNE in all patients having had FP in their series and demonstrated missed MGD in 22% of patients with Sestamibi as preoperative localisation, in 22% when US was used and in 20% of cases when combined Sestamibi and US were used as preoperative localisation (74).

The issue stems from the fact that preoperative localisation with US and Sestamibi is accurate for single gland disease, and has shown to be poorly sensitive for multi-gland disease (75). With Sestamibi, it is thought the inaccuracy with identifying the second adenoma in MGD results from the majority of radio-labelled tracer being concentrated in the first adenoma encountered, concealing the second. As a consequence, a FP in one of these patients with an unidentified second adenoma will result in inadequate cure, necessitating re-localisation and re-operation.

*Nichols et al (2012)* assessed 651 patients with primary HPT, 80% with single gland disease and 20% with MGD (76). Sestamibi was demonstrated to be significantly less sensitive for MGD compared to single gland disease (61% vs 97%).

*Milas et al (2003)* reviewed 828 patients with HPT undergoing BNE (77). They discovered MGD in 28% of patients and single gland disease in 71% of patients. Sestamibi correctly identified MGD in only 6% of patients, with the authors concluding FP, as opposed to BNE, may predispose to cases of persistent or recurrent HPT.

## Intraoperative Parathyroid Hormone (IOPTH) Assay

Fortuitously, the quick PTH, also known as the intraoperative parathyroid hormone (IOPTH) assay, was conceived. This is a chemiluminometric assay with rapid turnover in sample processing that bears results within a mere 15 – 20 minutes, allowing for intraoperative use. The utility in the test lies in its predictive power, whereby a 50% or greater drop in IOPTH, relative to baseline PTH, serves as confirmation of successful excision. A drop of less than 50% indicates additional PG pathology such as multi-gland disease, which would then necessitate a BNE to identify the additional elusive offending gland (78).

*Nussbaum et al (1988)* originally described the use of IOPTH, with a modification of the standard PTH testing by immunoradiometric assay (IRMA), producing a result within 15 minutes (79). The authors were able to demonstrate an IOPTH level fall to less than 40% of baseline PTH, from a 15 minute post excision sample, correlated with successful parathyroidectomy.

*Ryan et al (1990)* subsequently described their version of IOPTH, using IRMA with differing incubation methods (80). The result was a test with wider reference range, specifically a lesser lower limit, however with a turnaround time of 1 hour. They demonstrated an IOPTH sampled at 20 minutes post excision correlated to complete excision.

*Irvin et al (1991)* described their experience with IOPTH to assist surgeons with parathyroidectomies in an endeavour to improve cure rates by reducing the incidence of missed(MGD) (81). The authors used the IRMA modified IOPTH with an alteration to the incubation time, allowing a 15 minute turnaround. The issue encountered was the accuracy of the IOPTH being affected by PG manipulation. They discovered elevated level of IOPTH prior to excision, after manipulation alone. They concluded this form of IOPTH should rather be relative to post-manipulation, pre-excision PTH levels, rather than baseline PTH levels, to boost accuracy.

*Irvin et al (1993)* reported on their improved experience with IOPTH (82). The authors further modified the IRMA processing such that the IOPTH turnaround time reduced to 12 minutes. Additionally, accuracy was markedly improved. A 10 minute post excision IOPTH drop of 54% correlated with successful parathyroidectomy, and was associated with an overall accuracy of 97%.

*Irvin et al (1994)* described their up-to-date experience with IOPTH (83). Sampling PTH 5 minutes post excision, and processing the specimen with immunochemiluminometric assay (ICMA) rather than the traditional IRMA resulted in a sensitive test with rapid turnaround time of 10 minutes, which allowed intraoperative use to confirm completeness of resection. The authors demonstrated the sensitivity of this test to predict postoperative calcium levels of 94% and an overall accuracy of 95%.

Over the next few decades, numerous studies were published on IOPTH use for parathyroidectomy, with initial studies revealing marked improvements in cure rates, recommending this procedure be routine with all FP, to some later studies revealing no difference in cure with or without the IOPTH.

*Quinn et al (2021)* in their systematic review and meta-analysis assessed 12 studies describing FP with IOPTH for primary HPT (84). The authors revealed IOPTH to be of great benefit, with increased cure rates, reduced recurrence and need for re-operative surgery, increased conversion from FP to BNE from positive IOPTH results and no statistical increase in operative time. Additionally, the converse was found true. FP without IOPTH resulted in less conversions from FP to BNE, reduced cure rates and increased re-operative surgery for recurrence.

*Ishii et al (2018)* in their systematic review and meta-analysis assessed 14 studies describing FP with IOPTH for primary HPT (85). The authors revealed higher cure rates for FP without the use of IOPTH compared to FP with IOPTH (99,3% vs 98,1%) and lower recurrence rates without IOPTH (0,2% vs 1,5%). The concluding remark was that FP is a safe and effective procedure with concordant preoperative imaging, without the need of IOPTH.

The numerous publications on IOPTH revealed marked variation in sampling and interpretation, giving rise to differing cure rates and so differing opinions on utility. This dilemma required some stricter criteria to create uniformity.

The *Halle Criterion* published recommended a low normal IOPTH level ( $\leq 35$  ng/L) within 15 minutes post excision of the offending PG serving as confirmation of a successful procedure. The *Miami Criterion* recommended a 50% drop in IOPTH relative to preoperative baseline or pre-excision levels, sampled at 10 minutes post excision, serving as confirmation of success. The *Rome Criterion* recommended an IOPTH of greater than 50% drop relative to the highest pre-excision levels and/or IOPTH within the reference range at 20 minutes post excision and/or an IOPTH level of 7.5 ng/L or less than the 10 minutes post excision value. The *Vienna Criterion* recommended an IOPTH level drop of greater than 50% relative to baseline PTH levels. The *Miami Criterion* has demonstrated the highest accuracy and sensitivity rates (86).

The latest National Institute of Clinical Excellence (NICE) guidelines for primary HPT management recommend against routine IOPTH use during primary surgery due to minimal benefit and potential additional cost involved. The latest European Society of Endocrine Surgeons (ESES) guidelines for primary HPT management differ from the NICE guidelines. The ESES recommends IOPTH use for the primary surgery in instances of discordant preoperative localisation or when only single imaging is used as preoperative localisation (87).

#### Minimally Invasive Radio-Guided FP (MIRP)

Other intraoperative adjuncts have been described. *Norman et al (1999)* proposed the minimally invasive radio-guided FP (MIRP) whereby a radioactive tracer that concentrates in parathyroid tissue is injected preoperatively and serves as an intraoperative road map (88). Once in theatre, a handheld gamma probe is used to detect radioactivity which is present in the parathyroid tissue from the radioactive tracer. Once the offending gland is excised, the gamma probe is applied to excised gland and the neck. A confirmatory result of successful excision is an excised gland of 20% or more radioactivity and a background radioactivity of less than 20% in the neck. Any lower reading from the PG or higher neck background radioactivity reading indicates potential residual parathyroid tissue and thus further exploration is warranted. However this method was not reproduced in any other centres.

## Methylene Blue For Localisation

Methylene blue use for rapid intraoperative identification of the PG was originally described by *Dudley et al (1971)* (89). It has been shown by *Patel et al (2012)* in a systematic review of 39 studies that methylene blue is effective for rapid intraoperative PG identification (90). The dye is administered intravenously and subsequently concentrates and stains the PG, allowing for easy identification. However, despite the impressive results reported, the studies included in the systematic review were all observational studies of low quality evidence. There is no randomised controlled trial or other high quality evidence supporting its use. Additionally, the great deterrent with using this technique is the grave adverse effects. Numerous studies have demonstrated severe neurotoxicity associated with this technique (91,92).

*Candell et al (2014)* described their experience with intraoperative US guided methylene blue dye injection into the PG for re-operative parathyroid surgery (93). They demonstrated immense benefit with this technique in assisting PG identification in an already 'hostile' field. Cure was achieved in all but 1 patient with parathyroid carcinoma who developed recurrent hypercalcaemia. The complication rate was low, with 1 case of transient recurrent laryngeal nerve palsy.

## Frozen Section

Frozen section examination refers to the freezing and sectioning of intraoperative specimen with subsequent rapid microscopic pathological examination to provide swift intraoperative feedback to the operating surgeon (94). The benefit is the rapid turnaround time, however this is at the expense of accuracy. Currently its routine use is not recommended for parathyroid surgery. The benefit it confers is identification of parathyroid tissue over other tissue types, however is not able to distinguish between a parathyroid adenoma and parathyroid hyperplasia. Additionally, there have been cases of missed diagnoses due to inaccurate frozen section results. Since the advent of IOPTH, the reliance on frozen section for parathyroid surgery has declined (95,96).

## Newer Intra-Operative Adjuncts

A multitude of additional intraoperative adjuncts to assist in PG identification in the form of optical techniques have been published within the last 5 years. The majority of clinical studies have evaluated autofluorescence and indocyanine green with good results thus far, however more rigorous evidence with randomised controlled trials is warranted to truly validate their benefit (97).

## Minimally Invasive Video Assisted Parathyroidectomy (MIVAP)

As a result of the impressive performance of FP and its subsequent general adoption, much effort has been made to perfect FP even further, with numerous evolutions. Originally, FP, like BNE, has been performed as an 'open' procedure only, involving a skin incision and followed by deep dissection and PG excision. Newer minimally invasive techniques have been applied to FP with the use of an endoscope.

Minimally invasive video assisted parathyroidectomy (MIVAP) first described by *Miccoli et al (1997)* involves a targeted miniature skin incision (via either a central or lateral approach) followed by open dissection with subsequent insertion of an endoscope to facilitate accurate

PG identification and excision (98,99). This technique relies on external traction to create a working room, with no insufflation whatsoever. Results have shown equivalent operative time, cure rates and length of hospital stay to the traditional open technique (100).

*Barczynski et al (2006)* in their randomised blinded trial compared traditional open FP to MIVAP and demonstrated equivalent cure and morbidity, however MIVAP was associated with lower postoperative pain requiring reduced analgesic usage; improved convalescence, and better cosmesis with the disadvantage of increased cost from endoscopic instrument use, when compared to open FP (101).

The purely endoscopic parathyroidectomy (EP) was first described by *Gagner et al (1996)* and entails a skin incision followed by endoscope insertion, insufflation and dissection (102). Two access options exist, central and lateral. The central access technique, performed entirely endoscopically, is best used for anteriorly located PG. For deep, posteriorly located PG, a lateral approach to EP, as described by *Henry et al (1999)*, is better suited (103,104). The overall concern with this technique is the conversion rate. *Fouquet et al (2010)* demonstrated excellent cure rates (98%), however 28% of cases had to be converted to the open procedure due to lack of intraoperative localisation; bleeding; inadequate IOPTH drop and difficult dissection (105). The authors conclude that this may be the reason this technique has not been widely adopted.

### Problem Statement

Parathyroidectomy from primary HPT has undergone substantial evolutionary change over the past 98 years. FP has been demonstrated to be safe, effective and cost friendly and is the current standard of care. However, these recommendations stem from literature emanating from high-income countries

### Research Question

It remains to be demonstrated as to whether FP is safe and cost effective in low and middle-income countries, considering the resource constraints these regions face.

### References

1. Johansson H. The Uppsala anatomist Ivar Sandström and the parathyroid gland. *Ups J Med Sci.* 2015;120(2):72-7.
2. SANDSTROM I. Om en ny Kortl has menniskan och atskilliga daggdjur. *Upsala lakarforening Forh.* 1880;15:441.
3. Policeni BA, Smoker WRK, Reede DL. Anatomy and Embryology of the Thyroid and Parathyroid Glands. *Seminars in Ultrasound, CT and MRI.* 2012;33(2):104-14.
4. Noussios G, Anagnostis P, Natsis K. Ectopic parathyroid glands and their anatomical, clinical and surgical implications. *Exp Clin Endocrinol Diabetes.* 2012;120(10):604-10.
5. Tattera D, Wong LM, Vikse J, Sanna B, Pękala P, Walocha J, et al. The prevalence and anatomy of parathyroid glands: a meta-analysis with implications for parathyroid surgery. *Langenbeck's Archives of Surgery.* 2019;404(1):63-70.
6. Lofrese JJ, Basit H, Lappin SL. *Physiology, Parathyroid: StatPearls Publishing, Treasure Island (FL); 2022 2022.*
7. Fraser WD. Hyperparathyroidism. *The Lancet.* 2009;374(9684):145-58.

8. S Whyte R. In search of stones, bones, abdominal groans and psychic moans: putting the pieces together-a case of parathyroid adenoma. *Perspectives (Pre-2012)*. 2009;33(2):5.
9. Kunstman JW, Udelsman R. Superiority of Minimally Invasive Parathyroidectomy. *Advances in Surgery*. 2012;46(1):171-89.
10. Cameron JL. William Stewart Halsted. Our surgical heritage. *Ann Surg*. 1997;225(5):445-58.
11. Halsted WS, Evans HM. I. The Parathyroid Glandules. Their Blood Supply and their Preservation in Operation upon the Thyroid Gland. *Ann Surg*. 1907;46(4):489-506.
12. Maccallum WG, Voegtlin C. ON THE RELATION OF TETANY TO THE PARATHYROID GLANDS AND TO CALCIUM METABOLISM. *J Exp Med*. 1909;11(1):118-51.
13. Udelsman R. Six hundred fifty-six consecutive explorations for primary hyperparathyroidism. *Ann Surg*. 2002;235(5):665-70; discussion 70-2.
14. Cope O. The study of hyperparathyroidism at the Massachusetts General Hospital. *N Engl J Med*. 1966;274(21):1174-82.
15. Cope O. Hyperparathyroidism: Diagnosis and management. *The American Journal of Surgery*. 1960;99(4):394-403.
16. Cope O, Barnes BA, Castleman B, Mueller GC, Roth SI. Vicissitudes of parathyroid surgery: trials of diagnosis and management in 51 patients with a variety of disorders. *Ann Surg*. 1961;154(4):491-508.
17. Cope O, Keynes WM, Roth SI, Castleman B. Primary chief-cell hyperplasia of the parathyroid glands: a new entity in the surgery of hyperparathyroidism. *Ann Surg*. 1958;148(3):375-88.
18. Cope O, Nardi GL, Castleman B. Carcinoma of the parathyroid glands: 4 cases among 148 patients with hyperparathyroidism. *Ann Surg*. 1953;138(4):661-71.
19. Churchill ED. THE OPERATIVE TREATMENT OF HYPERPARATHYROIDISM. *Ann Surg*. 1934;100(4):606-12.
20. CHURCHILL ED. Principles of parathyroid surgery. *New England Journal of Medicine*. 1937;216(9):376-7.
21. Churchill ED, Cope O. THE SURGICAL TREATMENT OF HYPERPARATHYROIDISM: BASED ON 30 CASES CONFIRMED BY OPERATION. *Ann Surg*. 1936;104(1):9-35.
22. Norris EH. The parathyroid adenoma; a study of 322 cases. *Surg Gynecol Obstet*. 1947;84(1):1-41.
23. Black B. Surgical aspects of hyperparathyroidism. *Surgery, gynecology & obstetrics*. 1948;87(2):172-82.
24. Black BM. *Hyperparathyroidism*: Thomas Springfield, Ill.; 1953.
25. BLACK BM, ZIMMER JF. Hyperparathyroidism, with Particular Reference to Treatment: Review of Two Hundred Seven Proved Cases. *AMA Archives of Surgery*. 1956;72(5):830-7.
26. Rienhoff WF, Jr. The surgical treatment of hyperparathyroidism, with a report of 27 cases. *Ann Surg*. 1950;131(6):917-44.
27. McGeown MG, Morrison E. Hyperparathyroidism. *Postgraduate Medical Journal*. 1959;35(404):330-7.
28. Golden A, Canary JJ, Kerwin DM. Concurrence of hyperplasia and neoplasia of the parathyroid glands. *The American Journal of Medicine*. 1965;38(4):562-78.
29. Brennan MF, Doppman JL, Kurdy AG, Marx SJ, Spiegel AM, Aurbach GD. Assessment of techniques for preoperative parathyroid gland localization in patients undergoing reoperation for hyperparathyroidism. *Surgery*. 1982;91(1):6-11.

30. Doppman JL, Hammond WG, Melson GL, Evens RG, Ketcham AS. Staining of Parathyroid Adenomas by Selective Arteriography. *Radiology*. 1969;92(3):527-30.
31. DiGiulio W, Morales JO. The Value of the Selenomethionine Se 75 Scan in Preoperative Localization of Parathyroid Adenomas. *JAMA*. 1969;209(12):1873-80.
32. Yeager RM, Kremenz ET. Toluidine blue in identification of parathyroid glands at operation. *Ann Surg*. 1969;169(6):829-38.
33. Karo JJ, Maas LC, Kaine H, Gelzayd EA. Ultrasonography and parathyroid adenoma. *Jama*. 1978;239(20):2163-4.
34. Shimshak RR, Schoenrock GJ, Taekman HP, Cianci P, Chambers RF. Case report. Preoperative localization of a parathyroid adenoma using computed tomography and thyroid scanning. *J Comput Assist Tomogr*. 1979;3(1):117-9.
35. Roth SI, Wang C-a, Potts JT. The team approach to primary hyperparathyroidism. *Human Pathology*. 1975;6(6):645-8.
36. Edis AJ, Beahrs OH, van Heerden JA, Akwari OE. "Conservative" versus "liberal" approach to parathyroid neck exploration. *Surgery*. 1977;82(4):466-73.
37. Tibblin S, Bondeson AG, Ljungberg O. Unilateral parathyroidectomy in hyperparathyroidism due to single adenoma. *Annals of surgery*. 1982;195(3):245-52.
38. Sample WF, Mitchell SP, Bledsoe RC. Parathyroid ultrasonography. *Radiology*. 1978;127(2):485-90.
39. Edis AJ, Evans TC, Jr. High-resolution, real-time ultrasonography in the preoperative location of parathyroid tumors. Pilot study. *N Engl J Med*. 1979;301(10):532-4.
40. Duffy P, Picker RH, Duffield S, Reeve T, Hewlett S. Parathyroid sonography: a useful aid to preoperative localization. *J Clin Ultrasound*. 1980;8(2):113-6.
41. Moreau JF, Dubost C, Buy JN, Ferry J. [Pre-operative detection of parathyroid adenomas with ultrasound echography (author's transl)]. *Nouv Presse Med*. 1981;10(23):1923-7.
42. Simeone JF, Mueller PR, Ferrucci JT, Jr., vanSonnenberg E, Wang CA, Hall DA, et al. High-resolution real-time sonography of the parathyroid. *Radiology*. 1981;141(3):745-51.
43. Vogel LM, Lucas R, Czako P. Unilateral parathyroid exploration. *Am Surg*. 1998;64(7):693-6; discussion 6-7.
44. Stark DD, Gooding GA, Moss AA, Clark OH, Ovenfors CO. Parathyroid imaging: comparison of high-resolution CT and high-resolution sonography. *AJR Am J Roentgenol*. 1983;141(4):633-8.
45. MacFarlane SD, Hanelin LG, Taft DA, Ryan JA, Jr., Fredlund PN. Localization of abnormal parathyroid glands using thallium-201. *Am J Surg*. 1984;148(1):7-13.
46. Kiratli PÖ, Peksoy İ, Erbaş B, Gedikoğlu G, Karabulut N. Technetium-99m pertechnetate uptake in ectopic parathyroid adenoma. *Annals of nuclear medicine*. 1999;13:113-5.
47. Coakley AJ, Kettle AG, Wells CP, O'Doherty MJ, Collins RE. 99Tcm sestamibi--a new agent for parathyroid imaging. *Nucl Med Commun*. 1989;10(11):791-4.
48. Taillefer R, Boucher Y, Potvin C, Lambert R. Detection and localization of parathyroid adenomas in patients with hyperparathyroidism using a single radionuclide imaging procedure with technetium-99m-sestamibi (double-phase study). *J Nucl Med*. 1992;33(10):1801-7.
49. Kwan W, Morita E, Clark O. 99M-TECHNETIUM SESTAMIBI IS A SUPERIOR AGENT TO 201-THALLIUM/PERTECHNETATE SUBTRACTION IN EVALUATING PATIENTS FOR REPEAT PARATHYROIDECTOMY FOR EITHER ADENOMA OR HYPERPLASIA. *Clinical Nuclear Medicine*. 1993;18(10):927.
50. Norman J, Chheda H, Farrell C, Britt LG, et al. Minimally Invasive Parathyroidectomy for Primary Hyperparathyroidism: Decreasing Operative Time and

Potential Complications while Improving Cosmetic Results / Discussion. *The American Surgeon*. 1998;64(5):391-5; discussion 5-6.

51. van Heerden JA, Grant CS. Surgical treatment of primary hyperparathyroidism: An institutional perspective. *World Journal of Surgery*. 1991;15(6):688-92.
52. Udén P, Chan A, Duh Q-Y, Siperstein A, Clark OH. Primary hyperparathyroidism in younger and older patients: Symptoms and outcome of surgery. *World Journal of Surgery*. 1992;16(4):791-7.
53. Chen H, Zeiger MA, Gordon TA, Udelsman R. Parathyroidectomy in Maryland: effects of an endocrine center. *Surgery*. 1996;120(6):948-53.
54. Delbridge LW, Younes NA, Guinea AI, Reeve TS, Clifton-Bligh P, Robinson BG. Surgery for primary hyperparathyroidism 1962-1996: indications and outcomes. *Med J Aust*. 1998;168(4):153-6.
55. Aarum S, Nordenström J, Reihner E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for primary hyperparathyroidism: the new versus the old order: a randomised controlled trial of preoperative localisation. *Scandinavian journal of surgery*. 2007;96(1):26-30.
56. Slepavicius A, Beisa V, Janusonis V, Strupas K. Focused versus conventional parathyroidectomy for primary hyperparathyroidism: a prospective, randomized, blinded trial. *Langenbecks Arch Surg*. 2008;393(5):659-66.
57. Baliski C, Nosyk B, Melck A, Bugis S, Rosenberg F, A HA. The cost-effectiveness of three strategies for the surgical treatment of symptomatic primary hyperparathyroidism. *Ann Surg Oncol*. 2008;15(10):2653-60.
58. Delbridge LW, Dolan SJ, Hop TT, Reeve TS, Robinson BG, Wilkinson MR. Minimally invasive parathyroidectomy: 50 consecutive cases. *Medical journal of Australia*. 2000;172(9):418-22.
59. Norman J, Lopez J, Politz D. Abandoning unilateral parathyroidectomy: why we reversed our position after 15,000 parathyroid operations. *J Am Coll Surg*. 2012;214(3):260-9.
60. Bergenfelz A, Lindblom P, Tibblin S, Westerdahl J. Unilateral versus bilateral neck exploration for primary hyperparathyroidism: a prospective randomized controlled trial. *Ann Surg*. 2002;236(5):543-51.
61. Udelsman R, Donovan PI, Sokoll LJ. One hundred consecutive minimally invasive parathyroid explorations. *Ann Surg*. 2000;232(3):331-9.
62. Burkey SH, Snyder WH, III, Nwariaku F, Watumull L, Mathews D. Directed parathyroidectomy: Feasibility and performance in 100 consecutive patients with primary hyperparathyroidism. *Archives of Surgery*. 2003;138(6):604-8; discussion 8-9.
63. Pang T, Stalberg P, Sidhu S, Sywak M, Wilkinson M, Reeve T, et al. Minimally invasive parathyroidectomy using the lateral focused mini-incision technique without intraoperative parathyroid hormone monitoring. *Journal of British Surgery*. 2007;94(3):315-9.
64. Norlén O, Wang KC, Tay YK, Johnson WR, Grodski S, Yeung M, et al. No need to abandon focused parathyroidectomy: a multicenter study of long-term outcome after surgery for primary hyperparathyroidism. *Ann Surg*. 2015;261(5):991-6.
65. Udelsman R, Lin Z, Donovan P. The superiority of minimally invasive parathyroidectomy based on 1650 consecutive patients with primary hyperparathyroidism. *Ann Surg*. 2011;253(3):585-91.
66. Wong W, Foo FJ, Lau M, Sarin A, Kiruparan P. Simplified minimally invasive parathyroidectomy: a series of 100 cases and review of the literature. *The Annals of The Royal College of Surgeons of England*. 2011;93(4):290-3.

67. Eslamy HK, Ziessman HA. Parathyroid Scintigraphy in Patients with Primary Hyperparathyroidism: 99mTc Sestamibi SPECT and SPECT/CT. *RadioGraphics*. 2008;28(5):1461-76.
68. Rodgers SE, Hunter GJ, Hamberg LM, Schellingerhout D, Doherty DB, Ayers GD, et al. Improved preoperative planning for directed parathyroidectomy with 4-dimensional computed tomography. *Surgery*. 2006;140(6):932-41.
69. Wong KK, Fig LM, Gross MD, Dwamena BA. Parathyroid adenoma localization with 99mTc-sestamibi SPECT/CT: a meta-analysis. *Nuclear Medicine Communications*. 2015;36(4):363-75.
70. Yeh R, Tay Y-KD, Tabacco G, Dercle L, Kuo JH, Bandeira L, et al. Diagnostic Performance of 4D CT and Sestamibi SPECT/CT in Localizing Parathyroid Adenomas in Primary Hyperparathyroidism. *Radiology*. 2019;291(2):469-76.
71. Barczyński M, Bränström R, Dionigi G, Mihai R. Sporadic multiple parathyroid gland disease—a consensus report of the European Society of Endocrine Surgeons (ESES). *Langenbeck's Archives of Surgery*. 2015;400(8):887-905.
72. Lee NC, Norton JA. Multiple-Gland Disease in Primary Hyperparathyroidism: A Function of Operative Approach? *Archives of Surgery*. 2002;137(8):896-900.
73. Gough I. Reoperative parathyroid surgery: the importance of ectopic location and multigland disease. *ANZ J Surg*. 2006;76(12):1048-50.
74. Siperstein A, Berber E, Barbosa GF, Tsinberg M, Greene AB, Mitchell J, et al. Predicting the Success of Limited Exploration for Primary Hyperparathyroidism Using Ultrasound, Sestamibi, and Intraoperative Parathyroid Hormone: Analysis of 1158 Cases. *Annals of Surgery*. 2008;248(3):420-8.
75. Chiu B, Sturgeon C, Angelos P. What is the link between nonlocalizing sestamibi scans, multigland disease, and persistent hypercalcemia? A study of 401 consecutive patients undergoing parathyroidectomy. *Surgery*. 2006;140(3):418-22.
76. Nichols KJ, Tomas MB, Tronco GG, Palestro CJ. Sestamibi parathyroid scintigraphy in multigland disease. *Nucl Med Commun*. 2012;33(1):43-50.
77. Milas M, Wagner K, Easley KA, Siperstein A, Weber CJ. Double adenomas revisited: nonuniform distribution favors enlarged superior parathyroids (fourth pouch disease). *Surgery*. 2003;134(6):995-1003; discussion -4.
78. Chen H, Pruhs Z, Starling JR, Mack E. Intraoperative parathyroid hormone testing improves cure rates in patients undergoing minimally invasive parathyroidectomy. *Surgery*. 2005;138(4):583-90.
79. Nussbaum SR, Thompson AR, Hutcheson KA, Gaz RD, Wang CA. Intraoperative measurement of parathyroid hormone in the surgical management of hyperparathyroidism. *Surgery*. 1988;104(6):1121-7.
80. Ryan MF, Jones SR, Barnes AD. Modification to a commercial immunoradiometric assay permitting intraoperative monitoring of parathyroid hormone levels. *Ann Clin Biochem*. 1990;27 ( Pt 1):65-8.
81. Irvin GL, 3rd, Dembrow VD, Prudhomme DL. Operative monitoring of parathyroid gland hyperfunction. *Am J Surg*. 1991;162(4):299-302.
82. Irvin GL, Dembrow VD, Prudhomme DL. Clinical usefulness of an intraoperative "quick parathyroid hormone" assay. *Surgery*. 1993;114(6):1019-23.
83. Irvin GL, 3rd, Deriso GT, 3rd. A new, practical intraoperative parathyroid hormone assay. *Am J Surg*. 1994;168(5):466-8.
84. Quinn AJ, Ryan É J, Garry S, James DL, Boland MR, Young O, et al. Use of Intraoperative Parathyroid Hormone in Minimally Invasive Parathyroidectomy for Primary Hyperparathyroidism: A Systematic Review and Meta-analysis. *JAMA Otolaryngol Head Neck Surg*. 2021;147(2):135-43.

85. Ishii H, Mihai R, Watkinson JC, Kim DS. Systematic review of cure and recurrence rates following minimally invasive parathyroidectomy. *BJS Open*. 2018;2(6):364-70.
86. Barczynski M, Konturek A, Hubalewska-Dydejczyk A, Cichon S, Nowak W. Evaluation of Halle, Miami, Rome, and Vienna intraoperative iPTH assay criteria in guiding minimally invasive parathyroidectomy. *Langenbeck's Archives of Surgery*. 2009;394(5):843-9.
87. Zammit M, Pierce K, Bailey L, Rowland M, Waghorn A, Shore S. Challenging NICE guidelines on parathyroid surgery. *Surgeon*. 2022;20(4):e105-e11.
88. Norman J, Murphy C. Minimally invasive radio-guided parathyroidectomy. *Operative Techniques in General Surgery*. 1999;1(1):28-33.
89. Dudley N. Methylene blue for rapid identification of the parathyroids. *Br Med J*. 1971;3(5776):680-1.
90. Patel H, Chadwick D, Harrison B, Balasubramanian S. Systematic review of intravenous methylene blue in parathyroid surgery. *Journal of British Surgery*. 2012;99(10):1345-51.
91. Khavandi A, Whitaker J, Gonna H. Serotonin toxicity precipitated by concomitant use of citalopram and methylene blue. *Medical Journal of Australia*. 2008;189(9):534-5.
92. Patel AS, Singh-Ranger D, Lowery KA, Crinnion JN. Adverse neurologic effect of methylene blue used during parathyroidectomy. *Head & neck*. 2006;28(6):567-8.
93. Candell L, Campbell MJ, Shen WT, Gosnell JE, Clark OH, Duh Q-Y. Ultrasound-Guided Methylene Blue Dye Injection for Parathyroid Localization in the Reoperative Neck. *World Journal of Surgery*. 2014;38(1):88-91.
94. Brender E, Burke A, Glass RM. Frozen Section Biopsy. *JAMA*. 2005;294(24):3200-.
95. Iacobone M, Scarpa M, Lumachi F, Favia G. Are frozen sections useful and cost-effective in the era of intraoperative qPTH assays? *Surgery*. 2005;138(6):1159-65.
96. Aygün N, Uludağ M. Intraoperative Adjunct Methods for Localization in Primary Hyperparathyroidism. *Sisli Etfal Hastan Tip Bul*. 2019;53(2):84-95.
97. Abbaci M, De Leeuw F, Breuskin I, Casiraghi O, Lakhdar AB, Ghanem W, et al. Parathyroid gland management using optical technologies during thyroidectomy or parathyroidectomy: A systematic review. *Oral Oncology*. 2018;87:186-96.
98. Miccoli P, Pinchera A, Cecchini G, Conte M, Bendinelli C, Vignali E, et al. Minimally invasive, video-assisted parathyroid surgery for primary hyperparathyroidism. *J Endocrinol Invest*. 1997;20(7):429-30.
99. Henry JF, Raffaelli M, Iacobone M, Volot F. Video-assisted parathyroidectomy via the lateral approach vs conventional surgery in the treatment of sporadic primary hyperparathyroidism: results of a case-control study. *Surg Endosc*. 2001;15(10):1116-9.
100. Lanitis S, Chortis P, Sourtse G, Gkanis V, Lainas S, Tournis S, et al. Shifting from open to video-assisted parathyroidectomy: effect of the adjustment period on safety, clinical outcomes and cost. *Ann R Coll Surg Engl*. 2022;104(4):295-301.
101. Barczynski M, Cichon S, Konturek A, Cichon W. Minimally Invasive Video-Assisted Parathyroidectomy Versus Open Minimally Invasive Parathyroidectomy for a Solitary Parathyroid Adenoma: A Prospective, Randomized, Blinded Trial. *World Journal of Surgery*. 2006;30(5):721-31.
102. Gagner M. Endoscopic subtotal parathyroidectomy in patients with primary hyperparathyroidism. *Br J Surg*. 1996;83(6):875.
103. Henry J, Defechereux T, Gramatica L, De Boissezon C, editors. Endoscopic parathyroidectomy via a lateral neck incision. *Annales de Chirurgie*; 1999.
104. Brunaud L, Li Z, Van Den Heede K, Cuny T, Van Slycke S. Endoscopic and robotic parathyroidectomy in patients with primary hyperparathyroidism. *Gland Surg*. 2016;5(3):352-60.

105. Fouquet T, Germain A, Zarnegar R, Klein M, De Talance N, Claude Mayer J, et al. Totally endoscopic lateral parathyroidectomy: prospective evaluation of 200 patients. *Langenbeck's Archives of Surgery*. 2010;395(7):935-40.

**FEASABILITY OF FOCUSED PARATHYROIDECTOMY IN DEVELOPING COUNTRIES – A Scoping Review**

K. Rugnath<sup>1</sup>, S. Mewa Kinoo<sup>2</sup>

**1. Kapil Rugnath**

MBChB (UKZN), FCS (SA)

General Surgery Medical Officer, King Edward VIII Hospital, Durban, South Africa

Email: rugnath@gmail.com

**2. Suman Mewa Kinoo (corresponding author)**

MBChB (UKZN), FCS (SA), MMed (UKZN), Phd (UKZN)

Honorary Lecturer, Department of Surgery, University of KwaZulu Natal, Nelson R Mandela School of Medicine

General Surgeon, King Edward VIII Hospital, Durban, South Africa

Email: smewakinoo@gmail.com

## **Abstract**

### **Background**

The mainstay of treatment of primary hyperparathyroidism (HPT) involves a parathyroidectomy, which, depending on the number of affected parathyroid glands and the availability of resources, may involve a bilateral neck exploration (BNE) with 4 gland assessment or a minimally invasive, focused parathyroidectomy (FP) necessitating pre-operative localisation.

### **Objective**

We assess the feasibility of FP, in developing countries where limited resources may impede its use based on the potential lack of availability and or cost of these radiological studies and biochemical tests.

### **Methods**

A scoping review was performed with published literature evaluated from the past 15 years (2007 & onwards). Articles were screened and only included if they discussed focused parathyroidectomy, preoperative localisation, economic impact and they originated from a developing country (upper middle or lower middle-income).

### **Results**

A total of 18 articles met the inclusion criteria, comprising 7 developing countries (2 upper middle-income and 5 lower middle-income countries). Preoperative localisation was performed in all studies, with overall accuracy rates of 75,5% for ultrasound and 85,7% for <sup>99m</sup>Tc Sestamibi. A total 1202 patients (70%) had FP. Five hundred and fifty-five patients underwent FP without intraoperative adjuncts and 647 underwent FP with intraoperative adjuncts, with cure rates of 95,4% and 97,8% respectively. Overall cure rate for FP was 96,6%.

### **Conclusion**

With access to accurate preoperative localisation and excellent cure rates with and without intraoperative adjuncts, we conclude that FP is feasible in developing countries.

## Background

The ‘gold standard’ operative treatment of primary HPT has been BNE since the first parathyroidectomy in 1925, with excellent cure rates of greater than 95% demonstrated (1). The rationale at the time, was the knowledge of multi-gland disease coupled with the inability to identify specific diseased glands preoperatively. *Van Heerden et al (1991)* demonstrated a cure rate of 99,5% in 379 patients with primary HPT undergoing BNE (2). Consistently high success rates were further demonstrated by multiple authors, revealing this procedure to be reliable and easily reproducible (3, 4).

The FP was conceived due to the realisation that in 80% of cases of primary HPT the aetiology is a single gland adenoma, allowing a more targeted approach to parathyroidectomy (5). However, FP is naturally entirely reliant on adequate preoperative localisation. The greatest advancement being the <sup>99m</sup>Technetium Sestamibi, as described by *Coakley et al (1989)* (6). Today, there are numerous preoperative localisation studies available viz different types of Sestamibi radionucleotide scans, CT scans, MRI scans and ultrasound (US), each with varying degrees of sensitivities.

Initially, certain authors were firmly against the notion of FP, with some advising against the technique as they simply considered it to be a passing trend and misguided (7). They identified 2 main concerns with this technique.

The first was that of inferior cure rates. FP was highlighted by *Delbridge et al (2000)*, who assessed FP in 50 patients and demonstrated an inferior 84% cure rate with FP, significantly lower compared to the standard of care BNE’s established 95% cure rate (8). These inferior cure rates were mostly due to “double adenomas” which can account for up to 10% of all parathyroid adenomas (9). However, since then, intraoperative adjuncts used during parathyroidectomy evolved significantly, with intraoperative nuclear scanning and rapid PTH test (IOPTH) cure rates equalled that of BNE (10,11). When coupled with intraoperative assays, it also obviates inspection of the remaining PG’s thus preventing the complications associated with it like those that accompany BNE.

The second concern with FP was the cost, with some authors demonstrating a higher cost associated with FP compared to BNE, resulting from the requisite preoperative localisation tests and the addition of intraoperative adjuncts (12, 13). However, since then, the cost of previously elusive preoperative localisation studies decreased, and the literature reflected this by subsequently revealing FP to be cost effective comparative to BNE alone, balanced by the added cost of increased operative time & longer hospitalisation associated with the latter (14).

Subsequently, FP has been shown to be superior to BNE. *Udelsman et al* in 2000 with 100 cases, in 2002 with 255 cases and in 2011 with 1037 cases of FP demonstrated a higher cure rate with FP (99,4% vs 97,1% with BNE) with reduced cost, morbidity and length of hospital stay (15-17). Numerous other authors have demonstrated consistent findings (18-20). The benefits of FP over traditional BNE include equivalent rates of cure (95 – 98%), with added benefits of: the ability to be performed under a local anaesthetic; outpatient recovery; shorter operative time; desirable cosmesis; reduced nerve injuries and elimination of permanent hypoparathyroidism (21). Thus, in the setting of positive preoperative localisation, FP has become the new standard of care.

Importantly, however, the mounds of literature we are guided by emanate from the developed, ‘high-income world’, where resources are far more abundant, with remarkable strides made in healthcare resource allocation. Unfortunately, in a developing (middle-income) setting, with marked disparity between the private and public healthcare sector, resources are drastically limited, often to the point of tailoring guideline recommendations to local resource availability (23).

Therefore, despite the described cost-effectiveness of FP with requisite localisation & IOPTH in studies from developed world literature, a developing country may not be able to rationalise this cost. In addition to the cost, availability is a concern, which, when considered, would drastically affect the feasibility of this procedure, over a traditional BNE that requires no additional localization or intra-operative adjuncts.

## **Objective**

This scoping review serves to ascertain the feasibility of FP in developing countries, considering the additional cost and unclear availability of mandatory preoperative localisation studies along with requisite intraoperative adjuncts and the necessary skill and experience needed to facilitate this procedure successfully.

## **Methods**

This scoping review evaluated primary, secondary and tertiary literature, drawing from both published peer-reviewed and grey literature. The framework guide laid out by *Arksey & O’Malley (2005)*, including modifications published by *Levac et al (2010)*, were followed (24,25). The suggested stages traversed include: 1. identifying the research question; 2. identifying the relevant studies; 3. study selection, with specified inclusion and exclusion criteria; 4. charting the data and lastly 5. collating, summarising and reporting the results. Primarily, the PRISMA-Scr extension framework and checklist was adhered to, with reporting of eligible studies via the PRISMA flow diagram. The eligibility of the research question was explored via the PCC framework (26).

### Eligibility of Research Question

The PCC framework – namely; ‘population, concept and context’ framework, as recommended by the Joanna Briggs Institute (JBI), was employed (27). The population was identified as primary hyperparathyroid patients warranting parathyroidectomy. The concept was identified as cost and availability accounting for feasibility of focused parathyroidectomy comparative to bilateral neck exploration in developing countries. Finally, context was identified as developing countries.

## Search Criteria

An extensive literature search was conducted, using Boolean logic operators such as ‘AND’ & ‘OR’, with the following key terms: ‘primary hyperparathyroidism’, ‘parathyroidectomy’, ‘surgery’, ‘nuclear medicine’, ‘developing countries’, ‘cost’ and ‘South Africa’. The keywords used complied with Medical Subject Headings (MeSH) terms, adjoined with relevant subheadings & qualifiers, falling within the jurisdictions of the MeSH tree structures. To broaden the exposure to relevant articles, the following non-MeSH conforming keywords were additionally employed: ‘focused parathyroidectomy’, ‘feasibility’, ‘low-income’, ‘economic viability’, ‘Sestamibi’ and ‘intra-operative PTH’.

## Screening

Articles were retrieved upon exploration of the literature via the aforementioned library databases. Screening was initiated firstly by means of confirming relevance based on review of each title of the studies retrieved. Thereafter, abstracts were scanned, followed by introductions and statements of purpose. If the studies fulfilled the purpose of this review, the countries of origin were then scrutinized and cross checked with the United Nations Development Program (UNDP) & World Bank (WB) list of developing/middle income countries to ensure the articles to be perused were relevant ([28,29](#)). Finally, the full texts were engaged.

## Eligibility Criteria

### Inclusion Criteria

The basis for which these studies were selected includes: 1. studies providing evidence on surgical management of primary hyperthyroidism, specifically discussing focused or minimally invasive parathyroidectomy; 2. studies providing evidence on preoperative localisation; 3. studies providing evidence on a developing world approach to parathyroidectomy; and 4. studies providing evidence on economic burden of either parathyroidectomy option. The period explored was literature from the past 15 years (from 2007 onward). No restriction on language was employed, allowing for greater acquaintance.

### Exclusion Criteria

Studies not meeting the inclusion criteria of: providing evidence on surgical management of primary hyperthyroidism; with specific discussion on focused/minimally invasive parathyroidectomy; providing evidence on preoperative localization; providing evidence on a developing world approach to parathyroidectomy and providing evidence on economic burden of either parathyroidectomy option were all excluded, including all studies published prior to 2007.

## Results

A total of 940 relevant articles were encountered upon exploration of the literature from 8 different databases (*see Table 1: Evidence Sources*). After duplicates were removed and elimination by screening of article titles, abstracts, full texts and countries of origin, a total of 18 remaining articles fulfilled the purpose of this review and thus were included for study (*see Figure 1 PRISMA Flow Chart of Included Studies*).

In the 18 articles, a total 1716 patients underwent parathyroidectomy from 7 developing countries (5 studies from South Africa (upper middle-income); 5 studies from India (lower middle-income); 3 studies from Pakistan (lower middle-income); 2 studies from Turkey (upper middle-income); 1 study from Egypt (lower middle-income); 1 study from Algeria (lower middle-income and 1 study from Nigeria (lower middle-income). (*see Table 2: Characteristics of Studies*).

FP was performed in 1202 patients (70%) with 7 studies (38,9%) reporting on 647 patients having had FP with intraoperative adjuncts (including IOPTH, methylene blue, frozen section or the gamma probe) and 11 studies (61,1%) reporting on 555 patients having had FP without intra-operative adjuncts.


Cure rates from the studies included in this review for FP were reported for 15 (7 with intraoperative adjuncts & 8 without intraoperative adjuncts) of the 18 studies, ranging from 83,3% to 100%, with an overall cure rate of 96,6%. Below is a table detailing FP with/without intraoperative adjuncts and cure rates. (*see Table 3: Cure Rates*)

Preoperative localisation in these developing countries were performed by US and <sup>99m</sup>Tc Sestamibi, with 1 study describing the use of C11 methionine PET/CT as an additional localisation modality. Accuracy rates of localisation studies in these developing countries ranged from; 44% - 94% with US, 66,5% - 100% with <sup>99m</sup>Tc Sestamibi and 71,4% for the C11 methionine PET/CT.

*Table 1: Evidence Sources*

Database	No. of Studies Encountered
Pubmed	446
Google Scholar	217
ProQuest	132
Science Direct	61
WorldCat	39
Academic Search Complete	32
Sabinet	7
Langenbeck's Archive of Surgery	6
Total	940

Figure 1 PRISMA Flow Chart of Included Studies



(30)

Table 2: Characteristics of Studies

Scoping Review Data Collection Tool								
	Author	Title	Year	Country & UNDP/WB Classification <sup>(31)</sup>	Study Type	No. of Patients	Main Outcomes	Ref
1	Usta et al	A 20-Year Study on 190 Patients With Primary Hyperparathyroidism in a Developing Country: Turkey Experience	2015	Turkey (Developing/Upper-Middle Income Country)	Retrospective descriptive study	190	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 82,6%, and 89,4% respectively. Combined <sup>99m</sup> TcS & US yielded a 92% localisation accuracy rate. FP with intraoperative frozen section analysis only was performed in 167 patients and resulted in good cure rate of 99,52% - similar cure rate to BNE, with significantly decreased operative time, hospital stay & hospital cost.	(32)
2	Siddiqui et al	Changing paradigms in the surgical management of hyperparathyroidism at a tertiary care hospital in a developing country	2019	Pakistan (Developing/Lower-Middle Income Country)	Retrospective study	72	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 71,1%, and 90,1% respectively. FP without intraoperative adjuncts, was performed in 27 of the total 72 cases with a cure rate of 92,6%, similar cure rate of BNE (90,5%) with fewer complications - demonstrating this technique to be both effective & reasonable.	(33)
3	Paruk et al	Characteristics, management and outcome of primary hyperparathyroidism in South Africa: a single-centre experience	2013	South Africa (Developing/Upper-Middle Income Country)	Retrospective study	28	Preoperative localisation was only performed with <sup>99m</sup> TcS, with an accuracy rate of 93,7%. FP without IOPTH monitoring was performed in 19 of 28 patients, with a cure rate of 94,7%. Demonstrating the safety & feasibility of this technique	(34)
4	Mallikarjuna et al.	Five-year Retrospective Study on Primary Hyperparathyroidism in South India: Emerging Roles of Minimally Invasive Parathyroidectomy and Preoperative Localization with Methionine Positron Emission Tomography-Computed Tomography Scan	2018	India (Developing/Lower-Middle Income Country)	Retrospective study	54	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 72,2%, and 70,6% respectively. Additionally, C11 methionine PET was used for localisation in a portion of patients, with an accuracy rate of 71,4%. 43 patients underwent FP without intraoperative adjuncts, more than three-quarters (79,6%) of the total patients during the 5 year period, demonstrating this technique to be feasible in this locality of South India. No cure rate was reported.	(35)
5	El-Hady et al.	Focused parathyroidectomy for single parathyroid adenoma: a clinical account of 20 patients	2018	Egypt (Developing/Lower-Middle Income Country)	Prospective Analysis	20	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 75%, and 90% respectively. Combined <sup>99m</sup> TcS & US yielded a localisation accuracy rate of 95% in this setting. FP without IOPTH monitoring was performed in 19 of 20 patients, with a cure rate of 95% & no major complications reported, with the	(36)

							authors concluding this technique to be safe and feasibility, even without intra-operative adjuncts.	
6	Sen et al	Focused Parathyroidectomy Under Local Anesthesia – A Feasibility Study	2019	India (Developing/Lower-Middle Income Country)	Prospective Analysis	65	Preoperative localisation was performed by US & <sup>99m</sup> TcS, however the accuracy rate was not determined in this study as only patients with 100% concordance with pre-operative localisation were included in this study. FP without IOPTH monitoring was performed in 48 of 65 patients. 30 of the 48 cases were performed under local anaesthetic, with significantly reduced cost demonstrated and a cure rate of 100% reported, concluding this procedure to be safe and feasible.	(37)
7	Rawat et al	Minimally Invasive Parathyroidectomy as the Surgical Management of Single Parathyroid Adenomas: A Tertiary Care Experience	2022	India (Developing/Lower-Middle Income Country)	Prospective Analysis	116	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 93.10%, and 96.55% respectively. Combined <sup>99m</sup> TcS & US yielded a localisation accuracy rate of 100% in this study. FP with IOPTH monitoring was performed in all patients with a cure rate of 99,13% reported, with the authors concluding this procedure to be safe and feasible.	(38)
8	Zitouni et al	Monocentric experience of primary hyperparathyroidism surgery in Algeria	2021	Algeria (Developing/Lower-Middle Income Country)	Retrospective study	62	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 60,7%, and 66.5% respectively. Combined <sup>99m</sup> TcS & US accuracy rate was 100% in this study. 28 of 62 patients had MIP (unilateral exploration + focused), 18 of whom specifically had a FP without IOPTH monitoring, with a cure rate of 83,3% - demonstrating the safety & feasibility of this procedure.	(39)
9	Van Wyngaard et al	Pre operative localisation and surgical outcomes for Primary Hyperparathyroidism (PHPT): an 11 year review at a South African hospital	2018	South Africa (Developing/Upper-Middle Income Country)	Retrospective study	98	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 44%, and 75% respectively. FP with IOPTH monitoring was performed in 76% of patients [75 patients] with the remaining 24% having had BNE. The overall cure rate was 94%. This study demonstrates the safety and feasibility of FP.	(40)
10	MacRobert et al	Single gland versus multigland disease in primary hyperparathyroidism at the Wits Donald Gordon Medical Centre in Johannesburg, South Africa	2021	South Africa (Developing/Upper-Middle Income Country)	Retrospective study	252	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS for single gland disease was 77,1%, and 72,9% respectively, Combined <sup>99m</sup> TcS & US yielded a sensitivity of 79.7% for single gland disease. FP was performed in 44.8% of patients [113 patients] with no documented use of intraoperative adjuncts, with the remaining having had BNE. No cure rate was established due to lack of 6-month postoperative results. This study re-enforced the notion that BNE is the gold standard, however, with a predominance of single gland disease (83,3%), this study advocates equally for FP and BNE.	(41)

11	Olatoke et al	Serial pathologic fractures of five long bones on four separate occasions in a patient with primary hyperparathyroidism, challenges of management in a developing country: a case report	2013	Nigeria (Developing/Lower-Middle Income Country)	Case Report	1	In this case report, the parathyroid adenoma was successfully localised with combined US and <sup>99m</sup> TcS. Despite numerous challenges such as the <sup>99m</sup> TcS facility being 250km away and PTH testing needing to be done in another African country, the Authors persisted and were able to perform a focused unilateral exploration, with a demonstrated drop in postoperative PTH levels. This case report demonstrates while it may be possible to perform a parathyroidectomy in this country, with PTH needing to be performed in another country and the distance of travel required for pre-operative localisation, there are significant challenges associated with both BNE & FP.	(42)
12	Bombil et al	Sonar guided focused parathyroidectomy under cervical block	2018	South Africa (Developing/Upper-Middle Income Country)	Retrospective study	15	This study looked specifically at FP performed under a cervical block +/- local anaesthetic. Although no exact rate was provided, combined US and <sup>99m</sup> TcS was used for preoperative localisation, with a report of good localisation/concordance, and mention of only one discordant finding. 100% of patients had a FP with complete resolution of hypercalcaemia in all patients, demonstrating this surgical technique to be safe and feasible.	(43)
13	Yadav et al	Surgical Management of Primary Hyperparathyroidism in the Era of Focused Parathyroidectomy. A Study in Tertiary Referral Centre of North India	2019	India (Developing/Lower-Middle Income Country)	Retrospective study	373	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 80%, and 86% respectively. FP was performed in 49,32% of patients [184 patients], with IOPTH monitoring being performed in 85% of FP cases [151 patients] and 66.21% of total cases [247]. There was no statistical difference in cure with (98,02%) or without IOPTH (100%). An overall cure rate of 97,8% was reported, with a cure rate of 98,92% for FP, statistically similar to that of BNE, with the authors concluding FP to be safe and feasible, with or without IOPTH monitoring.	(44)
14	Dahiya et al	Surgical outcome after focused parathyroidectomy: experience from a tertiary care centre in North India	2021	India (Developing/Lower-Middle Income Country)	Retrospective study	192	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 83%, and 88% respectively. FP without IOPTH was performed in all 192 patients, with a cure rate of 97,92%. The authors conclude the technique to be the treatment of choice for sporadic primary HPT, when there is accurate pre-operative localisation, thus demonstrating the safety and feasibility of this technique.	(45)

15	Alikor et al.	The Usefulness of <sup>99m</sup> technetium-sestamibi Parathyroid Scintigraphy in Preoperative Localization of Parathyroid Adenoma in Patients with Primary Hyperparathyroidism at an Academic Hospital in South Africa	2017	South Africa (Developing/Upper-Middle Income Country)	Retrospective study	11	Preoperative localisation was performed only with <sup>99m</sup> TcS, which demonstrated a sensitivity of 100% in this study, allowing FP in all 11 patients, with intraoperative use of the gamma probe (radio-guided). Cure rate was reported to be 100%, with 14 adenomas removed from the total 11 patients (2 with multiple adenomas detected at pre-operative localisation and confirmed intra-operatively). The authors conclude <sup>99m</sup> TcS to be useful in this setting, and have resultantly shown FP to be feasible.	(46)
16	Afzal et al	Management of hyperparathyroidism: a five year surgical experience	2011	Pakistan (Developing/Lower-Middle Income Country)	Retrospective study	35	32 of the total patients had primary HPT, with preoperative localisation with combined <sup>99m</sup> TcS & US, yielding a reported 100% accuracy rate. No individual localisation rates were provided. FP without IOPTH was performed in 27 cases (84,4%) and BNE in 5 cases (15,6%). 100% cure rate was revealed for both techniques, demonstrating safety and feasibility of this technique, with the authors concluding accurate pre-operative localisation & safe surgery allows total cure for these patients.	(47)
17	Baloch et al	Surgical management of hyperparathyroidism	2007	Pakistan (Developing/Lower-Middle Income Country)	Observational Case Series	16	11 of the total 16 patients in this case series presented with primary HPT, with preoperative localisation rate by combined US and <sup>99m</sup> TcS accurate in 100% of cases, allowing FP, with intraoperative methylene blue, in this group. The remaining patients had secondary HPT with parathyroid hyperplasia and total parathyroidectomy via BNE with auto-transplantation was performed for these cases. These techniques both demonstrated a 100% cure rate, with the authors concluding both techniques to be satisfactory treatments for their respective parathyroid pathology (primary vs secondary).	(48)
18	Koyuncu et al	Minimally invasive surgery in primary hyperparathyroidism	2023	Turkey (Developing/Upper-Middle Income Country)	Retrospective Analysis	116	Preoperative localisation accuracy rate by US & <sup>99m</sup> TcS was 94%, and 95% respectively. FP with intraoperative frozen section was performed in all 116 patients, with a cure rate of 94,2% and morbidity of 1,7%. The authors conclude the technique to be the preferred treatment of choice for single gland disease, demonstrating the safety and feasibility of this technique.	(49)

UNDP = United Nations Development Program / WB = World Bank / Ref = references / US = ultrasound / <sup>99m</sup>TcS = <sup>99m</sup>Tc Sestamibi / FP = focused parathyroidectomy / MIP = minimally invasive parathyroidectomy / BNE = bilateral neck exploration / IOPTH = intraoperative parathyroid hormone / HPT = hyperparathyroidism

*Table 3: Cure Rates*

	Number of Studies	Number of Patients	Cure Rate
<b>Focused Parathyroidectomy</b>	18	1202	96,6%
<b>Without Intra-operative Adjuncts</b>	11	555 (46,2%)	95,4%
<b>Intra-operative Adjuncts</b>	7	647 (53,8%)	97,8%
• <b>IOPTH</b>	3	342 (28,5%)	97,1%
• <b>Frozen Section</b>	2	283 (23,5%)	96,9%
• <b>Methylene Blue</b>	1	11 (0,9%)	100%
• <b>Gamma Probe</b>	1	11 (0,9%)	100%

## Discussion

Whilst the operative management of primary HPT has undergone profound evolutionary changes resulting in FP being considered the new standard of care for cases of primary HPT with unequivocally localised single gland disease, as recommended by the American Association of Endocrine Surgeons (AAES) and the Fifth International Workshop on Primary Hyperparathyroidism, the concern for practitioners in developing low and middle-income countries is the relevance of the evolutionary changes and the applicability of its outcome, namely FP, considering the basis for it all emanates from developed high-income countries ([50,51](#)).

As demonstrated by *Hurst et al (2005)* and *Adhikari et al (2021)*, evidence-based medicine and the guidelines they produce from high-income countries most often cannot be adhered to in resource constrained low and middle-income countries ([52,53](#)). As such, this review was conducted to address the question of feasibility of FP in terms of accessibility to and accuracy of preoperative localisation and adequacy of cure rates with this technique, in developing countries.

The first prerequisite to enable performing FP is adequate preoperative localisation. From this review, it was revealed that preoperative localisation was available in all developing countries from the articles included, with the most common modality used being the  $^{99m}\text{Tc}$  Sestamibi (used in all studies) followed by US (used in 16 of the 18 studies).

The overall accuracy of  $^{99m}\text{Tc}$  Sestamibi and US for preoperative localisation in these developing countries were 85,7% and 75,7% respectively. The overall accuracy of combined US and  $^{99m}\text{Tc}$  Sestamibi was 95,2%. These results are in keeping with preoperative localisation rates reported from high-income countries. *Cheung et al (2012)* in their meta-analysis revealed sensitivities for  $^{99m}\text{Tc}$  Sestamibi and US of 78,9% and 76,1% respectively ([54](#)). Combined  $^{99m}\text{Tc}$  Sestamibi/US use for preoperative localisation has been reported to have a sensitivity of 96%, in keeping with the sensitivity of preoperative localisation from the developing countries in this study ([55](#)).

In a single lower middle-income country, Nigeria, *Olatoke et al (2013)* demonstrated preoperative localisation to be possible, but not feasible, as each  $^{99m}\text{Tc}$  Sestamibi test required a 250km journey to access a facility with the necessary equipment ([42](#)).

Intraoperative adjuncts were used in 7 of the total 18 studies: IOPTH was used in 3 studies; frozen section in 2 studies; methylene blue in 1 study and the gamma probe in 1 study. The cure rate for patients undergoing FP with IOPTH was 97,1%, with an overall cure of 97,8% for patients undergoing FP with intraoperative adjuncts. In the remaining 11 studies of FP without intraoperative adjuncts, the overall cure rate was 95,4%.

These figures from the developing countries included are in keeping with reported cure rates from high-income countries. *Quinn et al (2021)* in their meta-analysis compared cure rates of patients undergoing FP with vs without IOPTH (56). The authors reported cure rates of 98% for FP with IOPTH and 94,8% for FP without.

The lowest cure rate in this review was reported by *Zitouni et al (2021)* from the lower middle-income Algeria, who reported a cure rate of 83,3% with FP (39). The limitation with this study was the low sample size of only 18 patients who underwent FP, thus the 83,3% represents 15 of the 18 cured, with 3 patients having persistence. Additionally, individual preoperative localisation accuracy rates were relatively low in this study, with US accurate in 60,7% of cases and <sup>99m</sup>Tc Sestamibi accurate in 66,5% of cases.

With the 18 studies from developing countries included in this review, a total of 1202 patients underwent FP with an overall cure rate of 96,6%. This is in keeping with reported cure rates from high income countries. *Ishii et al (2018)* in their meta-analysis of 5282 patients, from 14 included studies having undergone FP, demonstrated an overall cure rate of 96,9% (57).

From this review it is evident that not only is preoperative localisation accessible and accurate in these developing countries, but FP is also successfully being performed within these regions. The 2 lower middle-income countries with difficulties were Nigeria and Algeria, demonstrating that perhaps the upper-middle income developing regions are currently better suited to adapting to this minimally invasive technique.

## **Conclusion**

Preoperative localisation is available in developing countries, with adequate accuracy rates, in keeping with rates described in literature from high-income countries. FP is performed in developing countries with adequate cure rates in keeping with that of RCT's published from high-income countries. Although some developing countries lack availability of certain intraoperative adjuncts, this was not prohibitive and cure rates with or without these adjuncts were similar in these countries.

With the established availability & impressive accuracy rates of preoperative localisation and the demonstrated success rates of FP, it is evident this technique is safe and feasible in developing countries.

## **Limitations**

There were only 7 developing countries included in this review, thus limiting the relevance of this review to other developing countries. A small number of studies did not report individual preoperative localisation rates or overall cure rates, thus distorting representation of the overall localisation & cure rates.

## **Informed consent**

Not applicable

## **Ethics**

This study was approved for ethics exemption by the UKZN Biomedical Research Ethics Committee (BREC). BREC Reference: BREC/00005172/2023

## **Financial Support and Sponsorship**

None

## **Conflict of Interest**

None

## **References**

1. Lee NC, Norton JA. Multiple-Gland Disease in Primary Hyperparathyroidism: A Function of Operative Approach? *Archives of Surgery*. 2002;137(8):896-900.
2. van Heerden JA, Grant CS. Surgical treatment of primary hyperparathyroidism: An institutional perspective. *World Journal of Surgery*. 1991;15(6):688-92.
3. Udén P, Chan A, Duh Q-Y, Siperstein A, Clark OH. Primary hyperparathyroidism in younger and older patients: Symptoms and outcome of surgery. *World Journal of Surgery*. 1992;16(4):791-7.
4. Chen H, Zeiger MA, Gordon TA, Udelsman R. Parathyroidectomy in Maryland: effects of an endocrine center. *Surgery*. 1996;120(6):948-53.
5. Fraser WD. Hyperparathyroidism. *The Lancet*. 2009;374(9684):145-58.
6. Coakley AJ, Kettle AG, Wells CP, O'Doherty MJ, Collins RE. <sup>99</sup>Tc m sestamibi--a new agent for parathyroid imaging. *Nucl Med Commun*. 1989;10(11):791-4.
7. Delbridge LW, Younes NA, Guinea AI, Reeve TS, Clifton-Bligh P, Robinson BG. Surgery for primary hyperparathyroidism 1962-1996: indications and outcomes. *Med J Aust*. 1998;168(4):153-6.
8. Delbridge LW, Dolan SJ, Hop TT, Reeve TS, Robinson BG, Wilkinson MR. Minimally invasive parathyroidectomy: 50 consecutive cases. *Medical journal of Australia*. 2000;172(9):418-22.
9. Goodsell KE, Ermer JP, Zaheer S, Kelz RR, Fraker DL, Wachtel H. Double adenoma as a cause of primary hyperparathyroidism: Asymmetric hyperplasia or a distinct pathologic entity? *The American Journal of Surgery*. 2021;222(3):483-9.
10. Eslamy HK, Ziessman HA. Parathyroid Scintigraphy in Patients with Primary Hyperparathyroidism: <sup>99m</sup>Tc Sestamibi SPECT and SPECT/CT. *RadioGraphics*. 2008;28(5):1461-76.
11. Chen H, Pruhs Z, Starling JR, Mack E. Intraoperative parathyroid hormone testing improves cure rates in patients undergoing minimally invasive parathyroidectomy. *Surgery*. 2005;138(4):583-90.
12. Aarum S, Nordenström J, Reihner E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for primary hyperparathyroidism: the new versus the old order: a randomised controlled trial of preoperative localisation. *Scandinavian journal of surgery*. 2007;96(1):26-30.

13. Baliski C, Nosyk B, Melck A, Bugis S, Rosenberg F, A HA. The cost-effectiveness of three strategies for the surgical treatment of symptomatic primary hyperparathyroidism. *Ann Surg Oncol.* 2008;15(10):2653-60.
14. Fahy BN, Bold RJ, Beckett L, Schneider PD. Modern Parathyroid Surgery: A Cost-benefit Analysis of Localizing Strategies. *Archives of Surgery.* 2002;137(8):917-23.
15. Udelsman R, Donovan PI, Sokoll LJ. One hundred consecutive minimally invasive parathyroid explorations. *Ann Surg.* 2000;232(3):331-9.
16. Udelsman R. Surgery in primary hyperparathyroidism: the patient without previous neck surgery. *J Bone Miner Res.* 2002;17 Suppl 2:N126-32.
17. Udelsman R, Lin Z, Donovan P. The superiority of minimally invasive parathyroidectomy based on 1650 consecutive patients with primary hyperparathyroidism. *Ann Surg.* 2011;253(3):585-91.
18. Pang T, Stalberg P, Sidhu S, Sywak M, Wilkinson M, Reeve T, et al. Minimally invasive parathyroidectomy using the lateral focused mini-incision technique without intraoperative parathyroid hormone monitoring. *Journal of British Surgery.* 2007;94(3):315-9.
19. Wong W, Foo FJ, Lau M, Sarin A, Kiruparan P. Simplified minimally invasive parathyroidectomy: a series of 100 cases and review of the literature. *The Annals of The Royal College of Surgeons of England.* 2011;93(4):290-3.
20. Norlén O, Wang KC, Tay YK, Johnson WR, Grodski S, Yeung M, et al. No need to abandon focused parathyroidectomy: a multicenter study of long-term outcome after surgery for primary hyperparathyroidism. *Ann Surg.* 2015;261(5):991-6.
21. Slepavicius A, Beisa V, Janusonis V, Strupas K. Focused versus conventional parathyroidectomy for primary hyperparathyroidism: a prospective, randomized, blinded trial. *Langenbecks Arch Surg.* 2008;393(5):659-66.
22. Greene AK, Mowschenson P, Hodin RA. Is sestamibi-guided parathyroidectomy really cost-effective? *Surgery.* 1999;126(6):1036-40; discussion 40-1.
23. Peabody JW, Taguiwalo MM, Robalino DA, Frenk J. Improving the Quality of Care in Developing Countries. In: nd, Jamison DT, Breman JG, Measham AR, Alleyne G, Claeson M, et al., editors. *Disease Control Priorities in Developing Countries.* Washington (DC)2006.
24. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International journal of social research methodology.* 2005;8(1):19-32.
25. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci.* 2010;5:69.
26. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med.* 2018;169(7):467-73.
27. Peters MDJ GC, McInerney P, Munn Z, Tricco AC, Khalil, H. . Chapter 11: Scoping Reviews (2020 version). In: Aromataris E, Munn Z (Editors). *JBIM Manual for Evidence Synthesis*, JBI, 2020. Available from <https://synthesismanual.jbi.global>. <https://doi.org/10.46658/JBIMES-20-12>.
28. Program UND. Developing Regions, Documentation & Resources, Data Downloads 2023. <https://hdr.undp.org/data-center/documentation-and-downloads> [
29. Bank TW. Middle Income, Overview 2023 <https://data.worldbank.org/country/XP> [
30. Peters M, Godfrey C, Khalil H, McInerney P, Soares C, Parker D. 2017 Guidance for the Conduct of JBI Scoping Reviews. 2017.
31. Gbadamosi A. Understanding the Developed/Developing Country Taxonomy. *Advocates for International Development.* June 2013;Category Archives DOI <https://www.a4id.org/policy/understanding-the-developeddeveloping-country-taxonomy/>.

32. Usta A, Alhan E, Cinel A, Türkyılmaz S, Erem C. A 20-year study on 190 patients with primary hyperparathyroidism in a developing country: Turkey experience. *Int Surg.* 2015;100(4):648-55.
33. Siddiqui MI, Pasha HA, Asad R, Talati JJ. Changing paradigms in the surgical management of hyperparathyroidism at a tertiary care hospital in a developing country. *J Pak Med Assoc.* 2019;69(9):1360-4.
34. Paruk IM, Esterhuizen TM, Maharaj S, Pirie FJ, Motala AA. Characteristics, management and outcome of primary hyperparathyroidism in South Africa: a single-centre experience. *Postgrad Med J.* 2013;89(1057):626-31.
35. Mallikarjuna VJ, Mathew V, Ayyar V, Bantwal G, Ganesh V, George B, et al. Five-year Retrospective Study on Primary Hyperparathyroidism in South India: Emerging Roles of Minimally Invasive Parathyroidectomy and Preoperative Localization with Methionine Positron Emission Tomography-Computed Tomography Scan. *Indian J Endocrinol Metab.* 2018;22(3):355-61.
36. El-Hady HA, Radwan HS. Focused parathyroidectomy for single parathyroid adenoma: a clinical account of 20 patients. *Electron Physician.* 2018;10(6):6974-80.
37. Sen S, Cherian A, Ramakant P, Reka K, Paul M, Abraham D. Focused parathyroidectomy under local anesthesia – A feasibility study. *Indian Journal of Endocrinology and Metabolism.* 2019;23(1):67-71.
38. Rawat A, Grover M, Kataria T, Samdhani S, Mathur S, Sharma B. Minimally Invasive Parathyroidectomy as the Surgical Management of Single Parathyroid Adenomas: A Tertiary Care Experience. *Indian Journal of Otolaryngology and Head & Neck Surgery.* 2022:1-7.
39. Nouikes Zitouni S. Monocentric experience of primary hyperparathyroidism surgery in Algeria. *Surg Open Sci.* 2021;4:32-6.
40. Van Wyngaard T. Pre-operative localization and surgical outcomes for primary hyperparathyroidism (PHPT): An 11-year review at a South African hospital: University of Cape Town; 2018.
41. MacRobert NA, Kruger D, Schamm M. A Retrospective Audit of Single Versus Multigland Disease in Primary Hyperparathyroidism at a Single Centre in South Africa. *Wits Journal of Clinical Medicine.* 2021;3(2):93-100.
42. Olatoke SA, Agodirin OS, Rahman GA, Habeeb OG, Jimoh RO, Ahmed BA, et al. Serial pathologic fractures of five long bones on four separate occasions in a patient with primary hyperparathyroidism, challenges of management in a developing country: a case report. *Pan Afr Med J.* 2013;15:45.
43. Bombil I, Louw L, Mitchell C, Mahlobo F, Ramazani AM, Nthatheni RM. Sonar guided focused parathyroidectomy under cervical block. *South African Journal of Surgery.* 2018;56(2):30-3.
44. Yadav S, Mishra S, Mishra A, Mayilvagnan S, Chand G, Agarwal G, et al. Surgical management of primary hyperparathyroidism in the era of focused parathyroidectomy: A study in tertiary referral centre of North India. *Indian Journal of Endocrinology and Metabolism.* 2019;23(4):468-72.
45. Dahiya D, Abuji K, Kumari P, Gautam A, Bhadada S, Sood A, et al. Surgical outcome after focused parathyroidectomy: experience from a tertiary care centre in North India. *Pol Przegl Chir.* 2021;93(5):1-5.
46. Alikor C, Engelbrecht G. The Usefulness of <sup>99m</sup>technetium-sestamibi Parathyroid Scintigraphy in Preoperative Localization of Parathyroid Adenoma in Patients with Primary Hyperparathyroidism at an Academic Hospital in South Africa. *Asian Journal of Medicine and Health.* 2017;6:1-7.

47. Afzal A, Gauhar TM, Butt WT, Khawaja AA, Azim KM. Management of hyperparathyroidism: a five year surgical experience. JPMA The Journal of the Pakistan Medical Association. 2011;61(12):1194-8.
48. Baloch MN, Aslam T, Maher M. Surgical management of hyperparathyroidism. J Coll Physicians Surg Pak. 2007;17(11):683-5.
49. Koyuncu A, Ozer SP, Ozer B, Catal O, Sit M. Minimally invasive surgery in primary hyperparathyroidism. Journal of Bionic Memory. 2023;3(1):1-6.
50. Wilhelm SM, Wang TS, Ruan DT, Lee JA, Asa SL, Duh QY, et al. The American Association of Endocrine Surgeons Guidelines for Definitive Management of Primary Hyperparathyroidism. JAMA Surg. 2016;151(10):959-68.
51. Bilezikian JP, Khan AA, Silverberg SJ, Fuleihan GE-H, Marcocci C, Minisola S, et al. Evaluation and Management of Primary Hyperparathyroidism: Summary Statement and Guidelines from the Fifth International Workshop. Journal of Bone and Mineral Research. 2022;37(11):2293-314.
52. Adhikari S. Evidence-based medicine in low-income and middle-income countries. Lancet Glob Health. 2021;9(7):e903-e4.
53. Hurst SA, Hull SC, DuVal G, Danis M. Physicians' Responses to Resource Constraints. Archives of Internal Medicine. 2005;165(6):639-44.
54. Cheung K, Wang TS, Farrokhyar F, Roman SA, Sosa JA. A meta-analysis of preoperative localization techniques for patients with primary hyperparathyroidism. Ann Surg Oncol. 2012;19(2):577-83.
55. Arici C, Cheah WK, Ituarte PH, Morita E, Lynch TC, Siperstein AE, et al. Can localization studies be used to direct focused parathyroid operations? Surgery. 2001;129(6):720-9.
56. Quinn AJ, Ryan É J, Garry S, James DL, Boland MR, Young O, et al. Use of Intraoperative Parathyroid Hormone in Minimally Invasive Parathyroidectomy for Primary Hyperparathyroidism: A Systematic Review and Meta-analysis. JAMA Otolaryngol Head Neck Surg. 2021;147(2):135-43.
57. Ishii H, Mihai R, Watkinson JC, Kim DS. Systematic review of cure and recurrence rates following minimally invasive parathyroidectomy. BJS Open. 2018;2(6):364-70.

## Appendices

### Appendix 1: The Final Study Protocol

# Title: FEASIBILITY OF FOCUSED PARATHYROIDECTOMY FOR PRIMARY HYPERPARATHYROIDISM IN A DEVELOPING COUNTRY

<b>Principal Investigator:</b>	Kapil <u>Rugnath</u>
<b>Institution:</b>	University of Kwa-Zulu Natal Medical School 719 <u>Umbilo Road</u> , Durban Private Bag 7 <u>Congella</u> 4013  Tel: (031) 260 4216
<b>Supervisor:</b>	Suman <u>MewaKinoo</u>

## TABLE OF CONTENTS

<u>1.</u>	<u>BACKGROUND AND RATIONALE</u>	46
<u>1.1.</u>	<u>Introduction</u>	46
<u>1.2.</u>	<u>Background</u>	47
<u>1.3.</u>	<u>Rationale</u>	48
<u>2.</u>	<u>OBJECTIVES AND OUTCOME MEASURES/ENDPOINTS</u>	49
<u>2.1.</u>	<u>Overall Aim:</u>	49
<u>2.2.</u>	<u>Primary Objectives:</u>	49
<u>2.3.</u>	<u>Primary Hypothesis:</u>	49
<u>3.</u>	<u>METHODS [16-21]</u>	49
<u>3.1.</u>	<u>Study design</u>	49
<u>3.2.</u>	<u>Setting and study population</u>	50
<u>3.3.</u>	<u>Subjects</u>	50
<u>3.3.1</u>	<u>Sampling technique</u>	50
<u>3.3.2</u>	<u>Selection criteria</u>	50
<u>3.4.</u>	<u>Screening, enrolment and study procedures</u>	51
<u>3.5</u>	<u>Measures</u>	51
<u>3.5.1</u>	<u>Explanatory variables</u>	51
<u>3.5.2</u>	<u>Outcome variables</u>	51
<u>3.6</u>	<u>Data management plan</u>	51
<u>3.6.1</u>	<u>Data collection</u>	51
<u>3.6.2</u>	<u>Validity</u>	51
<u>3.6.3</u>	<u>Data entering, storage and validation</u>	52
<u>3.7</u>	<u>Statistical considerations, Sample size and power</u>	52
<u>Data analysis plan</u>		52
<u>3.8</u>	<u>Ethical considerations</u>	52
<u>3.8.1</u>	<u>Informed consent</u>	52
<u>3.8.2</u>	<u>Confidentiality</u>	52
<u>3.8.3</u>	<u>Conflict of interest</u>	52
<u>4.</u>	<u>STUDY TIMELINE</u>	52
<u>5.</u>	<u>STUDY LIMITATIONS</u>	53
<u>6.</u>	<u>FUTURE DIRECTIONS</u>	53

<a href="#">7. BUDGET</a> .....	53
- <a href="#">Justification of budget</a> .....	53
<a href="#">8. REFERENCES [1-25]</a> .....	53
.....	55
<a href="#">9. Appendices</a> .....	55

# 1. BACKGROUND AND RATIONALE

## 1.1 Introduction

The parathyroid glands (PG) are specialised endocrine structures located in the neck, posterior to the thyroid gland, within the pre-tracheal fascia. Their main function is to produce parathyroid hormone (PTH), which is critical for calcium homeostasis, and thus normal physiologic function. Ordinarily, 4 glands are present: 1 pair superiorly and 1 pair inferiorly. However, variability in number (2 – 6) and location ( $\approx$  16% ectopia including: mediastinum, retro-oesophageal and even lateral neck sites) is well documented. [1]

Parathyroid disease most often presents in the form of hypersecretion of PTH – termed hyperparathyroidism, which is the third most common endocrine disorder, following diabetes mellitus and thyroid disease. [2] Considering PTH is responsible for maintaining eucalcaemia, any hypersecretion results in hypercalcaemia. Thus, the symptomatic presentation in these patients involve hypercalcaemic symptoms; so-called ‘stones, bones, groans and psychic moans’ – derived from kidney stones, osteoporotic bone pain, peptic ulcer disease with abdominal pain groans and impaired cognition with psychic moans.

Hyperparathyroidism (HPT) may be classified broadly into 3 categories:

- Primary
- Secondary
- Tertiary

Primary HPT results from a direct PTH over-production from the PG, most often as a result of a parathyroid adenoma: and less commonly as a result of multiple adenoma’s, parathyroid hyperplasia, parathyroid carcinoma and Multiple Endocrine Neoplasia (MEN) 1 and 2A.

Secondary HPT occurs most commonly as a consequence of chronic kidney disease (CKD). Specifically, secondary HPT results from PG over stimulation, PTH hypersecretion and resultant PG hyperplasia, in response to hypocalcaemia of CKD.

Tertiary HPT represents a persistence and progression of disease from secondary HPT in the setting of hypocalcaemic induced chronic PG over-stimulation, to profound hyperplasia and autonomous PTH secretion, persisting incessantly, post renal transplant.

The management of primary HPT necessitates parathyroidectomy in symptomatic patients and asymptomatic patients in certain instances. There is controversy as to which approach viz, bilateral neck exploration (BNE) or focused parathyroidectomy (FP) is superior in terms of cost and complications.

Surgical intervention for secondary & tertiary HPT, when indicated, always involves BNE as the pathophysiology in these two entities entails diffuse hyperplasia, and thus all four glands need to be inspected and the approach is therefore not an area of debate.

## 1.2 Background

Parathyroidectomy by way of BNE was first described by Felix Mandl in 1925, and subsequently became the recommended gold standard for all HPT requiring surgical intervention, with a reported cure rate of 95%. [3],[4] The rationale at the time, was the knowledge of multi-gland disease coupled with the inability to identify specific diseased glands pre-operatively. Since then, the standard of care has undergone a myriad of evolutionary advancements.

Minimally Invasive Parathyroidectomy (MIP), also known as FP or ‘targeted’ parathyroidectomy soon came to the fore. FP involves identification of the specific diseased PG pre-operatively and when coupled with intra-operative assays, affords the benefit of providing the surgeon with a roadmap, allowing a simple localised miniature incision directly over the affected gland, with easier excision, and obviation of inspection of the remaining PG’s.

Born from the realisation that primary HPT is often a consequence of a solitary parathyroid adenoma (involving a single gland) in 80% of cases, the first less invasive rung on the parathyroidectomy evolutionary ladder, promoted by Roth et al in 1975 & Tibblin et al in 1982 [5],[6],[7] began with and entailed a unilateral neck exploration, without any pre-operative localisation. The side chosen for exploration was determined by either an oesophagogram, vascular PTH sampling, palpation or guess. The technique involved removal of all parathyroid tissue from one side, with the use of an intra-operative saline float test & oil red O staining. Excellent cure rates were realised, unfortunately only 43 of 102 patients revealed parathyroid pathology on the initial unilateral exploration, with 45 of 102 requiring contra-lateral exploration, thus a solution for correctly identifying the side of the offending PG pathology prior to exploration was desperately needed. [8]

With the advent of the nuclear scan Technetium 99m (Tc<sup>99</sup>) Sestamibi – also known as parathyroid MIBI (99m Tc-methoxy isobutyl isonitrile) in 1989 [9], pre-operative localization was realized. A radiotracer (Tc<sup>99</sup>) was injected intravenously where it flowed briefly until it was rapidly taken up by and localized in hyperfunctioning parathyroid tissue. A planar (2D) image was then taken with a nuclear medicine gamma camera which revealed the site of the offending hyperfunctioning tissue. This allowed identification of parathyroid adenomas and their exact location such that a single 10-20mm incision could be placed directly over the identified adenoma site, under a local, regional or general anaesthetic. Consequently, the modern FP was conceived.

This was revolutionary for cases with solitary adenomas, that were accurately localised. However, the accuracy of pre-op localization came to the fore as cases of missed multi-gland disease (multiple adenomas & multi-gland hyperplasia) arose. To address this problem, pre-operative localization progressed to entail a combination of the MIBI with existing ultrasonography. This realized a 13% increase in yield, however sensitivities for multi-gland disease remained concerning. [10]

Fortuitously, chemiluminometrics entered the fray, in the form of a quick PTH assay (QPTH). The QPTH is an intra-operative serum test, sampled post excision of adenoma, with a mere turn-around time of 15-20 minutes. Results revealing a more than 50% decline, compared to pre-operative levels, serve as confirmation of successful adenoma removal, eliminating the possibility of a potentially missed second adenoma, additional PG hyperplasia or hyperfunctioning ectopia. And, in the face of failed adequate QPTH decrease, then only a BNE

would be warranted to explore the remaining glands for additional unidentified pathology. The addition of the QPTH has reportedly improved cure rates significantly from 91.9% to 99.1%. [11]

In the decades following the advent of MIBI, pre-operative localisation would see drastic improvements in technology and resultant sensitivities. The MIBI scan itself advanced drastically with evolutions in:

- Timing sequence – resulting in dual phase imaging
- Additional radiotracer isotope – resulting in dual isotope usage
- Image acquisition – resulting in progress from the original planar (2D) images to 3D images, making use of single photon emission computed tomography (SPECT) technology

In addition to the advancements in MIBI, further newer pre-operative localization studies have been developed with exceptional sensitivities demonstrated. [12]

Localisation Study	Reported Sensitivity	Study
MIBI SPECT/CT	86%	Wong et al, 2015
4D CT (contrasted triple phase CT)	94%	Bahl et al, 2015
MRI	97.8%	Agiro et al, 2018
PET/MRI	96.2%	Huber et al, 2018
4D MRI	100%	Merchavy et al, 2016
<sup>18</sup> F-Fluorocholine PET ( <sup>18</sup> F-FCH PET)	100%	Huber et al, 2018
<sup>18</sup> F-FCH PET/4D CT	100%	Piccardo et al, 2019

An additional adjunct in the FP armamentarium, so-called minimally invasive radio-guided parathyroidectomy (MIRP), described by Murphy et al. (1999), details the injection of radioactive tracer Tc99 2-4 hours prior to surgery, coupled with the use of an intra-operative gamma probe. A prescribed reading of  $\geq 20\%$  background radioactivity (BRA) of a removed suspected adenoma serves as confirmatory, obviating the need for frozen section. [13] Additionally, less than 20% BRA in any remaining tissue, serves as confirmation of no remaining hyperfunction pathological tissue, thus additionally obviating the need for intra-operative QPTH. [13]

The benefits of FP over traditional BNE include equivalent rates of cure (95 – 98%) [4], with added benefits of; the ability to be performed under a local anaesthetic, outpatient recovery, shorter operative time, desirable cosmesis, reduced nerve injuries and elimination of permanent hypoparathyroidism. Thus, in the setting of positive pre-operative localization, FP has become the new standard of care.

### 1.3 Rationale

With the enormous benefits and de rigueur standing of FP, it would be tempting to neglect the additional costs involved in the obligatory pre-operative localization studies and intra-operative QPTH assays required. Thus, FP cost-effectiveness studies ensued, with initial conclusions weighing against the overall benefits of FP, with the economic burden of requisite investigations being unjustified. [3]

With improvements in technology and resource availability over time, the cost of previously elusive pre-operative localisation studies decreased, and the literature reflected this by subsequently revealing FP's to be cost effective comparative to BNE alone, balanced by the added cost of increased operative time and longer hospitalization associated with the latter. [14]

Importantly, however, the mounds of literature we are guided by emanates from the 'developed, high-income world', where resources are far more abundant, with remarkable strides made in healthcare resource allocation. Unfortunately, in a developing world (middle-income) setting, with marked disparity between the private and public healthcare sector, resources are drastically limited, often to the point of tailoring guideline recommendations to local resource availability. [15]

Therefore, despite the described cost-effectiveness of FP with requisite localization & QPTH in studies from developed world literature, a developing country, such as South Africa, may not be able to rationalise this cost. In addition to the cost, availability is a concern, which, when considered, would drastically affect the feasibility of this procedure, over a traditional BNE that requires no additional localization or intra-operative adjuncts.

## **2. OBJECTIVES AND OUTCOME MEASURES/ENDPOINTS**

### **2.1 Overall Aim:**

The aim of this scoping review serves to ascertain the feasibility, in terms of cost and availability, of the mandatory pre-operative localization studies, and intra-operative QPTH, necessary to facilitate a successful FP, for primary HPT in a developing country, South Africa.

### **2.2 Primary Objectives:**

1. To ascertain if focused parathyroidectomy is performed in South Africa
2. To ascertain if pre-operative localization with Sestamibi/SPECT is readily available
3. To ascertain if intra-operative quick-PTH assay is readily available
4. To assess the cost of Sestamibi and QPTH locally
5. To assess the cost of a BNE locally
6. To ascertain if there are any 'high volume' parathyroidectomy centers in South Africa
7. To assess the outcomes of FP vs BNE locally

### **2.3 Primary Hypothesis:**

Despite resource constraints in South Africa, focused parathyroidectomy requiring pre-operative localization and intra-operative QPTH is feasible

## **3. METHODS [16-21]**

### **3.1 Study design**

The research design will be in the form of a scoping review, making use of primary, secondary and tertiary literature, drawing from both published peer-reviewed and grey literature, specifically from within the last 15 years. The framework guide laid out by Arksey & O'Malley, including modifications published by Levac et al, will be followed. The suggested stages to be traversed include (1) identifying the research question; (2) identifying the relevant

studies; (3) study selection, with specified inclusion and exclusion criteria; (4) charting the data and lastly (5) collating, summarising and reporting the results.

### **3.2 Setting and study population**

The identified setting for this study is literature from developing countries, where resource constraints, in the form of prohibitive cost and lack of availability, often result in modification of 'best-practice' western guidelines.

### **3.3 Subjects**

#### 3.3.1 Sampling technique

##### - Search Criteria

An extensive literature search will be conducted, using Boolean logic operators such as 'AND' & 'OR', with the following key terms: 'primary hyperparathyroidism', 'parathyroidectomy', 'surgery', 'nuclear medicine', 'neck', 'developing countries', 'cost' and 'South Africa'. The keywords to be used comply with Medical Subject Headings (MeSH) terms, adjoined with relevant subheadings & qualifiers, falling within the jurisdictions of the MeSH tree structures. To broaden the exposure to relevant articles, the following non-MeSH conforming keywords will be additionally employed: 'focused parathyroidectomy', 'feasibility', 'developing world', 'economic viability', 'Sestamibi' and 'quick PTH'. The search will only look for literature from within the last 15 years.

##### - Identification of Evidence Sources

The extensive search will be conducted utilizing the following library databases: Pubmed/MEDLINE, Google Scholar, World Cat, Academic Search Complete via EBSCO host, Open Dissertations via EBSCO host, SA ePublications via SABINET African journals, SACat via OCLC and ProQuest.

These prodigious evidence sources were identified through their phenomenal reputations as leaders in library database provision, enabling access to a large cohort of invaluable research papers. The inclusion of local library databases, which forms part of the identified respectable evidence sources, stand to improve exposure to publications emanating from developing countries, thereby improving yield in addressing the research question.

#### 3.3.2 Selection criteria

##### - Inclusion criteria

The basis for which studies will be selected includes: (1) studies providing evidence on surgical management of primary hyperthyroidism; (2) studies providing evidence on pre-operative localization; (3) studies providing evidence on a developing world approach to parathyroidectomy; and (4) studies providing evidence on economic burden of either parathyroidectomy option. No restriction on language will be employed, allowing for greater acquaintance.

##### - Exclusion criteria

International studies on surgical approaches for Primary HPT not providing cost information.

Local studies not providing information on operative approach for Primary HPT.

### 3.4 Screening, enrolment and study procedures

Upon exploration of the literature, screening will be initiated firstly by means of confirming relevance based on review of each title of the studies retrieved. Thereafter, abstracts will be scanned, followed by introductions and statements of purpose. If the studies fulfill the purpose of this review, the full text will be engaged.

### 3.5 Measures

#### 3.5.1 Explanatory variables

1. Availability of Sestamibi scans in developing countries
2. Availability of QPTH developing countries
3. Cost of Sestamibi and QPTH
4. Availability of requisite parathyroid surgery expertise in developing countries
5. Availability of high-volume parathyroid surgery ‘centers of excellence’ in developing countries

#### 3.5.2 Outcome variables

1. Prevalence of focused parathyroidectomy in developing countries
2. Cost associated with the various parathyroidectomy options

### 3.6 Data management plan

#### 3.6.1 Data collection

All studies identified on the aforementioned library databases will be tabulated via use of Microsoft Excel. Further individual lists with specifics pertaining to included and excluded studies will additionally be tabulated as separate sheets, on one comprehensive document.

#### 3.6.2 Validity

The eligibility of the research question will be tested using the PCC framework – namely; ‘population, concept and context’ framework, as recommended by the Joanna Briggs Institute (JBI). The population is identified as primary hyperparathyroid patients warranting parathyroidectomy. The concept is identified as cost and availability accounting for feasibility of focused parathyroidectomy comparative to bilateral neck exploration. Finally, context is identified as the developing world setting, South Africa.

Population	Human subjects undergoing investigation and surgery for primary hyperparathyroidism
Concept	Feasibility of focused parathyroidectomy vs bilateral neck exploration
Context	Developing world setting (middle-income), South Africa

Furthermore, to ensure validity of the scoping review, the quality of studies included will be assessed by means of the 'JBI critical appraisal checklist for systematic reviews and research syntheses' tool. The utility of this tool lies in its ability to appraise a studies methodological quality, ensuring the possibility of bias in either its design, conduct or analysis, is identified. (Appendix 1)

### 3.6.3 Data entering, storage and validation

The database making use of Microsoft Excel will be directly created and curated by the PI. Ongoing quality assurance will be completed prior to data entry. After entry, data will be edited and cross-checked such that no omissions and/or errors occur. Additionally, the database will be scrutinized by the Supervisor ensuring complete accuracy. Data will be stored on the study PI's computer and will be backed up daily on the central server; access will be limited to the PI.

## 3.7 Statistical considerations, Sample size and power

### Data analysis plan

- Descriptive Analysis:

Primarily, the PRISMA-Scr extension framework and checklist will be adhered to in traversing data analysis, with data reporting of eligible studies via the PRISMA flow diagram. (Appendix 2)

## 3.8 Ethical considerations

The protocol will be submitted to the Biomedical Research Ethics Committee of UKZN for approval prior to study initiation. No literature research will be conducted, collected or captured prior.

### 3.8.1 Informed consent

The scoping review will require no informed consent as there is no participation of individuals required for this study

### 3.8.2 Confidentiality

This study will pose no risk to patient confidentiality as private medical records are not within the scope of this review.

Data will be stored on the study PI's computer and will be backed up daily on the central server. Access to database records and data files will be restricted to the PI.

### 3.8.3 Conflict of interest

None declared

## 4. STUDY TIMELINE

This review will require extensive literature research throughout a planned 6 month period

## 5. STUDY LIMITATIONS

This study takes the form of a scoping review, and as such, is reliant solely on available and accessible literature.

## 6. FUTURE DIRECTIONS

The study findings will be able to inform surgeons locally of feasibility of the attractive focused parathyroidectomy based on objective costs and availability of the procedure requisites.

## 7. BUDGET

R500

Justification of budget

Coverage of data cost required to search the relevant library databases.

## 8. REFERENCES [1-25]

1. Ilahi, A., E. Muco, and T.B. Ilahi, *Anatomy, Head and Neck, Parathyroid*, in *In: StatPearls [Internet]*. . Jan 2022: Treasure Island (FL); StatPearls Publishing.
2. Saleem, M. and H. Iftikhar, *Stones, Bones, Groans, and Psychic Moans: Primary Hyperparathyroidism Presenting as Surgical Emergency*. *Cureus*, 2019. **11**(6): p. e4989.
3. Roe, S.M., et al., *Cost-effectiveness of preoperative localization studies in primary hyperparathyroid disease*. *Ann Surg*, 1994. **219**(5): p. 582-6.
4. Slepavicius, A., et al., *Focused versus conventional parathyroidectomy for primary hyperparathyroidism: a prospective, randomized, blinded trial*. *Langenbecks Arch Surg*, 2008. **393**(5): p. 659-66.
5. Kunstman, J.W. and R. Udelsman, *Superiority of Minimally Invasive Parathyroidectomy*. *Advances in Surgery*, 2012. **46**(1): p. 171-189.
6. Roth, S.I., C.-a. Wang, and J.T. Potts, *The team approach to primary hyperparathyroidism*. *Human Pathology*, 1975. **6**(6): p. 645-648.
7. Tibblin, S., A.G. Bondeson, and O. Ljungberg, *Unilateral parathyroidectomy in hyperparathyroidism due to single adenoma*. *Annals of surgery*, 1982. **195**(3): p. 245-252.
8. Aygün, N. and M. Uludağ, *Surgical Treatment of Primary Hyperparathyroidism: Which Therapy to Whom?* *Sisli Etfal Hastanesi tip bulteni*, 2019. **53**(3): p. 201-214.
9. Swanson, T.W., et al., *Determinants of Tc-99m sestamibi SPECT scan sensitivity in primary hyperparathyroidism*. *The American Journal of Surgery*, 2010. **199**(5): p. 614-620.
10. Tay, D., J.P. Das, and R. Yeh, *Preoperative Localization for Primary Hyperparathyroidism: A Clinical Review*. *Biomedicines*, 2021. **9**(4).
11. Barczynski, M., et al., *Intraoperative parathyroid hormone assay improves outcomes of minimally invasive parathyroidectomy mainly in patients with a presumed solitary parathyroid adenoma and missing concordance of preoperative imaging*. *Clinical Endocrinology*, 2007. **66**(6): p. 878-885.

12. Morris, M.A., et al., *Parathyroid Imaging: Past, Present, and Future*. *Frontiers in Endocrinology*, 2022. **12**.
13. Murphy, C. and J. Norman, *The 20% rule: a simple, instantaneous radioactivity measurement defines cure and allows elimination of frozen sections and hormone assays during parathyroidectomy*. *Surgery*, 1999. **126**(6): p. 1023-8; discussion 1028-9.
14. Fahy, B.N., et al., *Modern Parathyroid Surgery: A Cost-benefit Analysis of Localizing Strategies*. *Archives of Surgery*, 2002. **137**(8): p. 917-923.
15. Peabody, J.W., et al., *Improving the Quality of Care in Developing Countries*, in *Disease Control Priorities in Developing Countries*, nd, et al., Editors. 2006: Washington (DC).
16. Levac, D., H. Colquhoun, and K.K. O'Brien, *Scoping studies: advancing the methodology*. *Implement Sci*, 2010. **5**: p. 69.
17. Tricco, A.C., et al., *PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation*. *Ann Intern Med*, 2018. **169**(7): p. 467-473.
18. Page, M.J., et al., *The PRISMA 2020 statement: an updated guideline for reporting systematic reviews*. *BMJ*, 2021. **372**: p. n71.
19. Glonti, K., et al., *A scoping review protocol on the roles and tasks of peer reviewers in the manuscript review process in biomedical journals*. *BMJ Open*, 2017. **7**(10): p. e017468.
20. Ma, L.L., et al., *Methodological quality (risk of bias) assessment tools for primary and secondary medical studies: what are they and which is better?* *Mil Med Res*, 2020. **7**(1): p. 7.
21. Aromataris, E., et al., *Summarizing systematic reviews: methodological development, conduct and reporting of an umbrella review approach*. *Int J Evid Based Healthc*, 2015. **13**(3): p. 132-40.
22. Prager, G., et al., *Minimally invasive open parathyroidectomy A review*. *Acta Chirurgica Austriaca*, 2007. **31**: p. 221-226.
23. Nafisi Moghadam, R., et al., *Comparative Diagnostic Performance of Ultrasonography and 99mTc-Sestamibi Scintigraphy for Parathyroid Adenoma in Primary Hyperparathyroidism; Systematic Review and Meta- Analysis*. *Asian Pac J Cancer Prev*, 2017. **18**(12): p. 3195-3200.
24. Denham, D.W. and J. Norman, *Cost-effectiveness of preoperative sestamibi scan for primary hyperparathyroidism is dependent solely upon the surgeon's choice of operative procedure*. *J Am Coll Surg*, 1998. **186**(3): p. 293-305.
25. Lubitz, C.C., et al., *Preoperative localization strategies for primary hyperparathyroidism: an economic analysis*. *Ann Surg Oncol*, 2012. **19**(13): p. 4202-9.

## 1. Protocol Appendices

Protocol Appendix 1

# JBI Critical Appraisal Checklist for Systematic Reviews and Research Syntheses

Reviewer \_\_\_\_\_ Date \_\_\_\_\_  
Author \_\_\_\_\_ Year \_\_\_\_\_ Record Number \_\_\_\_\_


	Yes	No	Unclear	Not applicable
1. Is the review question clearly and explicitly stated?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Were the inclusion criteria appropriate for the review question?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Was the search strategy appropriate?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Were the sources and resources used to search for studies adequate?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Were the criteria for appraising studies appropriate?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Was critical appraisal conducted by two or more reviewers independently?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. Were there methods to minimize errors in data extraction?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. Were the methods used to combine studies appropriate?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. Was the likelihood of publication bias assessed?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
10. Were recommendations for policy and/or practice supported by the reported data?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
11. Were the specific directives for new research appropriate?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Overall appraisal:      Include       Exclude       Seek further info

---

## Protocol Appendix 2

PRISMA 2020 flow diagram for new systematic reviews which included searches of databases, registers and other sources



\*Consider, if feasible to do so, reporting the number of records identified from each database or register searched (rather than the total number across all databases/registers).

\*\*If automation tools were used, indicate how many records were excluded by a human and how many were excluded by automation tools.

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71. For more information, visit: <http://www.prisma-statement.org/>

## Supervisor-Student Memorandum of Understanding

*Prepared by Prof MJ Chimbari*

This memorandum states the responsibilities of the supervisor(s) and postgraduate student and requires both parties to accept the responsibilities by signing.

### Details of Student, Supervisors, and Project

**Student Name:** .....**Kapil Rugnath**.....  
**Student Number:** .....**207503757**.....  
**School:** .....**College of Health Sciences - School of Medicine**.....  
**Degree:** .....**MMED**.....  
**Supervisor(s):** .....**Suman Mewa Kinoo**.....  
**Research Topic:** .....**Feasibility of Focused Parathyroidectomy for Primary**.....  
**Hyperparathyroidism In A Developing Country**.....  
**Date**.....**25 October 2022**.....

### Responsibilities of the Postgraduate Student

While there are many responsibilities carried by a student in pursuing postgraduate studies the following are the minimum expected.

1. Student should identify a research topic acceptable to the supervisor in order to register
2. Student must show commitment to the degree programme and undertake to produce a full proposal within 3 month of registering
3. Student must produce written work that is their best effort for comments by the supervisor
4. Student should meet at least once per month (in person or through skype) with the supervisor and have the courage to request for such meetings. In all such meetings the student should provide a brief report of their work and take minutes of the discussions and retain such records until the degree has been awarded
5. Students must keep a laboratory manual where all experimental procedures and data are recorded. This laboratory manual remains the property of the university
6. Student must demonstrate the highest level of scientific honesty at all *stages (proposal writing, seeking ethical approval, collecting data, analyzing data and writing thesis or manuscripts)* of the degree programme.
7. Students must familiarize themselves with the university's policy on Plagiarism
8. Students should follow the advice provided by the supervisor and if they choose not to they should discuss the matter with the supervisor immediately
9. Student must always inform the supervisor of their whereabouts
10. Student should keep up to date with literature in their field of study and share any new literature they come across with the supervisor
11. Student must agree to complete studies within the time specified in the CHS handbook for the specific degree programme
12. Student should allow the supervisor to publish their work if they do not do so or show interest one year after graduating on the understanding that the student will be co-author


## Responsibilities of the Supervisor

1. Supervisor must support student at all stages of the degree programme (*settling down, proposal writing, ethical applications, data collection, data analysis and write up of thesis or manuscripts*)
2. Supervisor must be sensitive to the overall well-being of the student
3. Supervisor must have good knowledge of the research area of the student
4. Supervisor must be available to the student and should have regular meetings (face to face or by skype) with the student. If the supervisor must be away for an extended period they should identify a co-supervisor to assist the student during that period
5. Supervisor must read work submitted by student for comments and give feedback within 3 weeks depending on the nature of the work submitted
6. Supervisor must be constructively critical to the student's work
7. Supervisor must have sufficient interest in the work of the student
8. In instances of co-supervision the supervisors must avoid confusing the student by giving conflicting opinions/comments. If there are differences in opinion those should be discussed among the supervisors and the student given the agreed opinion.
9. Supervisor should, where funds permit, facilitate arrangements for masters and doctoral students to present a paper or a poster at an international conference as part of training
10. Supervisor must provide an annual progress report on the research and progression of the student to the discipline
11. Supervisor must protect the work of the student by not pre-maturely publishing it or assigning another student to similar work
12. Student must always be the first author of their work and any co-authorship with other people not on the supervision team should be clarified at an early stage of the project

## Conflict Resolution

Should there be a conflict or disagreement between supervisor and student which cannot be resolved by the parties involved, then either party can approach the Academic Leader Research or Dean and Head of School (or the College Dean of Research if the Dean and Head of School is one of the conflicting parties) about the conflict. The Dean and Head of School (or College Dean of Research) will then either arbitrate or choose a senior academic of the School not involved in the conflict to arbitrate. The arbitrator's decision is final and cannot be appealed.

## Signatures:

*Student*.....

*Supervisor*.....

*Co-Supervisor(s)*.....

*Academic Leader Research or D&HoS*.....



# Zertifikat Certificat

# Certificado Certificate

Promouvoir les plus hauts standards éthiques dans la protection des participants à la recherche biomédicale  
Promoting the highest ethical standards in the protection of biomedical research participants

## Certificat de formation - Training Certificate

Ce document atteste que - this document certifies that

**Suman Mewa-Kinoo**

a complété avec succès - has successfully completed

**Introduction to Research Ethics**

du programme de formation TRREE en évaluation éthique de la recherche  
of the TRREE training programme in research ethics evaluation



Release Date: 2022/09/23

CID : 5z4p1mTvq3

Professeur Dominique Sprumont  
Coordinateur TRREE Coordinator



Programmes de formation continue (2 crédits)  
Continuing Education Programs (2 credits)

Foederatio  
Pharmaceutica  
Helvetiae

**FPH**

Programmes de formation continue  
Continuing Education Programs

Programmes de formation  
postgraduée et continue

Ce programme est soutenu par - This program is supported by :

European and Developing Countries Clinical Trials Partnership (EDCTP) ([www.edctp.org](http://www.edctp.org)) - Swiss National Science Foundation ([www.snf.ch](http://www.snf.ch)) - Canadian Institutes of Health Research (<http://www.cihr-irsc.gc.ca/e/2891.html>) - Swiss Academy of Medical Science (SAMS/ASSM/SAMW) ([www.samw.ch](http://www.samw.ch)) - Commission for Research Partnerships with Developing Countries ([www.kfpe.ch](http://www.kfpe.ch))



# Zertifikat Certificat

# Certificado Certificate

Promouvoir les plus hauts standards éthiques dans la protection des participants à la recherche biomédicale  
Promoting the highest ethical standards in the protection of biomedical research participants

## Certificat de formation - Training Certificate

Ce document atteste que - this document certifies that

**Suman Mewa-Kinoo**

a complété avec succès - has successfully completed

**South Africa**

du programme de formation TRREE en évaluation éthique de la recherche  
of the TRREE training programme in research ethics evaluation



Release Date: 2022/09/23

CID : RXgRzUFsob

Professeur Dominique Sprumont  
Coordinateur TRREE Coordinator



Programmes de formation continue (2 crédits)  
Continuing Education Programs (2 credits)

Foederatio  
Pharmaceutica  
Helvetiae

**FPH**

Programmes de formation continue  
Continuing Education Programms

Programmes de formation  
postgraduée et continue

Ce programme est soutenu par - This program is supported by :

European and Developing Countries Clinical Trials Partnership (EDCTP) ([www.edctp.org](http://www.edctp.org)) - Swiss National Science Foundation ([www.snf.ch](http://www.snf.ch)) - Canadian Institutes of Health Research (<http://www.cihr-irsc.gc.ca/e/2891.html>) - Swiss Academy of Medical Science (SAMS/ASSM/SAMW) ([www.samw.ch](http://www.samw.ch)) - Commission for Research Partnerships with Developing Countries ([www.kfpe.ch](http://www.kfpe.ch))



# Zertifikat Certificat

# Certificado Certificate

Promouvoir les plus hauts standards éthiques dans la protection des participants à la recherche biomédicale  
Promoting the highest ethical standards in the protection of biomedical research participants

## Certificat de formation - Training Certificate

Ce document atteste que - this document certifies that

**kapil rugnath**

a complété avec succès - has successfully completed

**South Africa**

du programme de formation TRREE en évaluation éthique de la recherche  
of the TRREE training programme in research ethics evaluation



Release Date: 2022/10/19

CID : JIQhR9Stl

Professeur Dominique Sprumont  
Coordinateur TRREE Coordinator



Programmes de formation continue (2 crédits)  
Continuing Education Programs (2 credits)

Foederatio  
Pharmaceutica  
Helvetiae

**FPH**

Programmes de formation continue  
Continuing Education Programms

Programmes de formation  
postgraduée et continue

Ce programme est soutenu par - This program is supported by :

European and Developing Countries Clinical Trials Partnership (EDCTP) ([www.edctp.org](http://www.edctp.org)) - Swiss National Science Foundation ([www.snf.ch](http://www.snf.ch)) - Canadian Institutes of Health Research (<http://www.cihr-irsc.gc.ca/e/2891.html>) - Swiss Academy of Medical Science (SAMS/ASSM/SAMW) ([www.samw.ch](http://www.samw.ch)) - Commission for Research Partnerships with Developing Countries ([www.kfpe.ch](http://www.kfpe.ch))



# Zertifikat Certificat

# Certificado Certificate

Promouvoir les plus hauts standards éthiques dans la protection des participants à la recherche biomédicale  
Promoting the highest ethical standards in the protection of biomedical research participants

## Certificat de formation - Training Certificate

Ce document atteste que - this document certifies that

**kapil rugnath**

a complété avec succès - has successfully completed

**Introduction to Research Ethics**

du programme de formation TRREE en évaluation éthique de la recherche  
of the TRREE training programme in research ethics evaluation



Release Date: 2022/10/19

CID : 8QumTbPxLA

Professeur Dominique Sprumont  
Coordinateur TRREE Coordinator



Programmes de formation continue (2 crédits)  
Continuing Education Programs (2 credits)

Foederatio  
Pharmaceutica  
Helvetiae

**FPH**

Programmes de formation continue  
Continuing Education Programs

Programmes de formation  
postgraduée et continue

Ce programme est soutenu par - This program is supported by :

European and Developing Countries Clinical Trials Partnership (EDCTP) ([www.edctp.org](http://www.edctp.org)) - Swiss National Science Foundation ([www.snf.ch](http://www.snf.ch)) - Canadian Institutes of Health Research (<http://www.cihr-irsc.gc.ca/e/2891.html>) - Swiss Academy of Medical Science (SAMS/ASSM/SAMW) ([www.samw.ch](http://www.samw.ch)) - Commission for Research Partnerships with Developing Countries ([www.kfpe.ch](http://www.kfpe.ch))

## Appendix 2: The Guidelines for Authorship for the Journal selected for submission of the manuscript

### South African Journal of Surgery (SAJS)

#### Author Guidelines

Submitted manuscripts that are not in the correct format and without the required supporting documentation specified in these guidelines will be returned to the author(s) for correction and will delay publication.

#### AUTHORSHIP

Named authors must consent to publication by signing a covering letter which should be submitted as a supplementary file. Authorship should be based on substantial contribution to:

- (i) conception, design, analysis and interpretation of data;
- (ii) drafting or critical revision for important intellectual content; and
- (iii) approval of the version to be published. These conditions must all be met (uniform requirements for manuscripts submitted to biomedical journals; refer to [www.icmje.org](http://www.icmje.org)); and
- (iv) exact contribution of each author must be stated.

#### DECLARATION OF CONFLICT OF INTEREST

Authors must declare all sources of support for the research and any association with a product or subject that may constitute a conflict of interest. If there is no conflict of interest to declare please include the following: The authors declare no conflict of interest.

#### FUNDING SOURCE

All sources of funding should be declared. Also define the involvement of study sponsors in the study design, collection, analysis and interpretation of data; the writing of the manuscript; the decision to submit the manuscript for publication. If the study sponsors had no such involvement, this should be stated as follows: No funding source to be declared.

#### RESEARCH ETHICS COMMITTEE APPROVAL

The submitting author must provide written confirmation of Research Ethics Committee approval for all studies including case reports. The ethics committee as well as the approval number should be included. Please provide the Ethics Committee approval letter.

#### STATISTICAL ANALYSIS

Authors are advised to involve medical statisticians at the protocol stage of their research project: to plan sample size, and the selection of appropriate statistical tests for analysis and presentation.

#### PROTECTION OF PATIENT'S RIGHTS TO PRIVACY

Identifying information should not be published in written descriptions, photographs, and pedigrees unless the information is essential for scientific purposes and the patient (or parent or guardian) gives informed written consent for publication. The patient should be shown the manuscript to be published. Refer to [www.icmje.org](http://www.icmje.org).

#### ETHNIC CLASSIFICATION

The rationale for analysis based on racio-ethnic-cultural categorisation should be indicated.

## **CATEGORIES OF SUBMISSIONS**

Shorter items are more likely to be accepted for publication, owing to space constraints and reader preferences.

### ***Original articles***

Original articles on research relevant to surgery should not exceed 3 000 words, no more than 30 references, with up to 6 tables or figures. A structured abstract under the following headings, Background, Methods, Results, and Conclusions is a requirement and should not exceed 250 words.

### ***Scientific letters/short reports***

Short reports should not exceed 1 500 words with a maximum of 10 references. Only one table or illustration is permissible. A structured abstract under the following headings, Background, Methods, Results, and Conclusions, is a requirement and should not exceed 250 words.

### ***Case reports***

Case reports should not exceed 1 500 words with no more than 10 references. Figures are limited to 2 figures and may include images or photographs. The case report should have three headings: Summary (not exceeding 100 words), Case report (with no introduction) and Discussion. Case reports will be published online only. The summary and the URL will appear in the printed version.

### ***Video case reports (SAJS-VIDEO)***

Video case reports should not exceed 1 500 words with 10 references and 6 figures. Heading should include Summary (not exceeding 100 words) and Case description (with three subheadings: Introduction, Case presentation and Discussion). The video file format must be only MP4 or MOV and should not exceed 300 MB and 8 minutes. Video case reports will be published online only. The summary and the URL will appear in the printed version.

### ***How to do it***

How to do it submissions should address a practical aspect of surgical or interventional (endoscopic or radiological) patient management in which a best practice technique or method to advance optimal patient management is presented in a standardised format. The submission should be structured with a short contextual introduction focussing on the indications for the procedure, followed by numbered sequential points that explain and illustrate the procedure and its complications. The total word count should not exceed 1500 words with a maximum of 10 references and 6 figures. Five keywords should be included.

### ***Editorials***

Opinions, etc. should not exceed 1 000 words and are welcome, but unless invited, will be subjected to the SAJS peer review process.

### ***Review articles***

Review articles relevant to surgery should not exceed 5 000 words, with a maximum of 50 references and no more than 6 tables or figures. A summary of 250 words or less is required.

### ***Letters to the editor***

Letters to the editor should be 400 words or less with only one image or table.

### ***Obituaries***

Obituaries should be 900 words or less and should be accompanied by a photograph.

## MANUSCRIPT PREPARATION

Refer to articles in recent issues for the presentation of headings and subheadings. If in doubt, refer to 'uniform requirements' - [www.icmje.org](http://www.icmje.org). Manuscripts must be provided in **UK English**.

The manuscript should contain the title, abstract, keywords, article text and references.

The title page should be submitted as a supplementary file and should include:

- Qualification, affiliation and contact details of ALL the authors.
- Email addresses of all author must be provided.
- ORCID number of ALL authors must be provided – if authors do not have ORCID, please register at <https://orcid.org/>.
- Disclaimers: Acknowledgements, Declaration of conflict of interest, Funding declaration, Ethics declaration and ORCID.
- *A signed copy of the title page including the declarations must be provided in PDF format. An unsigned copy of the title page MUST be submitted in MSWord format.*

## Abbreviations

All abbreviations should be spelt out when first used and thereafter used consistently, e.g. 'intravenous (IV)' or 'Department of Health (DoH)'.

## Scientific measurements

Scientific measurements must be expressed in SI units except blood pressure (mmHg) and haemoglobin (g/dl). Litres is denoted with a lowercase 'l' e.g. 'ml' for millilitres). Units should be preceded by a space (except for %), e.g. '40 kg' and '20 cm' but '50%'. Greater/smaller than signs (> and 40 years of age) should also be preceded by a space e.g. > 20 years. No spaces should precede ± and °, i.e. '35±6' and '19°C'.

**Numbers** should be written as grouped per thousand-units, i.e. 4 000, 22 160...

**Quotes** should be placed in single quotation marks: i.e. The respondent stated: '...' Round **brackets** (parentheses) should be used, as opposed to square brackets, which are reserved for denoting concentrations or insertions in direct quotes.

## General formatting

The manuscript must be in Microsoft Word. *Please DO NOT provide the manuscript in PDF format.* Text must be 1,5-spaced, in 12-point Times New Roman font, and contain no unnecessary formatting (such as text in boxes, except for Tables). *The manuscript must be free of track changes.*

## ILLUSTRATIONS AND TABLES

If tables or illustrations submitted have been published elsewhere, the author(s) should provide consent to republication obtained from the copyright holder.

**Tables** may be embedded in the manuscript file **and** provided as '**supplementary files**'. They must be numbered in Arabic numerals (1,2,3...) and referred to consecutively in the text (e.g. 'Table 1'). Tables should be constructed carefully and simply for intelligible data representation. Unnecessarily complicated tables are strongly discouraged. Tables must be cell-based (i.e. not constructed with text boxes, tabs or enters) and accompanied by a concise title and column headings. Footnotes must be indicated with consecutive use of the following symbols: \* † ‡ § ¶ || then \*\* †† ‡‡ etc.

**Figures** must be numbered in Arabic numerals and referred to in the text e.g. '(Figure 1)'. Figure legends: Figure 1: 'Title...'. All illustrations/figures/graphs must be of **high resolution/quality**: 300 dpi or more is preferable, but images must not be resized to increase resolution. Unformatted and uncompressed images must be attached as '**supplementary files**' upon submission (not embedded in the accompanying manuscript). TIFF and PNG formats are preferable; JPEG and PDF formats are accepted, but authors must be wary of image compression. Illustrations and graphs prepared in Microsoft PowerPoint or Excel must be accompanied by the original workbook.

## REFERENCES

Authors must verify references from the original sources. *Only complete, correctly formatted reference lists will be accepted.* Reference lists may be generated with the use of reference manager software, but the final document must be delinked from the reference database or otherwise generated manually. Citations should be inserted in the text as superscript, e.g. These regulations are endorsed by the World Health Organization,<sup>2</sup> and others.<sup>3,4-6</sup> The superscript reference number should come after the punctuation mark.

All references should be listed at the end of the article in numerical order of appearance in the **Vancouver style** (not alphabetical order). Approved abbreviations of journal titles must be used; see the List of Journals in Index Medicus. Names and initials of all authors should be given; if there are more than six authors, the first three names should be given followed by et al. First and last page, volume and issue numbers should be given. **Wherever possible, references must be accompanied by a digital object identifier (DOI) link and PubMed ID (PMID)/PubMed Central ID (PMCID).** Authors are encouraged to use the DOI lookup service offered by [CrossRef](#). Crossref DOIs should always be displayed as a full URL link in the form <https://doi.org/10.xxxx/xxxx>

**Journal references:** Price NC, Jacobs NN, Roberts DA, et al. Importance of asking about glaucoma. *Stat Med* 1998;289(1):350-355. [<http://dx.doi.org/10.1000/hgjr.182>] [PMID: 2764753]

**Book references:** Jeffcoate N. Principles of Gynaecology. 4th ed. London: Butterworth, 1975:96-101. *Chapter/section in a book:* Weinstein L, Swartz MN. Pathogenic Properties of Invading Microorganisms. In: Sodeman WA jun, Sodeman WA, eds. Pathologic Physiology: Mechanisms of Disease. Philadelphia: WB Saunders, 1974:457-472.

**Internet references:** World Health Organization. The World Health Report 2002 - Reducing Risks, Promoting Healthy Life. Geneva: World Health Organization, 2002. <http://www.who.int/whr/2002> (accessed 16 January 2010).

**Other references (e.g. reports)** should follow the same format: Author(s). Title. Publisher place: publisher name, year; pages. Cited manuscripts that have been accepted but not yet published can be included as references followed by '(in press)'. Unpublished observations and personal communications in the text must not appear in the reference list. The full name of the source person must be provided for personal communications e.g. '(Prof. Michael Jones, personal communication)';

## COVERING LETTER

A covering letter to the editor is mandatory and must include statements that the manuscript has not been published previously and is not under review elsewhere. It should state details of any prior publication of the research in abstract form or in Congress proceedings. The letter must declare if any of the authors have a conflict of interest and that the requirements for submission, including ethics approval and patient permission for case reports have been fulfilled. All authors must sign the covering letter.

Please provide the names and email addresses of three possible reviewers for this manuscript.

## REVIEW PROCESS

Manuscripts, after vetting by the editorial team, are assigned for peer-review to 3 reviewers, conversant with the particular field of research. The reviewers and the authors are blinded to each other's identity. The turn-around time for review and initial editorial decision notification aims to be within 6 weeks of submission.

## PROOFS

A PDF proof of an article may be sent to the corresponding author before publication to resolve remaining queries. At that stage, **only** typographical changes are permitted; the corresponding author is required, having conferred with his/her co-authors, to reply within 2 working days in order for the article to be published in the issue for which it has been scheduled.

#### **CHANGES OF ADDRESS**

Please notify the editorial department of any contact detail changes, including email, to facilitate communication.

#### **CHARGES**

There is no charge for the publication of manuscripts.

#### **COPYRIGHT NOTICE**

By submitting manuscripts to *SAJS*, authors of original articles are assigning copyright to the journal. Authors may use their own work after publication without written permission, provided they acknowledge the original source. Individuals and academic institutions may freely copy and distribute articles published in *SAJS* for educational and research purposes without obtaining permission.

The work is licensed under a [Creative Commons Attribution-Noncommercial Works 4.0 South Africa License](#). Material submitted for publication in the *SAJS* is accepted provided it has not been published elsewhere. The *SAJS* does not hold itself responsible for statements made by the authors.

#### **PRIVACY STATEMENT**

The *SAJS* is committed to protecting the privacy of the users of this journal website. The names, personal particulars and email addresses entered in this website will be used only for the stated purposes of this journal and will not be made available to third parties without the user's permission or due process. Users consent to receive communication from the *SAJS* for the stated purposes of the journal. Queries with regard to privacy may be directed to [robyn@jesser-point.co.za](mailto:robyn@jesser-point.co.za).

#### **Copyright Notice**

The *South African Journal of Surgery (SAJS)* reserves copyright of the material published. The work is licensed under a [Creative Commons Attribution-Noncommercial Works 4.0 South Africa License](#). Material submitted for publication in the *SAJS* is accepted provided it has not been published elsewhere. The *SAJS* does not hold itself responsible for statements made by the authors.

#### **Privacy Statement**

The *SAJS* is committed to protecting the privacy of the users of this journal website. The names, personal particulars and email addresses entered in this website will be used only for the stated purposes of this journal and will not be made available to third parties without the user's permission or due process. Users consent to receive communication from the *SAJS* for the stated purposes of the journal. Queries with regard to privacy may be directed to [robyn@jesser-point.co.za](mailto:robyn@jesser-point.co.za).

Online ISSN: 2078-5151 Print ISSN: 0038-2361

#### **COPYRIGHT NOTICE**

*South African Journal of Surgery (SAJS)* Copyright is held by Association of Surgeons of South Africa (ASSA). The work is licensed under a Creative Commons Attribution-Non-Commercial Works 4.0 South Africa License (CC-NC BY 4.0). Material submitted for publication in the *SAJS* is accepted provided it has not been published elsewhere. *SAJS* does not hold itself responsible for statements made by the authors. The opinions expressed in this publication are those of the authors. They do not purport to reflect the opinions or views of the ASSA or its members.

## Appendix 3: Ethical approvals



05 May 2023

Dr Kapil Rugnath (207503757)  
School of Clinical Medicine  
Medical School

Dear Dr Rugnath,

Protocol reference number: BREC/00005172/2023

Project title: Feasibility of focused parathyroidectomy for primary hyperparathyroidism in a developing country

Degree: MMed

### EXEMPTION LETTER

I refer to your application to BREC and wish to advise you that exemption from ethics review has been granted for this study.

This exemption will be **noted** at the next Biomedical Research Ethics Committee meeting to be held on **13 June 2023**.

Yours sincerely,



Prof D Wassenaar  
Chair: Biomedical Research Ethics Committee

---

Biomedical Research Ethics Committee  
Chair: Professor D R Wassenaar  
UKZN Research Ethics Office Westville Campus, Govan Mbeki Building  
Postal Address: Private Bag X54001, Durban 4000  
Email: [BREC@ukzn.ac.za](mailto:BREC@ukzn.ac.za)  
Website: <http://research.ukzn.ac.za/Research-Ethics/Biomedical-Research-Ethics.aspx>

Founding Campuses:  Edgewood  Howard College  Medical School  Pietermaritzburg  Westville

INSPIRING GREATNESS



**Appendix 5: Raw data – Articles Encountered Upon Search of The Literature**

1. Di Marco A, Mechera R, Glover A, Papachristos A, Clifton-Bligh R, Delbridge L, et al. Focused parathyroidectomy without intraoperative parathyroid hormone measurement in primary hyperparathyroidism: Still a valid approach? *Surgery*. 2021;170(5):1383-8.
2. Itani M, Middleton WD. Parathyroid Imaging. *Radiologic Clinics of North America*. 2020;58(6):1071-83.
3. Jha CK, Bichoo RA, Yadav SK, Sonthineni C, Bothra S. Ambulatory bilateral neck exploration for primary hyperparathyroidism: Is it safe? *The American Journal of Surgery*. 2017;213(6):1191-2.
4. Lalonde MN, Correia RD, Syktiotis GP, Schaefer N, Matter M, Prior JO. Parathyroid Imaging. *Seminars in Nuclear Medicine*. 2023.
5. Mohebbati A, Shaha AR. Imaging techniques in parathyroid surgery for primary hyperparathyroidism. *American Journal of Otolaryngology*. 2012;33(4):457-68.
6. Patel KR, Phillips DJ, Leibowitz JM, Scognamiglio T, Banuchi VE, Kuhel WI, et al. Value within otolaryngology: Assessment of the cost-utility analysis literature. *World Journal of Otorhinolaryngology-Head and Neck Surgery*. 2016;2(1):28-37.
7. Vidal O, Saavedra-Perez D, Vilaça J, Pantoja JP, Delgado-Oliver E, Lopez-Boado MA, et al. Minimally-invasive Endocrine Neck Surgery. *Cirugía Española (English Edition)*. 2019;97(6):305-13.
8. Walker JP. Status of the Rural Surgical Workforce. *Surgical Clinics of North America*. 2020;100(5):869-77.
9. Zammit M, Pierce K, Bailey L, Rowland M, Waghorn A, Shore S. Challenging NICE guidelines on parathyroid surgery. *The Surgeon*. 2022;20(4):e105-e11.
10. Di Marco A, Mechera R, Glover A, Papachristos A, Clifton-Bligh R, Delbridge L, et al. Focused parathyroidectomy without intraoperative parathyroid hormone measurement in primary hyperparathyroidism: Still a valid approach? *Surgery*. 2021;170(5):1383-8.
11. Herb J, Staley BS, Roberson M, Strassle PD, Kim LT. Use and disparities in parathyroidectomy for symptomatic primary hyperparathyroidism in the Medicare population. *Surgery*. 2021;170(5):1376-82.
12. Jang S, Mandabach M, Aburjania Z, Balentine CJ, Chen H. Wide variation in cost of surgical care for parathyroidectomy: is there a need for standardization of practice? *Surgery*. 2018;163(3):638-42.
13. Kiernan CM, Wang T, Perrier ND, Grubbs EG, Solórzano CC. Bilateral Neck Exploration for Sporadic Primary Hyperparathyroidism: Use Patterns in 5,597 Patients Undergoing Parathyroidectomy in the Collaborative Endocrine Surgery Quality Improvement Program. *Journal of the American College of Surgeons*. 2019;228(4):652-9.
14. Kuzu F, Arpaç D, Cakmak GK, Emre AU, Elri T, Ilikhan SU, et al. Focused parathyroidectomy without intra-operative parathormone monitoring: The value of PTH assay in preoperative ultrasound guided fine needle aspiration washout. *Annals of Medicine and Surgery*. 2016;6:64-7.
15. Narayan R, Perkins RM, Berbano EP, Yuan CM, Neff RT, Sawyers ES, et al. Parathyroidectomy Versus Cinacalcet Hydrochloride–Based Medical Therapy in the Management of Hyperparathyroidism in ESRD: A Cost Utility Analysis. *American Journal of Kidney Diseases*. 2007;49(6):801-13.
16. Neves Mcd, Abrahão AR, Abrahão M, Rosano M, Rocha Lad, Machado HKAG, et al. Sestamibi scan in renal parathyroidectomy: a worthwhile preoperative exam? *Brazilian Journal of Otorhinolaryngology*. 2022;88(5):740-4.
17. Ramonell KM, Fazendin J, Lovell K, Iyer P, Chen H, Lindeman B, et al. Outpatient parathyroidectomy in the pediatric population: An 18-year experience. *Journal of Pediatric Surgery*. 2022;57(3):410-3.
18. Werenski HE, Nguyen CJ, Johansson ED, Bunch PM, Randle RW. Value of Old Imaging for Patients Undergoing Parathyroidectomy for Primary Hyperparathyroidism. *Journal of Surgical Research*. 2023;282:147-54.
19. Calva-Cerqueira D, Smith BJ, Hostetler ML, Lal G, Menda Y, O'Dorisio TM, et al. Minimally Invasive Parathyroidectomy and Preoperative MIBI Scans: Correlation of Gland Weight and Preoperative PTH. *Journal of the American College of Surgeons*. 2007;205(4, Supplement):S38-S44.
20. Deutmeyer C, Weingarten M, Doyle M, Carneiro-Pla D. Case series of targeted parathyroidectomy with surgeon-performed ultrasonography as the only preoperative imaging study. *Surgery*. 2011;150(6):1153-60.
21. Elaraj D, Sturgeon C. Operative Treatment of Primary Hyperparathyroidism: Balancing Cost-effectiveness with Successful Outcomes. *Surgical Clinics of North America*. 2014;94(3):607-23.
22. Fraser S. Surgical management of parathyroid disease. *Surgery (Oxford)*. 2020;38(12):807-13.
23. Hanba C, Bobian M, Svider PF, Sheyn A, Siegel B, Lin H-S, et al. Perioperative considerations and complications in pediatric parathyroidectomy. *International Journal of Pediatric Otorhinolaryngology*. 2016;91:94-9.
24. Harari A, Allendorf J, Shifrin A, DiGorgi M, Inabnet WB. Negative preoperative localization leads to greater resource use in the era of minimally invasive parathyroidectomy. *The American Journal of Surgery*. 2009;197(6):769-73.
25. Iwata AJ, Wertz AS, Alluri S, Singer MC. A faster parathyroidectomy: Techniques to shorten non-surgical operating room time. *American Journal of Otolaryngology*. 2019;40(6):102292.
26. Kiernan CM, Schlegel C, Isom C, Kavalukas S, Peters MF, Solórzano CC. Ambulatory bilateral neck exploration for primary hyperparathyroidism: is it safe? *The American Journal of Surgery*. 2016;212(4):722-7.
27. Kunstman JW, Udelsman R. Superiority of Minimally Invasive Parathyroidectomy. *Advances in Surgery*. 2012;46(1):171-89.
28. Laird AM, Libutti SK. Minimally Invasive Parathyroidectomy Versus Bilateral Neck Exploration for Primary Hyperparathyroidism. *Surgical Oncology Clinics of North America*. 2016;25(1):103-18.
29. Lalonde MN, Correia RD, Syktiotis GP, Schaefer N, Matter M, Prior JO. Parathyroid Imaging. *Seminars in Nuclear Medicine*. 2023.
30. Lew JJ, Solorzano CC. Surgical Management of Primary Hyperparathyroidism: State of the Art. *Surgical Clinics of North America*. 2009;89(5):1205-25.
31. Li J, Vasilyeva E, Hiebert J, Britton H, Walker B, Wiseman SM. Limited clinical utility of intraoperative frozen section during parathyroidectomy for treatment of primary hyperparathyroidism. *The American Journal of Surgery*. 2019;217(5):893-8.
32. Medas F, Cappellacci F, Canu GL, Noordzij JP, Erdas E, Calò PG. The role of Rapid Intraoperative Parathyroid Hormone (ioPTH) assay in determining outcome of parathyroidectomy in primary hyperparathyroidism: A systematic review and meta-analysis. *International Journal of Surgery*. 2021;92:106042.
33. Norman J, Politz D, Lopez J. Reply: Unilateral Parathyroidectomy: Delayed Treatment with Higher Failures. *Journal of the American College of Surgeons*. 2012;215(2):297-300.
34. Ozderya A, Temizkan S, Cetin K, Ozugur S, Gul AE, Aydin K. The Results of Parathyroid Hormone Assay in Parathyroid Aspirates in Pre-Operative Localization of Parathyroid Adenomas for Focused Parathyroidectomy in Patients With Negative or Suspicious Technetium-99m-Sestamibi Scans. *Endocrine Practice*. 2017;23(9):1101-6.
35. Paravastu SCV, Chadwick DR. Parathyroidectomy in a district general hospital: Outcomes and evolution in the era of minimally invasive surgery. *International Journal of Surgery*. 2012;10(7):373-7.
36. Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *American Journal of Otolaryngology*. 2021;42(1):102819.
37. Rajeev P, Stechman MJ, Kirk H, Gleeson FV, Mihai R, Sadler GP. Safety and efficacy of minimally-invasive parathyroidectomy (MIP) under local anaesthesia without intra-operative PTH measurement. *International Journal of Surgery*. 2013;11(3):275-7.
38. Schneider DF, Mazeh H, Sippel RS, Chen H. Is minimally invasive parathyroidectomy associated with greater recurrence compared to bilateral exploration? Analysis of more than 1,000 cases. *Surgery*. 2012;152(6):1008-15.
39. Shah-Becker S, Goldenberg D. Minimally invasive parathyroidectomy. *Operative Techniques in Otolaryngology-Head and Neck Surgery*. 2016;27(3):152-6.
40. Shirali AS, Clemente-Gutierrez U, Perrier ND. Parathyroid Surgery: What Radiologists Need to Know. *Neuroimaging Clinics of North America*. 2021;31(3):397-408.
41. Solorzano CC, Carneiro-Pla D. Minimizing Cost and Maximizing Success in the Preoperative Localization Strategy for Primary Hyperparathyroidism. *Surgical Clinics of North America*. 2014;94(3):587-605.
42. Stalberg P, Delbridge L, van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy — current concepts. *The Surgeon*. 2007;5(5):301-8.
43. Stojadinovic A, Pribitkin E, Rosen D, Edwards M, Byrd DR. Unilateral vs Bilateral Parathyroidectomy: A Healthy Debate. *Journal of the American College of Surgeons*. 2012;215(2):300-2.
44. Wang TS, Cheung K, Farrakhyar F, Roman SA, Sosa JA. Would scan, but which scan? A cost-utility analysis to optimize preoperative imaging for primary hyperparathyroidism. *Surgery*. 2011;150(6):1286-94.
45. Zammit M, Pierce K, Bailey L, Rowland M, Waghorn A, Shore S. Challenging NICE guidelines on parathyroid surgery. *The Surgeon*. 2022;20(4):e105-e11.
46. Zhu CY, Nguyen DT, Yeh MW. Who Benefits from Treatment of Primary Hyperparathyroidism? *Surgical Clinics of North America*. 2019;99(4):667-79.
47. Bedi HK, Jedrzejko N, Nguyen A, Aspinall SR, Wiseman SM. Thyroid and parathyroid surgeon case volume influences patient outcomes: A systematic review. *Surgical Oncology*. 2021;38:101550.
48. Blanco-Saiz I, Goñi-Gironés E, Ribelles-Segura MJ, Salvador-Egea P, Díaz-Tobarra M, Camarero-Salazar A, et al. Preoperative parathyroid localization. Relevance of MIBI SPECT-CT in adverse scenarios. *Endocrinología, Diabetes y Nutrición*. 2022.
49. Bobanga ID, McHenry CR. Is intraoperative parathyroid hormone monitoring necessary for primary hyperparathyroidism with concordant preoperative imaging? *The American Journal of Surgery*. 2017;213(3):484-8.
50. Choi J-H, Jayaram A, Bresnahan E, Pletcher E, Steinmetz D, Owen R, et al. The Role of Surgeon-Performed Office and Preincision Ultrasounds in Localization of Parathyroid Adenomas in Primary Hyperparathyroidism. *Endocrine Practice*. 2022;28(7):660-6.
51. Douthwaite SA, Young JE, Pasternak JD, Yoo J. Surgical Management of Primary Hyperparathyroidism. *Journal of Clinical Densitometry*. 2013;16(1):48-53.
52. Garner EF, Chen H. Parathyroid Surgery☆. In: Huhtaniemi I, Martini L, editors. *Encyclopedia of Endocrine Diseases (Second Edition)*. Oxford: Academic Press; 2017. p. 151-9.

53. Graves CE, Duh Q-Y, Suh I. Innovations in Parathyroid Localization Imaging. *Surgical Oncology Clinics of North America*. 2022;31(4):631-47.
54. Lee L, Steward DL. Techniques for Parathyroid Localization with Ultrasound. *Ultrasound Clinics*. 2012;7(2):211-8.
55. Lo C-Y, Lang BH, Chan WF, Kung AWC, Lam KSL. A prospective evaluation of preoperative localization by technetium-99m sestamibi scintigraphy and ultrasonography in primary hyperparathyroidism. *The American Journal of Surgery*. 2007;193(2):155-9.
56. Lubitz CC, Duh Q-Y. 54 - Guide to Preoperative Parathyroid Localization Testing. In: Randolph GW, editor. *Surgery of the Thyroid and Parathyroid Glands (Third Edition)*; Elsevier; 2021. p. 494-501.e3.
57. Nguyen-Lee JJ, Paciuc M, Guerra R, Esnaola N, Zanocco KA, Zheng F. Derivation of a cost-saving screening strategy for asymptomatic primary hyperparathyroidism. *Surgery*. 2020;167(1):155-9.
58. Richards MK, Slavin ER, Tamarkin SW, McHenry CR. Technetium-99m sestamibi imaging: are the results dependent on the reviewer? *Journal of Surgical Research*. 2012;177(1):97-101.
59. Singh VP, Mir R. Present Trends in Parathyroid Surgery. *Apollo Medicine*. 2009;6(4):327-34.
60. Van Udelsman B, Udelsman R. Surgery in Primary Hyperparathyroidism: Extensive Personal Experience. *Journal of Clinical Densitometry*. 2013;16(1):54-9.
61. Zarei A, Karthik S, Chowdhury FU, Patel CN, Scarsbrook AF, Vaidyanathan S. Multimodality imaging in primary hyperparathyroidism. *Clinical Radiology*. 2022;77(6):e401-e16.

#### Academic Search Complete

1. Hemmati H, Pourahmadi Y, Motamed B, Ashoobi MT, Kenarsari HE. Evaluation of the Consequences of Targeted or Focused Mini-Incision Parathyroidectomy in Patients With Primary Hyperparathyroidism. *Acta Medica Iranica*. 2022;60(12):742-8.
2. Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian Journal of Endocrinology & Metabolism*. 2019;23(3):347-52.
3. Sen S, Cherian A, Ramakant P, Reka K, Paul M, Abraham D. Focused parathyroidectomy under local anesthesia – A feasibility study. *Indian Journal of Endocrinology & Metabolism*. 2019;23(1):67-71.
4. El-hady HA, Radwan HS. Focused parathyroidectomy for single parathyroid adenoma: a clinical anecdote of 20 patients. *Electronic Physician*. 2018;10(6):6974-80.
5. Yadav S, Mishra S, Mishra A, Mayilvagnan S, Chand G, Agarwal G, et al. Surgical management of primary hyperparathyroidism in the era of focused parathyroidectomy: A study in tertiary referral centre of North India. *Indian Journal of Endocrinology & Metabolism*. 2019;23(4):468-72.
6. Yang Z, Guo M, Wu B, Zheng Q, Fan Y. Focused parathyroidectomy through an open-lateral approach for treating solitary parathyroid adenoma. *Surgical Practice*. 2015;19(4):160-5.
7. Hacıyanlı M, Genc H, Damburacı N, Oruk G, Tutuncuoğlu P, Erdogan N. Minimally invasive focused parathyroidectomy without using intraoperative parathyroid hormone monitoring or gamma probe. *Journal of Postgraduate Medicine*. 2009;55(4):242-6.
8. Chow TL, Choi CY, Lam SH. Focused parathyroidectomy without intra-operative parathyroid hormone monitoring for primary hyperparathyroidism: results in a low-volume hospital. *Journal of Laryngology & Otology*. 2015;129(8):788-94.
9. Slepavicius A, Beisa V, Janusonis V, Strupas K. Focused versus conventional parathyroidectomy for primary hyperparathyroidism: a prospective, randomized, blinded trial. *Langenbeck's Archives of Surgery*. 2008;393(5):659-66.
10. Guerin C, Lowery A, Gabriel S, Castinetti F, Philippon M, Vaillant-lombard J, et al. Preoperative imaging for focused parathyroidectomy: making a good strategy even better. *European Journal of Endocrinology*. 2015;172(5):519-26.
11. Rika M. Three-dimensional ultrasonography before minimally invasive focused parathyroidectomy: The importance of coronal images. *Surgery Today*. 2009;39(2):98-103.
12. Carling T, Udelsman R. Focused Approach to Parathyroidectomy. *World Journal of Surgery*. 2008;32(7):1512-7.
13. Soon PSH, Delbridge LW, Sywak MS, Barraclough BM, Edhouse P, Sidhu SB. Surgeon Performed Ultrasound Facilitates Minimally Invasive Parathyroidectomy by the Focused Lateral Mini-incision Approach. *World Journal of Surgery*. 2008;32(5):766-71.
14. Pang T, Stalberg P, Sidhu S, Sywak M, Wilkinson M, Reeve TS, et al. Minimally invasive parathyroidectomy using the lateral focused mini-incision technique without intraoperative parathyroid hormone monitoring. *British Journal of Surgery*. 2007;94(3):315-9.
15. Langusch CC, Norlen O, Titmuss A, Donoghue K, Holland AJA, Shun A, et al. Focused image-guided parathyroidectomy in the current management of primary hyperparathyroidism. *Archives of Disease in Childhood*. 2015;100(10):924-7.
16. B N, Y. P S, A P, P S, G S. Surgical Management of Primary Hyperparathyroidism: An Institutional Study on Surgical Quality Control. *Journal of Institute of Medicine Nepal (JIOMN)*. 2017;39(2):89-93.
17. Werenski HE, Nguyen CJ, Johansson ED, Bunch PM, Randle RW. Value of Old Imaging for Patients Undergoing Parathyroidectomy for Primary Hyperparathyroidism. *Journal of Surgical Research*. 2023;282:147-54.
18. Hye-Sun P, Namki H, Jong Ju J, Mijin Y, Yumie R. Update on Preoperative Parathyroid Localization in Primary Hyperparathyroidism. *Endocrinology & Metabolism*. 2022;37(5):744-55.
19. Lombardi CP, Raffaelli M, Traini E, Stasio E, Carrozza C, de Crea C, et al. Intraoperative PTH monitoring during parathyroidectomy: the need for stricter criteria to detect multiglandular disease. *Langenbeck's Archives of Surgery*. 2008;393(5):639-45.
20. McGill J, Sturgeon C, Kaplan SP, Chiu B, Kaplan EL, Angelos P. How Does the Operative Strategy for Primary Hyperparathyroidism Impact the Findings and Cure Rate? A Comparison of 800 Parathyroidectomies. *Journal of the American College of Surgeons*. 2008;207(2):246-9.
21. Utility of At-Excision Intraoperative PTH for Unilateral Focused Parathyroidectomy. *Journal of Surgical Research*. 2014;186(2):506-.
22. Naik AH, Wani MA, Wani KA, Laway BA, Malik AA, Shah ZA. Intraoperative Parathyroid Hormone Monitoring in Guiding Adequate Parathyroidectomy. *Indian Journal of Endocrinology & Metabolism*. 2018;22(3):410-6.
23. Kiernan CM, Wang T, Perrier ND, Grubbs EG, Solórzano CC. Bilateral Neck Exploration for Sporadic Primary Hyperparathyroidism: Use Patterns in 5,597 Patients Undergoing Parathyroidectomy in the Collaborative Endocrine Surgery Quality Improvement Program. *Journal of the American College of Surgeons*. 2019;228(4):652-9.
24. Philippon M, Guerin C, Taieb D, Vaillant J, Morange I, Brue T, et al. Bilateral neck exploration in patients with primary hyperparathyroidism and discordant imaging results: a single-centre study. *European Journal of Endocrinology*. 2014;170(5):719-25.
25. Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally Invasive Parathyroidectomy without Intraoperative PTH Performed after Positive Ultrasonography as the only Diagnostic Method in Patients with Primary Hyperparathyroidism. *World Journal of Surgery*. 2019;43(6):1525-31.
26. Bergenfelz AOJ, Jansson SKG, Wallin GK, Mårtensson HG, Rasmussen L, Eriksson HLO, et al. Impact of modern techniques on short-term outcome after surgery for primary hyperparathyroidism: a multicenter study comprising 2,708 patients. *Langenbeck's Archives of Surgery*. 2009;394(5):851-60.
27. Lee G, McKenzie T, Mullan B, Farley D, Thompson G, Richards M. A Multimodal Imaging Protocol, I/Tc-Sestamibi, SPECT, and SPECT/CT, in Primary Hyperparathyroidism Adds Limited Benefit for Preoperative Localization. *World Journal of Surgery*. 2016;40(3):589-94.
28. James B, Kaplan E, Grogan R, Angelos P. What's in a Name?: Providing Clarity in the Definition of Minimally Invasive Parathyroidectomy. *World Journal of Surgery*. 2015;39(4):975-80.
29. Miccoli P, Berti P, Materazzi G, Ambrosini CE, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTHa) during endoscopic parathyroidectomy: A prospective randomized trial. *Surgical Endoscopy*. 2008;22(2):398-400.
30. Karakas E, Schneider R, Rothmund M, Bartsch D, Schlosser K. Initial Surgery for Benign Primary Hyperparathyroidism: An Analysis of 1,300 Patients in a Teaching Hospital. *World Journal of Surgery*. 2014;38(8):2011-8.
31. Gupta A, Unawane A, Subhas G, Herschman BR, Silapaswan S, Kolachalam R, et al. Parathyroidectomies Using Intraoperative Parathormone Monitoring: When Should We Stop Measuring Intraoperative Parathormone Levels? *American Surgeon*. 2012;78(8):844-50.
32. Harari A, Allendorf J, Shifrin A, DiGorgi M, Inabnet WB. Negative preoperative localization leads to greater resource use in the era of minimally invasive parathyroidectomy. *American Journal of Surgery*. 2009;197(6):769-73.

#### Sabinet

1. 47th meeting of the Surgical Research Society of Southern Africa, 25-28 June 2019, University of Pretoria. *South African Journal of Surgery*. 2019;57(2):69-93.
2. Bert Myburgh Research Forum list of abstracts 2017. *Wits Journal of Clinical Medicine*. 2019;1(1):37-48.
3. Budge M, Conradie W, Beviss-Challinor K, Martin L, Conradie M, Coetzee A. Bone health in patients undergoing surgery for primary hyperparathyroidism at Tygerberg Hospital, Cape Town, South Africa. *Journal of Endocrinology, Metabolism and Diabetes in South Africa*. 2021;26(1):16-23.
4. MacRobert NA, Kruger D, Schamm M. A Retrospective Audit of Single Versus Multigland Disease in Primary Hyperparathyroidism at a Single Centre in South Africa. *Wits Journal of Clinical Medicine*. 2021;3(2):93-100.
5. Panieri E. What's new in endocrine surgery? *CME : Your SA Journal of CPD*. 2007;25(7):312-3.
6. Mitchell C, Madima NR, Louw L, Mahlobo F, Muganza RA, Bombil I. Sonar guided focused parathyroidectomy under cervical block. *South African Journal of Surgery*. 2018;56(2):30-3.
7. Mihai R, Simon D, Hellman P. Imaging for primary hyperparathyroidism—an evidence-based analysis. *Langenbeck's Archives of Surgery*. 2009;394(5):765-84.

#### Langenbeck's Archives of Surgery

1. Iacobone M, Scerrino G, Palazzo FF. Parathyroid surgery: an evidence-based volume—outcomes analysis. *Langenbeck's Archives of Surgery*. 2019;404(8):919-27.
2. Sitges-Serra A, Rosa P, Valero M, Membrilla E, Sancho JJ. Surgery for sporadic primary hyperparathyroidism: controversies and evidence-based approach. *Langenbeck's Archives of Surgery*. 2008;393(3):239-44.
3. Patel N, Whittet C, Zhao D, Rees J, Stechman MJ, Scott-Coombes DM. A 15-year experience: intraoperative parathyroid hormone assay for the management of primary hyperparathyroidism in a UK endocrine surgical unit. *Langenbeck's Archives of Surgery*. 2023;408(1):120.
4. Mihai R, Barczynski M, Iacobone M, Sitges-Serra A. Surgical strategy for sporadic primary hyperparathyroidism: an evidence-based approach to surgical strategy, patient selection, surgical access, and reoperations. *Langenbeck's Archives of Surgery*. 2009;394(5):785-98.
5. Lombardi CP, Raffaelli M, Traini E, Di Stasio E, Carrozza C, De Crea C, et al. Intraoperative PTH monitoring during parathyroidectomy: the need for stricter criteria to detect multiglandular disease. *Langenbeck's Archives of Surgery*. 2008;393(5):639-45.
6. Gimm O, Barczyński M, Mihai R, Raffaelli M. Training in endocrine surgery. *Langenbeck's Archives of Surgery*. 2019;404(8):929-44.

#### ProQuest

1. Sharanappa V, Mishra A, Bhatia V, Sabaretnam M, Chand G, Agarwal G, et al. Pediatric Primary Hyperparathyroidism: Experience in a Tertiary Care Referral Center in a Developing Country Over Three Decades. *World Journal of Surgery*. 2021;45(2):488-95.
2. Koman A. Primary Hyperparathyroidism: Nonclassical Symptoms and Benefits from Parathyroidectomy [Ph.D.]. Sweden: Karolinska Institutet (Sweden); 2021.
3. Yadav S, Mishra S, Mishra A, Mayilvagnan S, Chand G, Agarwal G, et al. Surgical management of primary hyperparathyroidism in the era of focused parathyroidectomy: A study in tertiary referral centre of North India. *Indian Journal of Endocrinology and Metabolism*. 2019;23(4):468-72.
4. Pradhan R, Gupta S, Agarwal A. Focused parathyroidectomy using accurate preoperative imaging and intraoperative PTH: Tertiary care experience. *Indian Journal of Endocrinology and Metabolism*. 2019;23(3):347-52.
5. Aygün N, Uludağ M. Surgical Treatment of Primary Hyperparathyroidism: Which Therapy to Whom? *Sisli Etfal Hastan Tip Bul*. 2019;53(3):201-14.
6. Al-Thani H, El-Matbouly M, Al-Sulaiti M, Asim M, Majzoub A, Tabeb A, et al. Management and outcomes of hyperparathyroidism: a case series from a single institution over two decades. *Therapeutics and Clinical Risk Management*. 2018;14:1337-45.
7. Yadav SK, Mishra SK, Mishra A, Mayilvagnan S, Chand G, Agarwal G, et al. Changing Profile of Primary Hyperparathyroidism Over Two and Half Decades: A Study in Tertiary Referral Center of North India. *World Journal of Surgery*. 2018;42(9):2732-7.
8. Jha CK, Raouef Ahmed B, Yadav SK, Chaitra S, Bothra S. Ambulatory bilateral neck exploration for primary hyperparathyroidism: Is it safe? *The American Journal of Surgery*. 2017;213(6):1191-2.
9. Khan AA, Hanley DA, Rizzoli R, Bollerslev J, Young JEM, Rejnmark L, et al. Primary hyperparathyroidism: review and recommendations on evaluation, diagnosis, and management. A Canadian and international consensus. *Osteoporosis International*. 2017;28(1):1-19.
10. Kaur P, Gattani R, Singhal A, Sarin D, Arora S, Mithal A. Impact of preoperative imaging on surgical approach for primary hyperparathyroidism: Data from single institution in India. *Indian Journal of Endocrinology and Metabolism*. 2016;20(5).
11. Paruk IM, Esterhuizen TM, Maharaj S, Pirie FJ, Motala AA. Characteristics, management and outcome of primary hyperparathyroidism in South Africa: a single-centre experience. *Postgraduate Medical Journal*. 2013;89(1057):626.
12. Agarwal S, Agarwal A, Chand G. The Necessity and Reliability of Intraoperative Parathyroid Hormone (PTH) Testing in Patients with Mild Hyperparathyroidism and PTH Levels in the Normal Range. *World Journal of Surgery*. 2012;36(2):483-5.
13. McLean T, Delbridge L. Comparison of Consumer Information on the Internet to the Current Evidence Base for Minimally Invasive Parathyroidectomy. *World Journal of Surgery*. 2010;34(6):1304-11.
14. Turkey K, Quan-Yang D, Kebebew E, Orlo Herrick C. Do patients undergoing parathyroidectomy for primary hyperparathyroidism in San Francisco, CA, and Bursa, Turkey, differ? *The American Journal of Surgery*. 2009;198(2):188-92.
15. Battistella E, Pomba L, Toniato R, Burei M, Gregianin M, Fernando SW, et al. Evolution of the Diagnosis and Treatment of Primary Hyperparathyroidism. *Journal of Clinical Medicine*. 2023;12(5):2057.
16. Pickering JM, Giles WH. Improving Intraoperative Parathyroid Hormone Lab Efficiency. *The American Surgeon*. 2022;88(5):915-21.
17. Mathew J, Arjunan R, Althaf S, Halkud R. Primary Hyperparathyroidism: Is Image Localization Alone Sufficient to Ensure Long-Term Cure in Unifocal Disease? *Cureus*. 2022;14(11).
18. Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *American Journal of Otolaryngology*. 2021;42(1).
19. Süleyman Özkan A, Adiyaman SC, Cevlik Ali Durubey Ç, Guray Durak Merih Güray D, Secil MS, Sevinc Ali İbrahim S. Intra-operative parathyroid hormone evaluation is superior to frozen section analysis in parathyroid surgery. *American Journal of Otolaryngology*. 2021;42(3).
20. Goldfarb M, Singer FR. Recent advances in the understanding and management of primary hyperparathyroidism. *F1000Research*. 2020;9.
21. Iwata AJ, Wertz AS, Alluri S, Singer MC. A faster parathyroidectomy: Techniques to shorten non-surgical operating room time. *American Journal of Otolaryngology*. 2019;40(6).
22. Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally Invasive Parathyroidectomy without Intraoperative PTH Performed after Positive Ultrasonography as the only Diagnostic Method in Patients with Primary Hyperparathyroidism. *World Journal of Surgery*. 2019;43(6):1525-31.
23. Pradhan R, Gupta S, Agarwal A. Focused parathyroidectomy using accurate preoperative imaging and intraoperative PTH: Tertiary care experience. *Indian Journal of Endocrinology and Metabolism*. 2019;23(3):347-52.
24. Li J, Vasilyeva E, Hiebert J, Britton H, Walker B, Wiseman SM. Limited clinical utility of intraoperative frozen section during parathyroidectomy for treatment of primary hyperparathyroidism. *The American Journal of Surgery*. 2019;217(5):893-8.
25. Walsh NJ, Sullivan BT, Duke WS, Terris DJ. Routine bilateral neck exploration and four-gland dissection remains unnecessary in modern parathyroid surgery. *Laryngoscope Investigative Otolaryngology*. 2019;4(1):188-92.
26. Aygün N, Uludağ M. Surgical Treatment of Primary Hyperparathyroidism: Which Therapy to Whom? *Sisli Etfal Hastan Tip Bul*. 2019;53(3):201-14.
27. Sen S, Cherian A, Ramakant P, Reka K, Paul M, Abraham D. Focused parathyroidectomy under local anesthesia – A feasibility study. *Indian Journal of Endocrinology and Metabolism*. 2019;23(1):67-71.
28. Ishii H, Mihai R, Watkinson JC, Kim DS. Systematic review of cure and recurrence rates following minimally invasive parathyroidectomy. *BJS Open*. 2018;2(6):364-70.
29. Ozderya AMD, Temizkan SMD, Cetin K, Ozugur SMD, Gul AEMD, Aydin KMD. THE RESULTS OF PARATHYROID HORMONE ASSAY IN PARATHYROID ASPIRATES IN PRE-OPERATIVE LOCALIZATION OF PARATHYROID ADENOMAS FOR FOCUSED PARATHYROIDECTOMY IN PATIENTS WITH NEGATIVE OR SUSPICIOUS TECHNETIUM-99M-SESTAMIBI SCANS. *Endocrine Practice*. 2017;23(9):1101-6.
30. Parnell KE, Oltmann SC. The surgical management of primary hyperparathyroidism: an updated review. *International Journal of Endocrine Oncology*. 2018;5(1).
31. Jinih MMBB, O'Connell EMBB, O'Leary DPMBBMP, Liew AMBBMMP, Redmond HPMFF. Focused Versus Bilateral Parathyroid Exploration for Primary Hyperparathyroidism: A Systematic Review and Meta-analysis. *Annals of Surgical Oncology*. 2017;24(7):1924-34.
32. Polistena AMDP, Lucchini RMD, Monacelli MMD, Triola RMD, Avenia SMD, Barillaro IMD, et al. Current Indications for Surgical Treatment of Primary Hyperparathyroidism in the Elderly. *The American Surgeon*. 2017;83(3):296-302.
33. Calò PG, Medas F, Loi G, Pisano G, Sorrenti S, Erdas E, et al. Parathyroidectomy for primary hyperparathyroidism in the elderly: experience of a single endocrine surgery center. *Aging Clinical and Experimental Research*, suppl 1. 2017;29:15-21.
34. Sharata AMD, Kelly TLBSC, Rozenfeld YMPH, Hammill CWMD, SchumanMd E, Carlisle JRMD, et al. Management of Primary Hyperparathyroidism: Can We Do Better? *The American Surgeon*. 2017;83(1):64-70.
35. Kiernan CM, Schlegel C, Isom C, Kavalukas S, Peters MF, Solórzano CC. Ambulatory bilateral neck exploration for primary hyperparathyroidism: is it safe? *The American Journal of Surgery*. 2016;212(4):722-7.
36. Singh Ospina NM, Rodriguez-gutierrez R, Maraka S, Espinosa De Ycaza AE, Jasim S, Castaneda-guarderas A, et al. Outcomes of Parathyroidectomy in Patients with Primary Hyperparathyroidism: A Systematic Review and Meta-analysis. *World Journal of Surgery*. 2016;40(10):2359-77.
37. Sun Moon K, Shu AD, Long J, Montez-Rath ME, Leonard MB, Norton JA, et al. Declining Rates of Inpatient Parathyroidectomy for Primary Hyperparathyroidism in the US. *PLoS One*. 2016;11(8).
38. Nair C, Babu M, Jacob P, Menon R, Mathew J. Is intraoperative parathyroid hormone monitoring necessary in symptomatic primary hyperparathyroidism with concordant imaging? *Indian Journal of Endocrinology and Metabolism*. 2016;20(4).
39. Mownah OA, Pafitanis G, Drake WM, Crinnion JN. Contemporary surgical treatment of primary hyperparathyroidism without intraoperative parathyroid hormone measurement. *Annals of the Royal College of Surgeons of England*. 2015;97(8):603-7.
40. Uslukaya O, Gumus M, Tasdemir B, Goya C, Kilinc F, Oguz A, et al. Improvement of minimally invasive parathyroidectomy outcomes by real time ultrasonography performed by a surgeon and radiologist team. *Medical Ultrasonography*. 2015;17(3):315-21.
41. Chow TL, Choi CY, Lam SH. Focused parathyroidectomy without intra-operative parathyroid hormone monitoring for primary hyperparathyroidism: results in a low-volume hospital. *The Journal of Laryngology and Otolaryngology*. 2015;129(8):788-94.
42. Habib Z, Kabaker A, Camacho P. Sporadic primary hyperparathyroidism: medical and surgical updates. *Expert Review of Endocrinology & Metabolism*. 2014;9(1):31-44.
43. Tee MCMDDMPH, Chan SKM, Nguyen VMD, Strugnell SSM, Yang JB, Jones SMD, et al. Incremental value and clinical impact of neck sonography for primary hyperparathyroidism: a risk-adjusted analysis. *Canadian Journal of Surgery*. 2013;56(5):325-31.
44. Opoku-Boateng AMD, Bolton JSMD, Corsetti RMD, Brown REM, Oxner CMD, Fuhrman GMMD. Use of a Sestamibi-only Approach to Routine Minimally Invasive Parathyroidectomy. *The American Surgeon*. 2013;79(8):797-801.

45. Calò PG, Pisano G, Loi G, Medas F, Barca L, Atzeni M, et al. Intraoperative parathyroid hormone assay during focused parathyroidectomy: the importance of 20 minutes measurement. *BMC Surgery*. 2013;13:36.
46. Karahan Ö, Okus A, Sevinç B, Eryılmaz M, Ay S, Çaycı M, et al. Minimally invasive parathyroidectomy under local anesthesia. *Journal of Postgraduate Medicine*. 2013;59(1):21-4.
47. Lubitz CCMDMPH, Stephen AEMD, Hodin RAMD, Pandharipande PMDMPH. Preoperative Localization Strategies for Primary Hyperparathyroidism: An Economic Analysis. *Annals of Surgical Oncology*. 2012;19(13):4202-9.
48. Zia SMD, Sippel RSMD, Chen HMD. Sestamibi Imaging for Primary Hyperparathyroidism: The Impact of Surgeon Interpretation and Radiologist Volume. *Annals of Surgical Oncology*. 2012;19(12):3827-31.
49. Gupta AMD, Unawane AMD, Subhas GMD, Herschman BRMD, Silapaswan SMD, Kolachalam RMD, et al. Parathyroidectomies Using Intraoperative Parathormone Monitoring: When Should We Stop Measuring Intraoperative Parathormone Levels? *The American Surgeon*. 2012;78(8):844-50.
50. Mohebbati A, Shaha AR. Imaging techniques in parathyroid surgery for primary hyperparathyroidism. *American Journal of Otolaryngology*. 2012;33(4):457-68.
51. Twigt BA, van Dalen T, Vollebregt AM, Kortlandt W, Vriens MR, Borel Rinkes IHM. The additional value of intraoperative parathyroid hormone assessment is marginal in patients with nonfamilial primary hyperparathyroidism: a prospective cohort study. *The American Journal of Surgery*. 2012;204(1):1-6.
52. Lubitz CCMDMPH, Chen HMD. Sestamibi-Negative Patients: To Operate or Image? *Annals of Surgical Oncology*. 2012;19(7):2086-7.
53. Gracie D, Hussain SSM. Use of minimally invasive parathyroidectomy techniques in sporadic primary hyperparathyroidism: systematic review. *The Journal of Laryngology and Otology*. 2012;126(3):221-7.
54. Nagar S, Reid D, Czako P, Long G, Shanley C. Outcomes analysis of intraoperative adjuncts during minimally invasive parathyroidectomy for primary hyperparathyroidism. *The American Journal of Surgery*. 2012;203(2):177-81.
55. Wong W, Foo FJ, Lau MI, Sarin A. Simplified minimally invasive parathyroidectomy: a series of 100 cases and review of the literature. *Annals of the Royal College of Surgeons of England*. 2011;93(4):290-3.
56. Quillo ARMD, Bumpous JMMD, Goldstein REMD, Fleming MMD, Flynn MBMD. Minimally Invasive Parathyroid Surgery, The Norman 20% Rule: Is it Valid? *The American Surgeon*. 2011;77(4):484-7.
57. Mazeh H, Chen H. Intraoperative adjuncts for parathyroid surgery. *Expert Review of Endocrinology & Metabolism*. 2011;6(2):245-53.
58. Neychev VK, Kouniavsky G, Shiue Z, Udall DN, Somervell H, Umbricht CB, et al. Chasing "Shadows": Discovering the Subtleties of Sestamibi Scans to Facilitate Minimally Invasive Parathyroidectomy. *World Journal of Surgery*. 2011;35(1):140-6.
59. McLean T, Delbridge L. Comparison of Consumer Information on the Internet to the Current Evidence Base for Minimally Invasive Parathyroidectomy. *World Journal of Surgery*. 2010;34(6):1304-11.
60. Ikeda Y, Takayama J, Takami H. Minimally invasive radioguided parathyroidectomy for hyperparathyroidism. *Annals of Nuclear Medicine*. 2010;24(4):233-40.
61. Morris LFMD, Zanocco KMD, Ituarte PH, G P, Ro K, Duh Q-yMDF, et al. The Value of Intraoperative Parathyroid Hormone Monitoring in Localized Primary Hyperparathyroidism: A Cost Analysis. *Annals of Surgical Oncology*. 2010;17(3):679-85.
62. Lombardi CP, Raffaelli M, Traini E, De Crea C, Corsello SM, Bellantone R. Video-Assisted Minimally Invasive Parathyroidectomy: Benefits and Long-Term Results. *World Journal of Surgery*. 2009;33(11):2266-81.
63. Barczynski M. Minimally invasive parathyroidectomy without intraoperative parathyroid hormone monitoring: When and why? *Journal of Postgraduate Medicine*. 2009;55(4):239-40.
64. Fraker DL, Harsono H, Lewis R. Minimally Invasive Parathyroidectomy: Benefits and Requirements of Localization, Diagnosis, and Intraoperative PTH Monitoring. Long-Term Results. *World Journal of Surgery*. 2009;33(11):2256-65.
65. Hacıyanlı M, Genc H, Damburaci N, Oruk G, Tutuncuoglu P, Erdogan N. Minimally invasive focused parathyroidectomy without using intraoperative parathyroid hormone monitoring or gamma probe. *Journal of Postgraduate Medicine*. 2009;55(4):242-6.
66. Harari A, Allendorf J, Shifrin A, DiGorgi M, Inabnet WB. Negative preoperative localization leads to greater resource use in the era of minimally invasive parathyroidectomy. *The American Journal of Surgery*. 2009;197(6):769-73.
67. Carling T, Udelsman R. Focused Approach to Parathyroidectomy. *World Journal of Surgery*. 2008;32(7):1512-7.
68. Soon PS, Delbridge LW, Sywak MS, Barraclough BM, Edhouse P, Sidhu SB. Surgeon Performed Ultrasound Facilitates Minimally Invasive Parathyroidectomy by the Focused Lateral Mini-incision Approach. *World Journal of Surgery*. 2008;32(5):766-71.
69. Miccoli P, Berti P, Materazzi G, Ambrosini CE, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTH) during endoscopic parathyroidectomy: A prospective randomized trial. *Surgical Endoscopy*. 2008;22(2):398-400.
70. Terris DJ, Stack BC, Jr., Gourin CG. Contemporary parathyroidectomy: exploiting technology. *American Journal of Otolaryngology*. 2007;28(6):408-14.
71. Beyer TD, Solorzano CC, Starr F, Nilubol N, Prinz RA. Parathyroidectomy outcomes according to operative approach. *The American Journal of Surgery*. 2007;193(3):368.
72. Battistella E, Pomba L, Toniato R, Burei M, Gregianin M, Fernando SW, et al. Evolution of the Diagnosis and Treatment of Primary Hyperparathyroidism. *Journal of Clinical Medicine*. 2023;12(5):2057.
73. Goldfarb M, Singer FR. Recent advances in the understanding and management of primary hyperparathyroidism. *F1000Research*. 2020;9.
74. Iwata AJ, Wertz AS, Alluri S, Singer MC. A faster parathyroidectomy: Techniques to shorten non-surgical operating room time. *American Journal of Otolaryngology*. 2019;40(6).
75. Li J, Vasilyeva E, Hiebert J, Britton H, Walker B, Wiseman SM. Limited clinical utility of intraoperative frozen section during parathyroidectomy for treatment of primary hyperparathyroidism. *The American Journal of Surgery*. 2019;217(5):893-8.
76. Mathew J, Arjunan R, Althaf S, Halkud R. Primary Hyperparathyroidism: Is Image Localization Alone Sufficient to Ensure Long-Term Cure in Unifocal Disease? *Cureus*. 2022;14(11).
77. Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *American Journal of Otolaryngology*. 2021;42(1).
78. Pickering JM, Giles WH. Improving Intraoperative Parathyroid Hormone Lab Efficiency. *The American Surgeon*. 2022;88(5):915-21.
79. Pradhan R, Gupta S, Agarwal A. Focused parathyroidectomy using accurate preoperative imaging and intraoperative PTH: Tertiary care experience. *Indian Journal of Endocrinology and Metabolism*. 2019;23(3):347-52.
80. Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally Invasive Parathyroidectomy without Intraoperative PTH Performed after Positive Ultrasonography as the only Diagnostic Method in Patients with Primary Hyperparathyroidism. *World Journal of Surgery*. 2019;43(6):1525-31.
81. Süleyman Özkan A, Adıyaman SC, Cevlik Ali Durubey Ç, Guray Durak Merih Güray D, Secil MS, Sevinc Ali İbrahim S. Intra-operative parathyroid hormone evaluation is superior to frozen section analysis in parathyroid surgery. *American Journal of Otolaryngology*. 2021;42(3).
82. Aygün N, Uludağ M. Surgical Treatment of Primary Hyperparathyroidism: Which Therapy to Whom? *Sisli Etfal Hastan Tip Bul.* 2019;53(3):201-14.
83. Ishii H, Mihai R, Watkinson JC, Kim DS. Systematic review of cure and recurrence rates following minimally invasive parathyroidectomy. *BJS Open*. 2018;2(6):364-70.
84. Sen S, Cherian A, Ramakant P, Reka K, Paul M, Abraham D. Focused parathyroidectomy under local anesthesia – A feasibility study. *Indian Journal of Endocrinology and Metabolism*. 2019;23(1):67-71.
85. Walsh NJ, Sullivan BT, Duke WS, Terris DJ. Routine bilateral neck exploration and four-gland dissection remains unnecessary in modern parathyroid surgery. *Laryngoscope Investigative Otolaryngology*. 2019;4(1):188-92.
86. Bombil I, Louw L, Mitchell C, Mahlobo F, Ramazani AM, Nthatheni RM. Sonar guided focused parathyroidectomy under cervical block. *South African Journal of Surgery*. 2018;56(2):30-3.
87. Parnell KE, Oltmann SC. The surgical management of primary hyperparathyroidism: an updated review. *International Journal of Endocrine Oncology*. 2018;5(1).
88. Yadav SK, Mishra SK, Mishra A, Mayilvagnan S, Chand G, Agarwal G, et al. Changing Profile of Primary Hyperparathyroidism Over Two and Half Decades: A Study in Tertiary Referral Center of North India. *World Journal of Surgery*. 2018;42(9):2732-7.
89. Al-Thani H, El-Matbouly M, Al-Sulaiti M, Asim M, Majzoub A, Tabea A, et al. Management and outcomes of hyperparathyroidism: a case series from a single institution over two decades. *Therapeutics and Clinical Risk Management*. 2018;14:1337-45.
90. Calò PG, Medas F, Loi G, Pisano G, Sorrenti S, Erdas E, et al. Parathyroidectomy for primary hyperparathyroidism in the elderly: experience of a single endocrine surgery center. *Aging Clinical and Experimental Research*, suppl 1. 2017;29:15-21.
91. Jinih MMBB, O'Connell EMBB, O'Leary DPMBBMP, Liew AMBBMMP, Redmond HPMFF. Focused Versus Bilateral Parathyroid Exploration for Primary Hyperparathyroidism: A Systematic Review and Meta-analysis. *Annals of Surgical Oncology*. 2017;24(7):1924-34.
92. Ozderya AMD, Temizkan SMD, Cetin K, Ozgur SMD, Gul AEMD, Aydin KMD. THE RESULTS OF PARATHYROID HORMONE ASSAY IN PARATHYROID ASPIRATES IN PRE-OPERATIVE LOCALIZATION OF PARATHYROID ADENOMAS FOR FOCUSED PARATHYROIDECTOMY IN PATIENTS WITH NEGATIVE OR SUSPICIOUS TECHNETIUM-99M-SESTAMIBI SCANS. *Endocrine Practice*. 2017;23(9):1101-6.
93. Polistena AMDD, Lucchini RMD, Monacelli MMD, Triola RMD, Avenia SMD, Barillaro IMD, et al. Current Indications for Surgical Treatment of Primary Hyperparathyroidism in the Elderly. *The American Surgeon*. 2017;83(3):296-302.
94. Chow TL, Choi CY, Lam SH. Focused parathyroidectomy without intra-operative parathyroid hormone monitoring for primary hyperparathyroidism: results in a low-volume hospital. *The Journal of Laryngology and Otology*. 2015;129(8):788-94.
95. Habib Z, Kabaker A, Camacho P. Sporadic primary hyperparathyroidism: medical and surgical updates. *Expert Review of Endocrinology & Metabolism*. 2014;9(1):31-44.
96. Kiernan CM, Schlegel C, Isom C, Kavalukas S, Peters MF, Solorzano CC. Ambulatory bilateral neck exploration for primary hyperparathyroidism: is it safe? *The American Journal of Surgery*. 2016;212(4):722-7.
97. Mownah OA, Pafitanis G, Drake WM, Crinnion JN. Contemporary surgical treatment of primary hyperparathyroidism without intraoperative parathyroid hormone measurement. *Annals of the Royal College of Surgeons of England*. 2015;97(8):603-7.

98. Nair C, Babu M, Jacob P, Menon R, Mathew J. Is intraoperative parathyroid hormone monitoring necessary in symptomatic primary hyperparathyroidism with concordant imaging? *Indian Journal of Endocrinology and Metabolism*. 2016;20(4).
99. Sharata AMD, Kelly TLBSC, Rozenfeld YMPH, Hammill CWMD, SchumanMd E, Carlisle JRMD, et al. Management of Primary Hyperparathyroidism: Can We Do Better? *The American Surgeon*. 2017;83(1):64-70.
- 100 Singh Ospina NM, Rodriguez-gutierrez R, Maraka S, Espinosa De Ycaza AE, Jasim S, Castaneda-guarderas A, et al. Outcomes of Parathyroidectomy in Patients with Primary Hyperparathyroidism: A Systematic Review and Meta-analysis. *World Journal of Surgery*. 2016;40(10):2359-77.
- 101 Sun Moon K, Shu AD, Long J, Montez-Rath ME, Leonard MB, Norton JA, et al. Declining Rates of Inpatient Parathyroidectomy for Primary Hyperparathyroidism in the US. *PLoS One*. 2016;11(8).
- 102 Usulukaya O, Gumus M, Tasdemir B, Goya C, Kilinc F, Oguz A, et al. Improvement of minimally invasive parathyroidectomy outcomes by real time ultrasonography performed by a surgeon and radiologist team. *Medical Ultrasonography*. 2015;17(3):315-21.
- 103 Calò PG, Pisano G, Loi G, Medas F, Barca L, Atzeni M, et al. Intraoperative parathyroid hormone assay during focused parathyroidectomy: the importance of 20 minutes measurement. *BMC Surgery*. 2013;13:36.
- 104 Gupta AMD, Unawane AMD, Subhas GMD, Herschman BRMD, Silapaswan SMD, Kolachalam RMD, et al. Parathyroidectomies Using Intraoperative Parathormone Monitoring: When Should We Stop Measuring Intraoperative Parathormone Levels? *The American Surgeon*. 2012;78(8):844-50.
- 105 Karahan Ö, Okus A, Sevinç B, Eryilmaz M, Ay S, Çaycı M, et al. Minimally invasive parathyroidectomy under local anesthesia. *Journal of Postgraduate Medicine*. 2013;59(1):21-4.
- 106 Lubitz CCMDMPH, Stephen AEMD, Hodin RAMD, Pandharipande PMDMPH. Preoperative Localization Strategies for Primary Hyperparathyroidism: An Economic Analysis. *Annals of Surgical Oncology*. 2012;19(13):4202-9.
- 107 Mohebaty A, Shaha AR. Imaging techniques in parathyroid surgery for primary hyperparathyroidism. *American Journal of Otolaryngology*. 2012;33(4):457-68.
- 108 Opoku-Boateng AMD, Bolton JSMD, Corsetti RMD, Brown REM, Oxner CMD, Fuhrman GMM. Use of a Sestamibi-only Approach to Routine Minimally Invasive Parathyroidectomy. *The American Surgeon*. 2013;79(8):797-801.
- 109 Paruk IM, Esterhuizen TM, Maharaj S, Pirie FJ, Motala AA. Characteristics, management and outcome of primary hyperparathyroidism in South Africa: a single-centre experience. *Postgraduate Medical Journal*. 2013;89(1057):626.
- 110 Tee MCMMDMPH, Chan SKM, Nguyen VMD, Strugnell SSM, Yang JB, Jones SMD, et al. Incremental value and clinical impact of neck sonography for primary hyperparathyroidism: a risk-adjusted analysis. *Canadian Journal of Surgery*. 2013;56(5):325-31.
- 111 Twigt BA, van Dalen T, Tollwegt AM, Kortlandt W, Vriens MR, Borel Rinkes IHM. The additional value of intraoperative parathyroid hormone assessment is marginal in patients with nonfamilial primary hyperparathyroidism: a prospective cohort study. *The American Journal of Surgery*. 2012;204(1):1-6.
- 112 Zia SMD, Sippel RSMD, Chen HMD. Sestamibi Imaging for Primary Hyperparathyroidism: The Impact of Surgeon Interpretation and Radiologist Volume. *Annals of Surgical Oncology*. 2012;19(12):3827-31.
- 113 Gracie D, Hussain SSM. Use of minimally invasive parathyroidectomy techniques in sporadic primary hyperparathyroidism: systematic review. *The Journal of Laryngology and Otology*. 2012;126(3):221-7.
- 114 Ikeda Y, Takayama J, Takami H. Minimally invasive radioguided parathyroidectomy for hyperparathyroidism. *Annals of Nuclear Medicine*. 2010;24(4):233-40.
- 115 Lubitz CCMDMPH, Chen HMD. Sestamibi-Negative Patients: To Operate or Image? *Annals of Surgical Oncology*. 2012;19(7):2086-7.
- 116 Mazeh H, Chen H. Intraoperative adjuncts for parathyroid surgery. *Expert Review of Endocrinology & Metabolism*. 2011;6(2):245-53.
- 117 McLean T, Delbridge L. Comparison of Consumer Information on the Internet to the Current Evidence Base for Minimally Invasive Parathyroidectomy. *World Journal of Surgery*. 2010;34(6):1304-11.
- 118 Nagar S, Reid D, Czako P, Long G, Shanley C. Outcomes analysis of intraoperative adjuncts during minimally invasive parathyroidectomy for primary hyperparathyroidism. *The American Journal of Surgery*. 2012;203(2):177-81.
- 119 Neychev VK, Kouniavsky G, Shiue Z, Udall DN, Somervell H, Umbricht CB, et al. Chasing "Shadows": Discovering the Subtleties of Sestamibi Scans to Facilitate Minimally Invasive Parathyroidectomy. *World Journal of Surgery*. 2011;35(1):140-6.
- 120 Quillo ARMD, Bumpous JMMD, Goldstein REMD, Fleming MME, Flynn MBMD. Minimally Invasive Parathyroid Surgery, The Norman 20% Rule: Is it Valid? *The American Surgeon*. 2011;77(4):484-7.
- 121 Wong W, Foo FJ, Lau MI, Sarin A. Simplified minimally invasive parathyroidectomy: a series of 100 cases and review of the literature. *Annals of the Royal College of Surgeons of England*. 2011;93(4):290-3.
- 122 Barczynski M. Minimally invasive parathyroidectomy without intraoperative parathyroid hormone monitoring: When and why? *Journal of Postgraduate Medicine*. 2009;55(4):239-40.
- 123 Beyer TD, Solorzano CC, Starr F, Nilubol N, Prinz RA. Parathyroidectomy outcomes according to operative approach. *The American Journal of Surgery*. 2007;193(3):368.
- 124 Carling T, Udelsman R. Focused Approach to Parathyroidectomy. *World Journal of Surgery*. 2008;32(7):1512-7.
- 125 Fraker DL, Harsono H, Lewis R. Minimally Invasive Parathyroidectomy: Benefits and Requirements of Localization, Diagnosis, and Intraoperative PTH Monitoring. Long-Term Results. *World Journal of Surgery*. 2009;33(11):2256-65.
- 126 Hacıyanlı M, Genc H, Damburacı N, Oruk G, Tutuncuoğlu P, Erdoğan N. Minimally invasive focused parathyroidectomy without using intraoperative parathyroid hormone monitoring or gamma probe. *Journal of Postgraduate Medicine*. 2009;55(4):242-6.
- 127 Harari A, Allendorf J, Shifrin A, DiGorgi M, Inabnet WB. Negative preoperative localization leads to greater resource use in the era of minimally invasive parathyroidectomy. *The American Journal of Surgery*. 2009;197(6):769-73.
- 128 Lombardi CP, Raffaelli M, Traini E, De Crea C, Corsello SM, Bellantone R. Video-Assisted Minimally Invasive Parathyroidectomy: Benefits and Long-Term Results. *World Journal of Surgery*. 2009;33(11):2266-81.
- 129 Miccoli P, Berti P, Materazzi G, Ambrosini CE, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTHa) during endoscopic parathyroidectomy: A prospective randomized trial. *Surgical Endoscopy*. 2008;22(2):398-400.
- 130 Morris LFMD, Zanocco KMD, Ituarte PH, G P, Ro K, Duh Q-yMDF, et al. The Value of Intraoperative Parathyroid Hormone Monitoring in Localized Primary Hyperparathyroidism: A Cost Analysis. *Annals of Surgical Oncology*. 2010;17(3):679-85.
- 131 Soon PS, Delbridge LW, Sywak MS, Barraclough BM, Edhouse P, Sidhu SB. Surgeon Performed Ultrasound Facilitates Minimally Invasive Parathyroidectomy by the Focused Lateral Mini-incision Approach. *World Journal of Surgery*. 2008;32(5):766-71.
- 132 Terris DJ, Stack BC, Jr., Gourin CG. Contemporary parathyroidectomy: exploiting technology. *American Journal of Otolaryngology*. 2007;28(6):408-14.

#### WorldCat

1. Kunstman JW, Kirsch JD, Mahajan A, Udelsman R. Parathyroid Localization and Implications for Clinical Management. *The Journal of Clinical Endocrinology & Metabolism*. 2013;98(3):902-12.
2. Carling T, Udelsman R. Minimally Invasive Parathyroidectomy. *Surgery of the Thyroid and Parathyroid Glands*: Springer Berlin Heidelberg; 2012. p. 475-85.
3. Udelsman R, Pasieka JL, Sturgeon C, Young JEM, Clark OH. Surgery for asymptomatic primary hyperparathyroidism: proceedings of the third international workshop. *The Journal of clinical endocrinology and metabolism*. 2009;94(2):366-72.
4. Noureldine SI, Gooi Z, Zufano RP. Minimally invasive parathyroid surgery. *Gland surgery*. 2015;4(5):410-9.
5. Update on Preoperative Parathyroid Localization in Primary Hyperparathyroidism. *Endocrinology and Metabolism* (구 대한내분비학회지) [Internet]. 2022; 37(5):[744-55 pp.].
6. Stålberg P, Delbridge L, van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy : current concepts. Uppsala universitet, Institutionen för kirurgiska vetenskaper; 2007.
7. Carling TMDPD, Udelsman RMDMBA. Minimally Invasive Parathyroidectomy. *Surgery of the Thyroid and Parathyroid Glands*: Berlin, Heidelberg : Springer Berlin Heidelberg : Springer; 2012. p. 475-85.
8. Nowlin WFMDFASCRS. Parathyroidectomy. *Advanced Surgical Techniques for Rural Surgeons*: New York, NY : Springer New York : Springer; 2015. p. 147-55.
9. Aarum S, Nordenström J, Reinhér E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for Primary Hyperparathyroidism: The New versus the Old Order: A Randomised Controlled Trial of Preoperative Localisation. *Scandinavian Journal of Surgery*. 2007;96(1):26-30.
10. Stalberg P, Delbridge L, van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy--current concepts. The surgeon : journal of the Royal Colleges of Surgeons of Edinburgh and Ireland. 2007;5(5):301-8.
11. Aarum S, Nordenström J, Reinhér E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for primary hyperparathyroidism: the new versus the old order. A randomised controlled trial of preoperative localisation. *Scandinavian journal of surgery : SJS : official organ for the Finnish Surgical Society and the Scandinavian Surgical Society*. 2007;96(1):26-30.
12. Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *European journal of endocrinology*. 2016;174(1):1-8.
13. Lee L, Steward DL. Techniques for parathyroid localization with ultrasound. *Otolaryngologic clinics of North America*. 2010;43(6):1229-39.
14. Kunstman JW, Kirsch JD, Mahajan A, Udelsman R. Clinical review: Parathyroid localization and implications for clinical management. *The Journal of clinical endocrinology and metabolism*. 2013;98(3):902-12.
15. Arora S, Balash PR, Yoo J, Smith GS, Prinz RA. Benefits of surgeon-performed ultrasound for primary hyperparathyroidism. *Langenbeck's archives of surgery*. 2009;394(5):861-7.
16. Park H-S, Hong N, Jeong JJ, Yun M, Rhee Y. Update on Preoperative Parathyroid Localization in Primary Hyperparathyroidism. *Endocrinology and metabolism* (Seoul, Korea). 2022;37(5):744-55.
17. Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: Feasibility and outcome. *Journal of visceral surgery*. 2018;155(4):253-8.

18. Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *American journal of otolaryngology*. 2021;42(1):102819.
19. Harari A, Allendorf J, Shifrin A, DiGorgi M, Inabnet WBDog, Endocrine Surgery DoSFWACoP, et al. Negative preoperative localization leads to greater resource use in the era of minimally invasive parathyroidectomy. *The American Journal of Surgery*. 2009;197(6):769-73.
20. Fang W-L, Tseng L-M, Chen J-Y, Chiou S-Y, Chou Y-H, Wu C-W, et al. The management of high-risk patients with primary hyperparathyroidism - minimally invasive parathyroidectomy vs. medical treatment. *Clinical endocrinology*. 2008;68(4):520-8.
21. Udelsman R, Lin Z, Donovan P. The superiority of minimally invasive parathyroidectomy based on 1650 consecutive patients with primary hyperparathyroidism. *Annals of surgery*. 2011;253(3):585-91.
22. Kunstman JW, Udelsman R. Superiority of minimally invasive parathyroidectomy. *Advances in surgery*. 2012;46:171-89.
23. Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *The Surgical clinics of North America*. 2014;94(3):607-23.
24. Slepavicius A, Beisa V, Janusonis V, Strupas K. Focused versus conventional parathyroidectomy for primary hyperparathyroidism: a prospective, randomized, blinded trial. *Langenbeck's archives of surgery*. 2008;393(5):659-66.
25. Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally Invasive Parathyroidectomy without Intraoperative PTH Performed after Positive Ultrasonography as the only Diagnostic Method in Patients with Primary Hyperparathyroidism. *World journal of surgery*. 2019;43(6):1525-31.
26. 43rd SEMDSA and 13th NOFSA Congresses and 11th DESSA Workshop : 12 - 15 April 2008, Cape Town : abstracts. *Journal of Endocrinology, Metabolism and Diabetes in South Africa*. 2008;13(1):20-44.
27. Anderson RL, Anderson MA. Rural General Surgery: A Review of the Current Situation and Realities from a Rural Community Practice in Central Nebraska. *Online Journal of Rural Research & Policy [Internet]*. 2012; 7(2).
28. Afzal A, Gauhar TM, Butt WT, Khawaja AA, Azim KM. Management of hyperparathyroidism: a five year surgical experience. *JPMA The Journal of the Pakistan Medical Association*. 2011;61(12):1194-8.
29. Usta A, Alhan E, Cinel A, Türkyılmaz S, Erem C. A 20-year study on 190 patients with primary hyperparathyroidism in a developing country: Turkey experience. *International surgery*. 2015;100(4):648-55.
30. Siddiqui MI, Pasha HA, Asad R, Talati JJ. Changing paradigms in the surgical management of hyperparathyroidism at a tertiary care hospital in a developing country. *JPMA The Journal of the Pakistan Medical Association*. 2019;69(9):1360-4.
31. Panieri E. What's new in endocrine surgery? : main article. *CME : Your SA Journal of CPD*. 2007;25(7):312-3.
32. Mitchell C, Madima NR, Louw L, Muganza RA, Mahlobo F, Bombil I. Sonar guided focused parathyroidectomy under cervical block. *South African Journal of Surgery*. 2018;56(2):30-3.
33. Calò PG, Medas F, Loi G, Erdas E, Pisano G, Nicolosi A. Feasibility of unilateral parathyroidectomy in patients with primary hyperparathyroidism and negative or discordant localization studies. *Updates in surgery*. 2016;68(2):155-61.
34. Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: Feasibility and outcome. *Journal of visceral surgery*. 2018;155(4):253-8.
35. Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia – A Feasibility Study. *Indian Journal of Endocrinology and Metabolism*. 2019;23(1):67-71.
36. Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia – A Feasibility Study. *Indian journal of endocrinology and metabolism*. 2019;23(1):67-71.
37. El-Hady HA, Radwan HS. Focused parathyroidectomy for single parathyroid adenoma: a clinical account of 20 patients. *Electronic physician*. 2018;10(6):6974-80.
38. Kandil E, Malazai AJ, Alrasheedi S, Tufano RP. Minimally invasive/focused parathyroidectomy in patients with negative sestamibi scan results. *Archives of otolaryngology--head & neck surgery*. 2012;138(3):223-5.
39. Bombil I, Louw L, Mitchell C, Mahlobo F, Muganza RA, Madima NR. Sonar guided focused parathyroidectomy under cervical block. *South African Journal of Surgery*. 2018;56(2):30-3.

#### Google Scholar

1. Noltes ME, Cottrell J, Madani A, Rotstein L, Gomez-Hernandez K, Devon K, et al. Quality indicators for the diagnosis and management of primary hyperparathyroidism. *JAMA Otolaryngology–Head & Neck Surgery*. 2022;148(3):209-19.
2. Yadav SK, Mishra SK, Mishra A, Mayilvagnan S, Chand G, Agarwal G, et al. Surgical management of primary hyperparathyroidism in the era of focused parathyroidectomy: A study in tertiary referral centre of North India. *Indian journal of endocrinology and metabolism*. 2019;23(4):468.
3. Aarum S, Nordenström J, Reihner E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for primary hyperparathyroidism: the new versus the old order: a randomised controlled trial of preoperative localisation. *Scandinavian journal of surgery*. 2007;96(1):26-30.
4. Adler Y, Tzelnick S, Shopen Y, Reifen E, Bachar G, Shpitzer T, et al. Retrospective analysis of the role of intra-operative parathyroid hormone monitoring during parathyroidectomy for primary hyperparathyroidism: a single center experience over 2 decades. *European Archives of Oto-Rhino-Laryngology*. 2022;279(11):5375-80.
5. Badii B, Staderini F, Foppa C, Tofani L, Skalamera I, Fiorenza G, et al. Cost–benefit analysis of the intraoperative parathyroid hormone assay in primary hyperparathyroidism. *Head & Neck*. 2017;39(2):241-6.
6. Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: feasibility and outcome. *Journal of Visceral Surgery*. 2018;155(4):253-8.
7. Bhutiani N, Graffree B, Martin RC, Quillo AR. Factors influencing time to decrease in intraoperative parathyroid hormone (ioPTH) levels in patients undergoing focused parathyroidectomy (FP). *The American Surgeon*. 2018;84(6):772-5.
8. Cheung K, Wang TS, Farrokhyar F, Roman SA, Sosa JA. A meta-analysis of preoperative localization techniques for patients with primary hyperparathyroidism. *Annals of surgical oncology*. 2012;19:577-83.
9. Di Marco A, Mechera R, Glover A, Papachristos A, Clifton-Bligh R, Delbridge L, et al. Focused parathyroidectomy without intraoperative parathyroid hormone measurement in primary hyperparathyroidism: Still a valid approach? *Surgery*. 2021;170(5):1383-8.
10. Di Stasio E, Carrozza C, Lombardi CP, Raffaelli M, Traini E, Bellantone R, et al. Parathyroidectomy monitored by intra-operative PTH: the relevance of the 20 min end-point. *Clinical biochemistry*. 2007;40(9-10):595-603.
11. Donatini G, Marcinjak C, Lenne X, Clément G, Brundet A, Sebag F, et al. Risk factors of redo surgery after unilateral focused parathyroidectomy: conclusions from a comprehensive nationwide database of 13,247 interventions over 6 years. *Annals of Surgery*. 2020;272(5):801-6.
12. Dowthwaite SA, Young JE, Pasternak JD, Yoo J. Surgical management of primary hyperparathyroidism. *Journal of Clinical Densitometry*. 2013;16(1):48-53.
13. Duke WS, Terris DJ. Minimally Invasive Parathyroidectomy: North American Modifications. *Parathyroid Surgery: Fundamental and Advanced Concepts*. 2013:119.
14. Elaraj DM, Sippel RS, Lindsay S, Sansano I, Duh Q-Y, Clark OH, et al. Are additional localization studies and referral indicated for patients with primary hyperparathyroidism who have negative sestamibi scan results? *Archives of Surgery*. 2010;145(6):578-81.
15. Erinjeri NJ, Udelsman R. Volume–outcome relationship in parathyroid surgery. *Best Practice & Research Clinical Endocrinology & Metabolism*. 2019;33(5):101287.
16. Greenleaf EK, Saunders BD, Schaefer EW, Hollenbeak CS. Health Services and Health Care Economics Related to Hyperparathyroidism and Parathyroid Surgery. *Medical and Surgical Treatment of Parathyroid Diseases: An Evidence-Based Approach*. 2017:489-500.
17. Guerin C, Lowery A, Gabriel S, Castinetti F, Philippon M, Vaillant-Lombard J, et al. Preoperative imaging for focused parathyroidectomy: making a good strategy even better. *European Journal of Endocrinology*. 2015;172(5):519-26.
18. Hanba C, Bobian M, Svider PF, Sheyn A, Siegel B, Lin H-S, et al. Perioperative considerations and complications in pediatric parathyroidectomy. *International Journal of Pediatric Otorhinolaryngology*. 2016;91:94-9.
19. Herb J, Staley BS, Roberson M, Strassle PD, Kim LT. Use and disparities in parathyroidectomy for symptomatic primary hyperparathyroidism in the Medicare population. *Surgery*. 2021;170(5):1376-82.
20. Iacobone M, Scerrino G, Palazzo FF. Parathyroid surgery: an evidence-based volume—outcomes analysis: European Society of Endocrine Surgeons (ESES) positional statement. *Langenbeck's archives of surgery*. 2019;404:919-27.
21. Iwata AJ, Wertz AS, Alluri S, Singer MC. A faster parathyroidectomy: Techniques to shorten non-surgical operating room time. *American Journal of Otolaryngology*. 2019;40(6):102292.
22. Kandil E, Malazai AJ, Alrasheedi S, Tufano RP. Minimally invasive/focused parathyroidectomy in patients with negative sestamibi scan results. *Archives of Otolaryngology–Head & Neck Surgery*. 2012;138(3):223-5.
23. Kiernan CM, Schlegel C, Isom C, Kavalukas S, Peters MF, Solórzano CC. Ambulatory bilateral neck exploration for primary hyperparathyroidism: is it safe? *The American Journal of Surgery*. 2016;212(4):722-7.
24. Kiernan CM, Wang T, Perrier ND, Grubbs EG, Solórzano CC. Bilateral neck exploration for sporadic primary hyperparathyroidism: use patterns in 5,597 patients undergoing parathyroidectomy in the collaborative endocrine surgery quality improvement program. *Journal of the American College of Surgeons*. 2019;228(4):652-9.
25. Kunstman JW, Kirsch JD, Mahajan A, Udelsman R. Parathyroid localization and implications for clinical management. *The Journal of Clinical Endocrinology & Metabolism*. 2013;98(3):902-12.
26. Kunstman JW, Udelsman R. Superiority of minimally invasive parathyroidectomy. *Advances in surgery*. 2012;46(1):171-89.
27. Kuzminski SJ, Sosa JA, Hoang JK. Update in parathyroid imaging. *Magnetic Resonance Imaging Clinics*. 2018;26(1):151-66.
28. Laird AM, Libutti SK. Minimally invasive parathyroidectomy versus bilateral neck exploration for primary hyperparathyroidism. *Surgical Oncology Clinics*. 2016;25(1):103-18.

29. Langusch CC, Norlen O, Titmuss A, Donoghue K, Holland AJ, Shun A, et al. Focused image-guided parathyroidectomy in the current management of primary hyperparathyroidism. *Archives of Disease in Childhood*. 2015;100(10):924-7.
30. Lee J, Kim S-j, Yu HW, Chai YJ, Choi JY, Lee KE. Changing Trends in Preoperative Localization and Surgical Techniques for the Treatment of Primary Hyperparathyroidism in a Single Tertiary Center. *Journal of Endocrine Surgery*. 2019;19(4):126-35.
31. Lew JI, Rivera M, Irvin GL, Solorzano CC. Operative failure in the era of focused parathyroidectomy: a contemporary series of 845 patients. *Archives of Surgery*. 2010;145(7):628-33.
32. Liddy S, Worsley D, Torreggiani W, Feeney J. Preoperative imaging in primary hyperparathyroidism: literature review and recommendations. *Canadian Association of Radiologists Journal*. 2017;68(1):47-55.
33. Lubitz CC, Duh Q-Y. Guide to Preoperative Parathyroid Localization Testing. *Surgery of the thyroid and parathyroid glands*: Elsevier; 2021. p. 494-501. e3.
34. Medas F, Cappellacci F, Canu GL, Noordzij JP, Erdas E, Calo PG. The role of Rapid Intraoperative Parathyroid Hormone (ioPTH) assay in determining outcome of parathyroidectomy in primary hyperparathyroidism: A systematic review and meta-analysis. *International Journal of Surgery*. 2021;92:106042.
35. Mihai R, Simon D, Hellman P. Imaging for primary hyperparathyroidism—an evidence-based analysis. *Langenbeck's archives of surgery*. 2009;394:765-84.
36. Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *European Journal of Endocrinology*. 2016;174(1):D1-D8.
37. Nagar S, Reid D, Czako P, Long G, Shanley C. Outcomes analysis of intraoperative adjuncts during minimally invasive parathyroidectomy for primary hyperparathyroidism. *The American journal of surgery*. 2012;203(2):177-81.
38. Parikh AM, Grogan RH, Moron FE. Localization of parathyroid disease in reoperative patients with primary hyperparathyroidism. *International Journal of Endocrinology*. 2020;2020.
39. Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *American Journal of Otolaryngology*. 2021;42(1):102819.
40. Quinn AJ, Ryan ÉJ, Garry S, James DL, Boland MR, Young O, et al. Use of intraoperative parathyroid hormone in minimally invasive parathyroidectomy for primary hyperparathyroidism: a systematic review and meta-analysis. *JAMA Otolaryngology—Head & Neck Surgery*. 2021;147(2):135-43.
41. Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally invasive parathyroidectomy without intraoperative PTH performed after positive ultrasonography as the only diagnostic method in patients with primary hyperparathyroidism. *World Journal of Surgery*. 2019;43:1525-31.
42. Shah-Becker S, Goldenberg D. Minimally invasive parathyroidectomy. *Operative Techniques in Otolaryngology-Head and Neck Surgery*. 2016;27(3):152-6.
43. Sharata A, Kelly TL, Rozenfeld Y, Hammill CW, Schuman E, Carlisle JR, et al. Management of primary hyperparathyroidism: can we do better? *The American Surgeon*. 2017;83(1):64-70.
44. Shifrin A. Advances in the diagnosis and surgical management of primary hyperparathyroidism. *Advances in treatment and management in surgical endocrinology*. 2020:71-83.
45. Shindo ML, Rosenthal JM. Minimal access parathyroidectomy using the focused lateral approach: technique, indication, and results. *Archives of Otolaryngology—Head & Neck Surgery*. 2007;133(12):1227-34.
46. Solorzano CC, Carneiro-Pla D. Minimizing cost and maximizing success in the preoperative localization strategy for primary hyperparathyroidism. *Surgical Clinics*. 2014;94(3):587-605.
47. Soon PS, Delbridge LW, Sywak MS, Barraclough BM, Edhouse P, Sidhu SB. Surgeon performed ultrasound facilitates minimally invasive parathyroidectomy by the focused lateral mini-incision approach. *World journal of surgery*. 2008;32:766-71.
48. Sosa JA, Tuggle CT, Wang TS, Thomas DC, Boudourakis L, Rivkees S, et al. Clinical and economic outcomes of thyroid and parathyroid surgery in children. *The Journal of Clinical Endocrinology & Metabolism*. 2008;93(8):3058-65.
49. Soylu L, Aydın OU, İlğan S, Özbaş S, Bilezikçi B, Gürsoy A, et al. Radioguided occult lesion localization for minimally-invasive parathyroidectomy without quick PTH monitoring and frozen section: impact of the learning curve. *Turkish Journal of Surgery*. 2020;36(3):297.
50. Udelsman R, Pasieka JL, Sturgeon C, Young J, Clark OH. Surgery for asymptomatic primary hyperparathyroidism: proceedings of the third international workshop. *The Journal of Clinical Endocrinology & Metabolism*. 2009;94(2):366-72.
51. UK NGC. Evidence review for surgical interventions. 2019.
52. UK NGC. Evidence review for surgical localisation. 2019.
53. Van Udelsman B, Udelsman R. Surgery in primary hyperparathyroidism: extensive personal experience. *Journal of Clinical Densitometry*. 2013;16(1):54-9.
54. Xie S, Kuriakose J, Beninato T, Carayannopoulos M, Trooskin SZ, Libutti SK, et al. Association of implementation of operating room-based parathyroid hormone testing with operative time and cost. *Journal of the American College of Surgeons*. 2022;235(6):906-12.
55. Yip L, Silverberg SJ, Fuleihan G, Rosen CJ. Preoperative localization for parathyroid surgery in patients with primary hyperparathyroidism. UpToDate Carty SE, Rosen CJ, Chen W (ed): UpToDate, Waltham, MA. 2020.
56. Zawawi F, Mlynarek AM, Cantor A, Varshney R, Black MJ, Hier MP, et al. Intraoperative parathyroid hormone level in parathyroidectomy: which patients benefit from it? *Journal of Otolaryngology-Head & Neck Surgery*. 2013;42:1-6.
57. Harari A, Allendorf J, Shifrin A, DiGorgi M, Inabnet WB. Negative preoperative localization leads to greater resource use in the era of minimally invasive parathyroidectomy. *The American journal of surgery*. 2009;197(6):769-73.
58. Abbott DE, Cantor SB, Grubbs EG, Santora R, Gomez HF, Evans DB, et al. Outcomes and economic analysis of routine preoperative 4-dimensional CT for surgical intervention in de novo primary hyperparathyroidism: does clinical benefit justify the cost? *Journal of the American College of Surgeons*. 2012;214(4):629-37.
59. Afzal A, Gauhar TM, Butt WT, Khawaja AA, Azim KM. Management of hyperparathyroidism: A live year surgical experience. *JPMA-Journal of the Pakistan Medical Association*. 2011;61(12):1194.
60. Agarwal S, Agarwal A, Chand G. The Necessity and Reliability of Intraoperative Parathyroid Hormone (PTH) Testing in Patients with Mild Hyperparathyroidism and PTH Levels in the Normal Range. *World journal of surgery*. 2012;36:483-.
61. Alikor CA, Engelbrecht G. The Usefulness of 99m technetium-sestamibi Parathyroid Scintigraphy in Preoperative Localization of Parathyroid Adenoma in Patients with Primary Hyperparathyroidism at an Academic Hospital in South Africa.
62. Baliski C, Nosyk B, Melck A, Bugis S, Rosenberg F, H. Anis A. The cost-effectiveness of three strategies for the surgical treatment of symptomatic primary hyperparathyroidism. *Annals of Surgical Oncology*. 2008;15:2653-60.
63. Bansal N, Mayilvaganan S. Primary hyperparathyroidism with negative imaging: a significant clinical problem. *Annals of Surgery*. 2017;265(4):e41.
64. Barczyński M. Introduction to focused issue on novel technologies in endocrine surgery. *Gland surgery*. 2020;9(Suppl 2):S65.
65. Bedi HK, Jedrejko N, Nguyen A, Aspinall SR, Wiseman SM. Thyroid and parathyroid surgeon case volume influences patient outcomes: a systematic review. *Surgical Oncology*. 2021;38:101550.
66. Bellantone R, Raffaelli M, De Crea C, Traini E, Lombardi CP. Minimally-invasive parathyroid surgery. *Acta Otorhinolaryngologica Italica*. 2011;31(4):207.
67. Bergenfelz AO, Jansson SK, Wallin GK, Mårtensson HG, Rasmussen L, Eriksson HL, et al. Impact of modern techniques on short-term outcome after surgery for primary hyperparathyroidism: a multicenter study comprising 2,708 patients. *Langenbeck's archives of surgery*. 2009;394:851-60.
68. Beyer TD, Solorzano CC, Starr F, Nilubol N, Prinz RA. Parathyroidectomy outcomes according to operative approach. *The American journal of surgery*. 2007;193(3):368-73.
69. Bhadada SK, Arya AK, Mukhopadhyay S, Khadgawat R, Sukumar S, Lodha S, et al. Primary hyperparathyroidism: insights from the Indian PHPT registry. *Journal of bone and mineral metabolism*. 2018;36:238-45.
70. Castaldi M, Roberts S, Latifi R. Prophylactic Parathyroidectomy. *Prophylactic Surgery*. 2021:51-60.
71. Chao TE, Sharma K, Mandigo M, Hagander L, Resch SC, Weiser TG, et al. Cost-effectiveness of surgery and its policy implications for global health: a systematic review and analysis. *The Lancet Global Health*. 2014;2(6):e334-e45.
72. Chow T-L. Intraoperative Parathyroid Hormone in Minimally Invasive Parathyroidectomy. *JAMA Otolaryngology—Head & Neck Surgery*. 2021;147(10):922-3.
73. Egan RJ, Scott-Coombes DM. The surgical management of sporadic primary hyperparathyroidism. *Best Practice & Research Clinical Endocrinology & Metabolism*. 2018;32(6):847-59.
74. Gaurav K, Mishra SK. Management of hyperparathyroidism—Influence of patient and disease factors. *The American Journal of Surgery*. 2018;216(1):176.
75. Gil-Cárdenas A, Gamino R, Reza A, Pantoja JP, Herrera MF. Is intraoperative parathyroid hormone assay mandatory for the success of targeted parathyroidectomy? *Journal of the American College of Surgeons*. 2007;204(2):286-90.
76. Goonewardena SA. Judicious use of frozen section in parathyroid surgery. *Sri Lanka Journal of Surgery*. 2019;37(2).
77. Hacıyanli M, Genc H, Damburaci N, Oruk G, Tutuncuoglu P, Erdogan N. Minimally invasive focused parathyroidectomy without using intraoperative parathyroid hormone monitoring or gamma probe. *Journal of Postgraduate Medicine*. 2009;55(4):242.
78. Hwang RS, Morris LF, Ro K, Park S, Ituarte PH, Hong JC, et al. A selective, Bayesian approach to intraoperative PTH monitoring. *Annals of surgery*. 2010;251(6):1122-6.
79. Jha CK, Bichoo RA, Yadav SK, Sonthineni C, Bothra S. Ambulatory bilateral neck exploration for primary hyperparathyroidism: Is it safe? *The American Journal of Surgery*. 2017;213(6):1191-2.
80. Judson BL, Shaha AR. Nuclear imaging and minimally invasive surgery in the management of hyperparathyroidism. *Journal of Nuclear Medicine*. 2008;49(11):1813-8.

81. KBhadada S, Singh P, Rao SD. Primary Hyperparathyroidism in the Indian Context. *Clinical Focus Series: Bone and Mineral Disorders*. 2019.
82. Khalid R, Hassaan M, Raza A. Clinical Presentation of Primary Hyperparathyroidism in Lahore, Pakistan—A Single Center Study. *Endocrinology and Disorders*. 2021;5(5):2640-1045.
83. Khan A, Hanley D, Rizzoli R, Bollerslev J, Young J, Rejnmark L, et al. Primary hyperparathyroidism: review and recommendations on evaluation, diagnosis, and management. A Canadian and international consensus. *Osteoporosis International*. 2017;28:1-19.
84. Khan ZF, Lew JI. Intraoperative parathyroid hormone monitoring in the surgical management of sporadic primary hyperparathyroidism. *Endocrinology and Metabolism*. 2019;34(4):327-39.
85. Kirdak T, Duh Q-Y, Kebebew E, Clark OH. Do patients undergoing parathyroidectomy for primary hyperparathyroidism in San Francisco, CA, and Bursa, Turkey, differ? *The American journal of surgery*. 2009;198(2):188-92.
86. Koman A. Primary hyperparathyroidism: nonclassical symptoms and benefits from parathyroidectomy: Karolinska Institutet (Sweden); 2021.
87. Kumar SR, Lakshmi KB, Reddy KVT, Deepika BA, Teja VV. Effect of open minimally invasive parathyroidectomy in the management of primary hyperparathyroidism. *International Surgery Journal*. 2017;4(11):3660-4.
88. Lala M. Management of Primary Hyperparathyroidism. *Indian journal of surgical oncology*. 2022;13(1):143-51.
89. MacRobert NA. Single gland versus multigland disease in primary hyperparathyroidism at the Wits Donald Gordon Medical Centre in Johannesburg, South Africa.
90. Madorin CA, Owen R, Coakley B, Lowe H, Nam K-H, Weber K, et al. Comparison of radiation exposure and cost between dynamic computed tomography and sestamibi scintigraphy for preoperative localization of parathyroid lesions. *JAMA surgery*. 2013;148(6):500-3.
91. McLean T, Delbridge L. Comparison of consumer information on the internet to the current evidence base for minimally invasive parathyroidectomy. *World journal of surgery*. 2010;34:1304-11.
92. McWilliams JM, Schwartz AL. Focusing on High-cost Patients: the Key to Addressing High Costs? *The New England journal of medicine*. 2017;376(9):807.
93. Mihai R, Barczynski M, Iacobone M, Sitges-Serra A. Surgical strategy for sporadic primary hyperparathyroidism an evidence-based approach to surgical strategy, patient selection, surgical access, and reoperations. *Langenbeck's archives of surgery*. 2009;394:785-98.
94. Mohebbati A, Shaha AR. Imaging techniques in parathyroid surgery for primary hyperparathyroidism. *American journal of otolaryngology*. 2012;33(4):457-68.
95. Morks A, Van Ginhouw T, Pekelharing J, Duschek E, Smit P, De Graaf P. Intra-operative parathyroid hormone measurements—experience of a non-academic hospital. *South African Journal of Surgery*. 2011;49(3):123-7.
96. Noureldine SI, Abbas A, Tufano RP, Srivastav S, Slakey DP, Friedlander P, et al. The impact of surgical volume on racial disparity in thyroid and parathyroid surgery. *Annals of surgical oncology*. 2014;21:2733-9.
97. Noureldine SI, Gooi Z, Tufano RP. Minimally invasive parathyroid surgery. *Gland surgery*. 2015;4(5):410.
98. O'Connell DA, Diamond C, Seikaly H, Harris JR. Objective and subjective scar aesthetics in minimal access vs conventional access parathyroidectomy and thyroidectomy surgical procedures: a paired cohort study. *Archives of otolaryngology—head & neck surgery*. 2008;134(1):85-93.
99. Paruk IM, Esterhuizen TM, Maharaj S, Pirie FJ, Motala AA. Characteristics, management and outcome of primary hyperparathyroidism in South Africa: a single-centre experience. *Postgraduate medical journal*. 2013;89(1057):626-31.
100. Pradeep P, Jayashree B, Mishra A, Mishra S. Systematic review of primary hyperparathyroidism in India: the past, present, and the future trends. *International journal of Endocrinology*. 2011;2011.
101. Pradhan R, Gupta S, Agarwal A. Focused parathyroidectomy using accurate preoperative imaging and intraoperative PTH: Tertiary care experience. *Indian Journal of Endocrinology and Metabolism*. 2019;23(3):347.
102. Puthenveetil P, Panchangam RB. Asymptomatic vs Symptomatic Primary Hyperparathyroidism: Comparison of Clinico-investigative Profile and Surgical Outcomes in Resource-Limited Setting. *Indian Journal of Surgery*. 2022;84(1):100-3.
103. Rago R, Forfori F, Frustaci G, Monzani R, Paracchini S, Franceschini F, et al. Day case parathyroidectomy: is this the right way for the patients? *Gland Surgery*. 2020;9(Suppl 1):S6.
104. Rajan S, Ravindhran B, George B, Bantwal G, Ayyar V, Manjuanth S. Simplified intraoperative parathormone assay for primary hyperparathyroidism in a resource-limited setting. *Biomarkers in medicine*. 2021;15(5):331-6.
105. Rawat A, Grover M, Kataria T, Samdhani S, Mathur S, Sharma B. Minimally Invasive Parathyroidectomy as the Surgical Management of Single Parathyroid Adenomas: A Tertiary Care Experience. *Indian Journal of Otolaryngology and Head & Neck Surgery*. 2022:1-7.
106. Sen S, Cherian AJ, Ramakant P, Reka K, Paul M, Abraham DT. Focused parathyroidectomy under local anesthesia—A feasibility study. *Indian Journal of Endocrinology and Metabolism*. 2019;23(1):67.
107. Serra C, Silveira L, Canudo A. Development of a low-cost portable device for identification of parathyroid glands using autofluorescence. *Indian Journal of Surgery*. 2020;82(6):1126-30.
108. Shapey I, Jabbar S, Khan Z, Nicholson J, Watson R. Scan-directed mini-incision focused parathyroidectomy: how accurate is accurate enough? *The Annals of The Royal College of Surgeons of England*. 2017;99(2):123-8.
109. Siddiqui MI, Pasha HA, Asad R, Talati JJ. Changing paradigms in the surgical management of hyperparathyroidism at a tertiary care hospital in a developing country. *JPMA*. 2019.
110. Suliburk J, Sywak M, Sidhu S, Delbridge L. ES11\* 10 YEAR HISTORY OF OVER 1000 CONSECUTIVE MINIMALLY INVASIVE PARATHYROIDECTOMIES (MIPS) WITHOUT INTRAOPERATIVE PTH AT A SINGLE INSTITUTION: LESSONS LEARNED. *ANZ Journal of Surgery*. 2009;79:A20-A.
111. Terris DJ, Duke WS, Pasieka JL. Parathyroid Surgery.
112. Traini E, Bellantone R, Tempera SE, Russo S, De Crea C, Lombardi CP, et al. Is parathyroidectomy safe and effective in patients with normocalcemic primary hyperparathyroidism? *Langenbeck's Archives of Surgery*. 2018;403:317-23.
113. Udelsman R, Åkerström G, Biagini C, Duh Q-Y, Miccoli P, Niederle B, et al. The surgical management of asymptomatic primary hyperparathyroidism: proceedings of the Fourth International Workshop. *The Journal of Clinical Endocrinology & Metabolism*. 2014;99(10):3595-606.
114. UK NGC. Evidence review for surgical localisation. 2019.
115. Usta A, Alhan E, Cinel A, Türkyilmaz S, Erem C. A 20-year study on 190 patients with primary hyperparathyroidism in a developing country: Turkey experience. *International Surgery*. 2015;100(4):648-55.
116. Van Wyngaard T. Pre-operative localization and surgical outcomes for primary hyperparathyroidism (PHPT): An 11-year review at a South African hospital: University of Cape Town; 2018.
117. Walker MD, Silverberg SJ. Primary hyperparathyroidism. *Nature Reviews Endocrinology*. 2018;14(2):115-25.
118. Wang R, Disharoon M, Frazier R, Xie R, Moses C, Gillis A, et al. Less Is More: Parathyroidectomy and Association with Postoperative Hypocalcemia in Dialysis Patients. *Journal of the American College of Surgeons*. 2023;236(4):639-45.
119. Wiseman SM, Saleh N, Tootooni A, Eshraghi P, Jama R, Saleh S. Parathyroid identification during thyroid and parathyroid operations: A pilot study evaluating a novel low cost autofluorescence based device. *The American Journal of Surgery*. 2021;221(6):1150-8.
120. Wojtczak B, Syrycka J, Kaliszewski K, Rudnicki J, Bolanowski M, Barczyński M. Surgical implications of recent modalities for parathyroid imaging. *Gland surgery*. 2020;9(Suppl 2):S86.
121. Wong IY-H, Lang BH-H. Minimally Invasive Parathyroidectomy for Primary Hyperparathyroidism—Current Views, Issues and Controversies. Edited by Gonzalo Diaz-Soto and Manuel Puig-Domingo. 2012:21.
122. Yadav SK, Mishra SK, Mishra A, Mayilvagnan S, Chand G, Agarwal G, et al. Changing profile of primary hyperparathyroidism over two and half decades: a study in tertiary referral center of North India. *World Journal of Surgery*. 2018;42:2732-7.
123. YP S. Surgical Management of Primary Hyperparathyroidism: An Institutional Study on Surgical Quality Control. *Journal of Institute of Medicine*. 2017;39(2).
124. Zachariah SK. Management of Primary Hyperparathyroidism: 'Past, Present and Future'. *Thyroid and parathyroid diseases—new insights into some old and some new issues*. 2012:147.
125. Abbas Y, Shahed QM, Quraishi MMS. Outcomes of minimally invasive parathyroidectomy with dual-modality imaging. *TriVantage| APS| NIM* 30. 2014;7(1):144.
126. Adam MA, Roman SA, Sosa JA. Outpatient parathyroidectomy. *Medical and Surgical Treatment of Parathyroid Diseases: An Evidence-Based Approach*. 2017:469-81.
127. Ahn D, Kwak JH. Role and Recent Trend of Intraoperative Parathyroid Hormone Monitoring During Parathyroidectomy in Patients With Primary Hyperparathyroidism. *Korean Journal of Otorhinolaryngology—Head and Neck Surgery*. 2022;65(5):253-9.
128. Barczynski M. Minimally invasive parathyroidectomy without intraoperative parathyroid hormone monitoring: When and why? *Journal of Postgraduate Medicine*. 2009;55(4):239.
129. Barczyński M, Golkowski F, Nawrot I. The current status of intraoperative iPTH assay in surgery for primary hyperparathyroidism. *Gland surgery*. 2015;4(1):36.
130. Barczynski M, Konturek A, Cichon S, Hubalewska-Dydejczyk A, Golkowski F, Huszno B. Intraoperative parathyroid hormone assay improves outcomes of minimally invasive parathyroidectomy mainly in patients with a presumed solitary parathyroid adenoma and missing concordance of preoperative imaging. *Clinical Endocrinology*. 2007;66(6):878-85.
131. Buzanakov DM, Slepsov IV, Semenov AA, Chernikov RA, Novokshonov KY, Kareлина YV, et al. Persistence of primary hyperparathyroidism: a single-center experience. *Langenbeck's Archives of Surgery*. 2022;407(8):3651-9.
132. Calò PG, Pisano G, Loi G, Medas F, Barca L, Atzeni M, et al. Intraoperative parathyroid hormone assay during focused parathyroidectomy: the importance of 20 minutes measurement. *BMC surgery*. 2013;13:1-5.
133. Calò PG, Pisano G, Tatti A, Medas F, Boi F, Mariotti S, et al. Intraoperative parathyroid hormone assay during focused parathyroidectomy for primary hyperparathyroidism: is it really mandatory? *Minerva Chirurgica*. 2012;67(4):337.
134. Carling T, Udelsman R. Focused approach to parathyroidectomy. *World journal of surgery*. 2008;32:1512-7.
135. Christakis I, Palazzo F. Minimally Invasive Parathyroidectomy. *Tips and Tricks in Endocrine Surgery*: Springer; 2013. p. 259-66.
136. Cvasciuc T, Lansdown M, Fraser S. Focused Parathyroidectomy without Intraoperative Parathyroid Hormone Testing: Acceptability after Preoperative Localization with SPECT-CT. *World*. 2019;11(1):2.

- 137 Deutmeyer C, Weingarten M, Doyle M, Carneiro-Pla D. Case series of targeted parathyroidectomy with surgeon-performed ultrasonography as the only preoperative imaging study. *Surgery*. 2011;150(6):1153-60.
- 138 Duke WS, Terris DJ. Minimally Invasive Non-endoscopic Parathyroidectomy. *Minimally Invasive and Robotic Thyroid and Parathyroid Surgery*. 2014:131-41.
- 139 Dyas AR, Lovell KM, Balentine CJ, Wang TN, Porterfield Jr JR, Chen H, et al. Reducing cost and improving operating room efficiency: examination of surgical instrument processing. *Journal of Surgical Research*. 2018;229:15-9.
- 140 Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *Surgical Clinics*. 2014;94(3):607-23.
- 141 Jang S, Mandabach M, Aburjania Z, Balentine CJ, Chen H. Wide variation in cost of surgical care for parathyroidectomy: is there a need for standardization of practice? *Surgery*. 2018;163(3):638-42.
- 142 Jinih M, O'Connell E, O'Leary DP, Liew A, Redmond HP. Focused versus bilateral parathyroid exploration for primary hyperparathyroidism: a systematic review and meta-analysis. *Annals of surgical oncology*. 2017;24:1924-34.
- 143 Karipineni F, Sahli Z, Somervell H, Mathur A, Prescott JD, Tufano RP, et al. Are preoperative sestamibi scans useful for identifying ectopic parathyroid glands in patients with expected multigland parathyroid disease? *Surgery*. 2018;163(1):35-41.
- 144 Kuzu F, Arpaci D, Cakmak GK, Emre AU, Elri T, Ilikhan SJ, et al. Focused parathyroidectomy without intra-operative parathormone monitoring: The value of PTH assay in preoperative ultrasound guided fine needle aspiration washout. *Annals of Medicine and Surgery*. 2016;6:64-7.
- 145 Lee GS, McKenzie TJ, Mullan BP, Farley DR, Thompson GB, Richards ML. A multimodal imaging protocol, 123 I/99 Tc-sestamibi, SPECT, and SPECT/CT, in primary hyperparathyroidism adds limited benefit for preoperative localization. *World journal of surgery*. 2016;40:589-94.
- 146 Lubitz CC, Chen H. Sestamibi-negative patients: to operate or image? *Annals of surgical oncology*. 2012;19:2086-7.
- 147 Melck AL, Armstrong MJ, Yip L, Carty SE. Case-controlled comparison of video-assisted and conventional minimally invasive parathyroidectomy. *The American Surgeon*. 2012;78(1):125-32.
- 148 Miccoli P, Berti P, Materazzi G, Ambrosini C, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTHa) during endoscopic parathyroidectomy: a prospective randomized trial. *Surgical endoscopy*. 2008;22:398-400.
- 149 Mihai R, Weisters M, Stechman MJ, Gleeson F, Sadler G. Cost-effectiveness of scan-directed parathyroidectomy. *Langenbeck's Archives of Surgery*. 2008;393:739-43.
- 150 Minuto M, Vera L, Ansaldo GL, Bargetto G, Cafiero F, Varaldo E. Modern approach to parathyroidectomy. *Giornale Italiano di Nefrologia: Organo Ufficiale Della Societa Italiana di Nefrologia*. 2014;31(6):gin/31.6. 10-gin/31.6. 10.
- 151 Miyabe R. Three-dimensional ultrasonography before minimally invasive focused parathyroidectomy: the importance of coronal images. *Surgery today*. 2009;39:98-103.
- 152 Morris LF, Zanocco K, Ituarte PH, Ro K, Duh Q-Y, Sturgeon C, et al. The value of intraoperative parathyroid hormone monitoring in localized primary hyperparathyroidism: a cost analysis. *Annals of surgical oncology*. 2010;17:679-85.
- 153 Mownah O, Pafitanis G, Drake W, Crinnion J. Contemporary surgical treatment of primary hyperparathyroidism without intraoperative parathyroid hormone measurement. *The Annals of The Royal College of Surgeons of England*. 2015;97(8):603-7.
- 154 Norlén O, Wang KC, Tay YK, Johnson WR, Grodski S, Yeung M, et al. No need to abandon focused parathyroidectomy: a multicenter study of long-term outcome after surgery for primary hyperparathyroidism. *Annals of surgery*. 2015;261(5):991-6.
- 155 Nowlin WF. Parathyroidectomy. *Advanced Surgical Techniques for Rural Surgeons*. 2015:147-55.
- 156 Opoku-Boateng A, Bolton JS, Corsetti R, Brown R, Oxner C, Fuhrman GM. Use of a sestamibi-only approach to routine minimally invasive parathyroidectomy. *The American Surgeon*. 2013;79(8):797-801.
- 157 Paek SH, Kim S-J, Choi JY, Lee KE. Clinical usefulness of intraoperative parathyroid hormone monitoring for primary hyperparathyroidism. *Annals of Surgical Treatment and Research*. 2018;94(2):69-73.
- 158 Pang T, Stalberg P, Sidhu S, Sywak M, Wilkinson M, Reeve T, et al. Minimally invasive parathyroidectomy using the lateral focused mini-incision technique without intraoperative parathyroid hormone monitoring. *Journal of British Surgery*. 2007;94(3):315-9.
- 159 Phitayakorn R, McHenry CR. Parathyroidectomy: overview of the anatomic basis and surgical strategies for parathyroid operations. *Clinical Reviews in Bone and Mineral Metabolism*. 2007;5:89-102.
- 160 Prasanna S, Davies G, Bochner M, Kollias J, Malycha P. Minimally invasive parathyroidectomy using surgeon-performed ultrasound and sestamibi. *ANZ Journal of Surgery*. 2007;77(9):774-7.
- 161 Rajaei MH, Oltmann SC, Adkisson CD, Elfelbein DM, Chen H, Carty SE, et al. Is intraoperative parathyroid hormone monitoring necessary with ipsilateral parathyroid gland visualization during anticipated unilateral exploration for primary hyperparathyroidism: a two-institution analysis of more than 2,000 patients. *Surgery*. 2014;156(4):760-8.
- 162 Sadacharan D, Agarwal G. Re-operative parathyroidectomy: An algorithm for imaging and monitoring of intraoperative parathyroid hormone levels that results in a successful focused approach. *Surgery*. 2009;146(3):524.
- 163 Sartori PV, Saibene AM, Leopaldi E, Boniardi M, Beretta E, Colombo S, et al. Intraoperative parathyroid hormone testing in primary hyperparathyroidism surgery: time for giving up? *European Archives of Oto-Rhino-Laryngology*. 2019;276:267-72.
- 164 Stalberg P, Delbridge L, Van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy—current concepts. *The Surgeon*. 2007;5(5):301-8.
- 165 Stojadinovic A, Pribitkin E, Rosen D, Edwards M, Byrd DR. Unilateral vs bilateral parathyroidectomy: a healthy debate. *Journal of the American College of Surgeons*. 2012;215(2):300-2.
- 166 Tublin ME, Pryma DA, Yim JH, Ogilvie JB, Mountz JM, Bencherif B, et al. Localization of parathyroid adenomas by sonography and technetium tc 99m sestamibi single-photon emission computed tomography before minimally invasive parathyroidectomy: are both studies really needed? *Journal of Ultrasound in Medicine*. 2009;28(2):183-90.
- 167 Wang TS, Cheung K, Farrokhyar F, Roman SA, Sosa JA. Would scan, but which scan? A cost-utility analysis to optimize preoperative imaging for primary hyperparathyroidism. *Surgery*. 2011;150(6):1286-94.
- 168 Werenski HE, Nguyen CJ, Johansson ED, Bunch PM, Randle RW. Value of Old Imaging for Patients Undergoing Parathyroidectomy for Primary Hyperparathyroidism. *Journal of Surgical Research*. 2023;282:147-54.
- 169 Westerdahl J, Bergenfelz A. Focused Parathyroidectomy. *Endocrine Surgery: Principles and Practice*. 2009:267-78.
- 170 Wong W, Foo FJ, Lau M, Sarin A, Kiruparan P. Simplified minimally invasive parathyroidectomy: a series of 100 cases and review of the literature. *The Annals of The Royal College of Surgeons of England*. 2011;93(4):290-3.
- 171 Yalçın O, Günay S. Focused Parathyroid Surgery: Minimally Invasive. *Thyroid and Parathyroid Diseases: A Case-Based Guide*. 2018:463.
- 172 Yalçın O, Günay S. Focused Parathyroid Surgery: Minimally Invasive Parathyroidectomy via a Mini-incision. *Thyroid and Parathyroid Diseases: A Case-Based Guide*. 2019:463-70.
- 173 Ypsilantis E, Charfare H, Wassif W. Intraoperative PTH assay during minimally invasive parathyroidectomy may be helpful in the detection of double adenomas and may minimise the risk of recurrent surgery. *International journal of endocrinology*. 2010;2010.
- 174 Zammit M, Pierce K, Bailey L, Rowland M, Waghorn A, Shore S. Challenging NICE guidelines on parathyroid surgery. *The Surgeon*. 2022;20(4):e105-e111.
- 175 Zanocco KA, Wu JX, Yeh MW. Parathyroidectomy for asymptomatic primary hyperparathyroidism: a revised cost-effectiveness analysis incorporating fracture risk reduction. *Surgery*. 2017;161(1):16-24.
- 176 Van Wyngaert T. Pre-operative localization and surgical outcomes for primary hyperparathyroidism (PHPT): An 11-year review at a South African hospital: University of Cape Town; 2018.
- 177 Alikor CA, Engelbrecht G. The Usefulness of 99m technetium-sestamibi Parathyroid Scintigraphy in Preoperative Localization of Parathyroid Adenoma in Patients with Primary Hyperparathyroidism at an Academic Hospital in South Africa.
- 178 Dumzela A, Bombil I. PROFILE OF PATIENTS OPERATED ON FOR PRIMARY HYPERPARATHYROIDISM AT CHRIS HANI BARAGWANATH ACADEMIC HOSPITAL. *Rheumatology*. 2013;17:1.
- 179 Sathe PA, Madiwale CV, Kandalkar BM, Bandgar TR, Shah NS, Menon PS. Primary hyperparathyroidism: a clinicopathological experience. *Indian Journal of Pathology and Microbiology*. 2009;52(3):313.
- 180 Bombil I, Louw L, Mitchell C, Mahlobo F, Muganza RA, Madima NR. Sonar guided focused parathyroidectomy under cervical block. *South African Journal of Surgery*. 2018;56(2):30-3.
- 181 KP<sup>1</sup> P, Bombil I, Mpanya D. Ectopic Parathyroid Adenoma Localized by Tc-99m Sestamibi SPECT/CT Localization Prior to Re-operation is Useful.
- 182 Medas F, Cappellacci F, Canu GL, Noordzij JP, Erdas E, Calo PG. The role of Rapid Intraoperative Parathyroid Hormone (ioPTH) assay in determining outcome of parathyroidectomy in primary hyperparathyroidism: A systematic review and meta-analysis. *International Journal of Surgery*. 2021;92:106042.
- 183 Lee S-W, Shim SR, Jeong SY, Kim S-J. Direct comparison of preoperative imaging modalities for localization of primary hyperparathyroidism: a systematic review and network meta-analysis. *JAMA Otolaryngology—Head & Neck Surgery*. 2021;147(8):692-706.
- 184 Wetzig N. ES03 GOITRE SURGERY IN AFRICA. *ANZ Journal of Surgery*. 2009;79:A18-A.
- 185 Doruyter AGG. Imaging of renal hyperparathyroidism using SPECT/CT with low-dose localizing CT: Stellenbosch: Stellenbosch University; 2013.
- 186 Yorke E, Atiase Y, Akpalu J. Primary Hyperparathyroidism: an Update and In-Depth Review of the Literature. *Int J Endocrinol Metab Disord*. 2018;4(2).
- 187 Malycha P. ES02 PARATHYROID SURGERY. A TRUE MARRIAGE OF SKILL, TECHNOLOGY AND SCIENCE. *ANZ Journal of Surgery*. 2009;79:A18-A.
- 188 Suliburk J, Sywak M, Sidhu S, Delbridge L. ES11\* 10 YEAR HISTORY OF OVER 1000 CONSECUTIVE MINIMALLY INVASIVE PARATHYROIDECTOMIES (MIPS) WITHOUT INTRAOPERATIVE PTH AT A SINGLE INSTITUTION: LESSONS LEARNED. *ANZ Journal of Surgery*. 2009;79:A20-A.
- 189 UK NGC. Evidence review for surgical localisation. 2019.
- 190 Petranović Ovcariček P, Giovannella L, Carrió Gasset I, Hindié E, Huellner MW, Luster M, et al. The EANM practice guidelines for parathyroid imaging. *European journal of nuclear medicine and molecular imaging*. 2021;48:2801-22.

- 191 Medas F, Cappellacci F, Canu GL, Noordzij JP, Erdas E, Calo PG. The role of Rapid Intraoperative Parathyroid Hormone (ioPTH) assay in determining outcome of parathyroidectomy in primary hyperparathyroidism: A systematic review and meta-analysis. *International Journal of Surgery*. 2021;92:106042.
- 192 Morks A, Van Ginhoven T, Pekelharing J, Duschek E, Smit P, De Graaf P. Intra-operative parathyroid hormone measurements—experience of a non-academic hospital. *South African Journal of Surgery*. 2011;49(3):123-7.
- 193 Alagaratnam S, Kurzawinski TR. Aetiology, diagnosis and surgical treatment of primary hyperparathyroidism in children: new trends. *Hormone research in paediatrics*. 2015;83(6):365-75.
- 194 Barnes W, Czako PF, Nagar S. Long-Term Success of Surgery for Primary Hyperparathyroidism: Focused Exploration using Intraoperative Parathyroid Hormone Monitoring Versus Four-Gland Exploration. *Difficult Decisions in Endocrine Surgery: An Evidence-Based Approach*. 2018:239-71.
- 195 Maharjan R, Maharjan R, Arun K. STUDY OF CLINICAL FEATURES, SURGICAL MANAGEMENT AND OUTCOMES OF PARATHYROID ADENOMA. *Nepalese Journal of ENT Head & Neck Surgery*. 2018;9(2):13-8.
- 196 Jabiev AA, Lew JJ, Garb JL, Sanchez YM, Solorzano CC. Primary hyperparathyroidism in the underinsured: A study of 493 patients. *Surgery*. 2012;151(3):471-6.
- 197 Al-Thani H, El-Matbouly M, Al-Sulaiti M, Asim M, Majzoub A, Tabeb A, et al. Management and outcomes of hyperparathyroidism: a case series from a single institution over two decades. *Therapeutics and Clinical Risk Management*. 2018:1337-45.
- 198 Morks A, Van Ginhoven T, Pekelharing J, Duschek E, Smit P, De Graaf P. Intra-operative parathyroid hormone measurements—experience of a non-academic hospital. *South African Journal of Surgery*. 2011;49(3):123-7.
- 199 Sardiwalla II, Mokhtari A, Sardiwalla Y. Parathyroid adenoma with concurrent toxic thyroid adenoma: a rare combination. *South African Journal of Surgery*. 2017;55(1):41-4.
- 200 Oltmann SC, Rajaei MH, Sippel RS, Chen H, Schneider DF. Primary hyperparathyroidism across the ages: presentation and outcomes. *Journal of Surgical Research*. 2014;190(1):185-90.
- 201 Alikor CA, Engelbrecht G. The Usefulness of 99m technetium-sestamibi Parathyroid Scintigraphy in Preoperative Localization of Parathyroid Adenoma in Patients with Primary Hyperparathyroidism at an Academic Hospital in South Africa.
- 202 Barnes W, Czako PF, Nagar S. Long-Term Success of Surgery for Primary Hyperparathyroidism: Focused Exploration using Intraoperative Parathyroid Hormone Monitoring Versus Four-Gland Exploration. *Difficult Decisions in Endocrine Surgery: An Evidence-Based Approach*. 2018:239-71.
- 203 Barnes W, Czako PF, Nagar S. Long-Term Success of Surgery for Primary Hyperparathyroidism: Focused Exploration using Intraoperative Parathyroid Hormone Monitoring Versus Four-Gland Exploration. *Difficult Decisions in Endocrine Surgery: An Evidence-Based Approach*. 2018:239-71.
- 204 Walker MD, Silverberg SJ. Primary hyperparathyroidism. *Nature Reviews Endocrinology*. 2018;14(2):115-25.
- 205 Bilezikian JP, Cusano NE, Khan AA, Liu J-M, Marcocci C, Bandeira F. Primary hyperparathyroidism. *Nature Reviews Disease Primers*. 2016;2(1):1-16.
- 206 Iputo R. Parathyroid disease. *Southern African Journal of Anaesthesia and Analgesia*. 2021;27(6):86-9.
- 207 MacRobert NA. Single gland versus multigland disease in primary hyperparathyroidism at the Wits Donald Gordon Medical Centre in Johannesburg, South Africa.
- 208 Wiseman SM, Saleh N, Tootooni A, Eshraghi P, Jama R, Saleh S. Parathyroid identification during thyroid and parathyroid operations: A pilot study evaluating a novel low cost autofluorescence based device. *The American Journal of Surgery*. 2021;221(6):1150-8.
- 209 Abbott DE, Cantor SB, Grubbs EG, Santora R, Gomez HF, Evans DB, et al. Outcomes and economic analysis of routine preoperative 4-dimensional CT for surgical intervention in de novo primary hyperparathyroidism: does clinical benefit justify the cost? *Journal of the American College of Surgeons*. 2012;214(4):629-37.
- 210 Mokoena T. Intra-operative serum calcium monitoring compares favourably with parathyroid hormone monitoring to determine completeness of surgery for parathyroid adenoma. 2020.
- 211 Shawky MS. Intraoperative parathyroid hormone monitoring in primary hyperparathyroidism; resolving controversies and debates. *Journal of Parathyroid Disease*. 2016;5(1):3-10.
- 212 Paruk IM, Esterhuizen TM, Maharaj S, Pirie FJ, Motala AA. Characteristics, management and outcome of primary hyperparathyroidism in South Africa: a single-centre experience. *Postgrad Med J*. 2013;89(1057):626-31.
- 213 Alikor C, Engelbrecht G. The Usefulness of 99mtechnetium-sestamibi Parathyroid Scintigraphy in Preoperative Localization of Parathyroid Adenoma in Patients with Primary Hyperparathyroidism at an Academic Hospital in South Africa. *Asian Journal of Medicine and Health*. 2017;6:1-7.
- 214 Akgün IE, Ünlü MT, Aygun N, Kostek M, Uludag M. Contribution of intraoperative parathyroid hormone monitoring to the surgical success in minimal invasive parathyroidectomy. *Frontiers in Surgery*. 2022;9.
- 215 Patel KN, Caso R. Intraoperative parathyroid hormone monitoring: optimal utilization. *Surgical Oncology Clinics*. 2016;25(1):91-101.
- 216 Ahmed R, Khan N, Ellemidin S, Gayaparsad K. Ultrasound-first imaging modality in the detection of parathyroid adenomas. *SA Journal of Radiology*. 2008;12(3):64-8
- 217 Koyuncu A, Ozer SP, Ozer B, Catal O, Sit M. Minimally invasive surgery in primary hyperparathyroidism. *Journal of Bionic Memory*. 2023 Mar 5;3(1):1-6.

#### Pubmed

1. Herb J, Staley BS, Roberson M, Strassle PD, Kim LT. Use and disparities in parathyroidectomy for symptomatic primary hyperparathyroidism in the Medicare population. *Surgery*. 2021;170(5):1376-82.
2. Alameer E, Omar M, Hoof M, Shalaby H, Abdelgawad M, Zora G, et al. Effects of Parathyroidectomy on Normocalcemic Primary Hyperparathyroidism and the Role of Intraoperative PTH Measurement. *Am Surg*. 2022;88(5):873-9.
3. Herb J, Staley BS, Roberson M, Strassle PD, Kim LT. Use and disparities in parathyroidectomy for symptomatic primary hyperparathyroidism in the Medicare population. *Surgery*. 2021;170(5):1376-82.
4. Kowalski GJ, Buła G, Źądło D, Gawrychowska A, Gawrychowski J. Primary hyperparathyroidism. *Endokrynol Pol*. 2020;71(3):260-70.
5. Singh Ospina NM, Rodriguez-Gutierrez R, Maraka S, Espinosa de Ycaza AE, Jasim S, Castaneda-Guarderas A, et al. Outcomes of Parathyroidectomy in Patients with Primary Hyperparathyroidism: A Systematic Review and Meta-analysis. *World J Surg*. 2016;40(10):2359-77.
6. Walker MD, Silverberg SJ. Primary hyperparathyroidism. *Nat Rev Endocrinol*. 2018;14(2):115-25.
7. Managing primary hyperparathyroidism in primary care. *Drug Ther Bull*. 2010;48(3):30-3.
8. Aarum S, Nordenström J, Reihné E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for primary hyperparathyroidism: the new versus the old order. A randomised controlled trial of preoperative localisation. *Scand J Surg*. 2007;96(1):26-30.
9. Abbott DE, Cantor SB, Grubbs EG, Santora R, Gomez HF, Evans DB, et al. Outcomes and economic analysis of routine preoperative 4-dimensional CT for surgical intervention in de novo primary hyperparathyroidism: does clinical benefit justify the cost? *J Am Coll Surg*. 2012;214(4):629-37; discussion 37-9.
10. Adil E, Adil T, Fedok F, Kauffman G, Goldenberg D. Minimally invasive radioguided parathyroidectomy performed for primary hyperparathyroidism. *Otolaryngol Head Neck Surg*. 2009;141(1):34-8.
11. Adler JT, Sippel RS, Chen H. New trends in parathyroid surgery. *Curr Probl Surg*. 2010;47(12):958-1017.
12. Agrawal R, Mayilvaganan S. Primary hyperparathyroidism: an analysis of failure of parathyroidectomy. *World J Surg*. 2014;38(9):2481-2.
13. Ahmadiéh H, Kreidieh O, Akl EA, El-Hajj Fuleihan G. Minimally invasive parathyroidectomy guided by intraoperative parathyroid hormone monitoring (IOPH) and preoperative imaging versus bilateral neck exploration for primary hyperparathyroidism in adults. *Cochrane Database Syst Rev*. 2020;10(10):Cd010787.
14. Al-Azem H, Khan A. Primary hyperparathyroidism. *Cmaj*. 2011;183(10):E685-9.
15. Alfadda A, Hagr A, Al-Qahtani K, Tabah R. Radio-guided minimally invasive parathyroidectomy under local anesthesia. *West Afr J Med*. 2006;25(2):134-7.
16. Alobuia WM, Meng T, Cisco RM, Lin DT, Suh I, Tamura MK, et al. Racial disparities in the utilization of parathyroidectomy among patients with primary hyperparathyroidism: Evidence from a nationwide analysis of Medicare claims. *Surgery*. 2022;171(1):8-16.
17. Alore EA, Massarweh NN, Ramsey DJ, Chen L, Chai CY, Singh H, et al. Variation in surgical management of primary hyperparathyroidism in the US Department of Veterans Affairs healthcare system: A 15-year observational study. *Surgery*. 2020;168(5):838-44.
18. Anagnostis P, Vaitis K, Veneti S, Potoupni V, Kenanidis E, Tsridis E, et al. Efficacy of parathyroidectomy compared with active surveillance in patients with mild asymptomatic primary hyperparathyroidism: a systematic review and meta-analysis of randomized-controlled studies. *J Endocrinol Invest*. 2021;44(6):1127-37.
19. Augustine MM, Bravo PE, Zeiger MA. Surgical treatment of primary hyperparathyroidism. *Endocr Pract*. 2011;17 Suppl 1:75-82.

20. Axelsson KF, Wallander M, Johansson H, Harvey NC, Vandenput L, McCloskey E, et al. Analysis of Comorbidities, Clinical Outcomes, and Parathyroidectomy in Adults With Primary Hyperparathyroidism. *JAMA Netw Open*. 2022;5(6):e2215396.
21. Balasubramanian SP, Harrison BJ. Simplified minimally invasive parathyroidectomy. *Ann R Coll Surg Engl*. 2011;93(7):563-4.
22. Baliski C, Nosyk B, Melck A, Bugis S, Rosenberg F, A HA. The cost-effectiveness of three strategies for the surgical treatment of symptomatic primary hyperparathyroidism. *Ann Surg Oncol*. 2008;15(10):2653-60.
23. Becker JL, Patel V, Johnson KJ, Guerrero M, Klein RR, Ranvier GF, et al. 4D-Dynamic Contrast-Enhanced MRI for Preoperative Localization in Patients with Primary Hyperparathyroidism. *AJNR Am J Neuroradiol*. 2020;41(3):522-8.
24. Bergenfelz A, van Slycke S, McKay O, Brunaud L. Author response to: European multicentre study on outcome of surgery for sporadic primary hyperparathyroidism. *Br J Surg*. 2021;108(12):e416.
25. Beyer TD, Solorzano CC, Starr F, Nilubol N, Prinz RA. Parathyroidectomy outcomes according to operative approach. *Am J Surg*. 2007;193(3):368-72; discussion 72-3.
26. Bhangu JS, Riss P. The role of intraoperative parathyroid hormone (IOPTH) determination for identification and surgical strategy of sporadic multiglandular disease in primary hyperparathyroidism (pHPT). *Best Pract Res Clin Endocrinol Metab*. 2019;33(5):101310.
27. Body JJ. [Primary hyperparathyroidism: diagnosis and management]. *Rev Med Brux*. 2012;33(4):263-7.
28. Bolland MJ, Grey A. Nonoperative Management of Mild Primary Hyperparathyroidism: A Reasonable, Evidence-Based Option. *Ann Intern Med*. 2022;175(6):899-900.
29. Buicko JL, Kichler KM, Amundson JR, Scurci S, Kozol RA. The Sestamibi Paradox: Improving Intraoperative Localization of Parathyroid Adenomas. *Am Surg*. 2017;83(8):832-5.
30. Bunch PM, Goyal A, Valenzuela CD, Randle RW. Parathyroid 4D CT in Primary Hyperparathyroidism: Exploration of Size Measurements for Identifying Multigland Disease and Guiding Biochemically Successful Parathyroidectomy. *AJR Am J Roentgenol*. 2022;218(5):888-97.
31. Callender GG, Udelsman R. Surgery for primary hyperparathyroidism. *Cancer*. 2014;120(23):3602-16.
32. Calò PG, Medas F, Loi G, Erdas E, Pisano G, Nicolosi A. Feasibility of unilateral parathyroidectomy in patients with primary hyperparathyroidism and negative or discordant localization studies. *Updates Surg*. 2016;68(2):155-61.
33. Calò PG, Pisano G, Tatti A, Medas F, Boi F, Mariotti S, et al. Intraoperative parathyroid hormone assay during focused parathyroidectomy for primary hyperparathyroidism: is it really mandatory? *Minerva Chir*. 2012;67(4):337-42.
34. Calzolari F, Misso C, D'Ajello M, Monacelli M, Iapadre M, Sanguinetti A, et al. [Role of minimally invasive surgery in the treatment of primary hyperparathyroidism]. *Chir Ital*. 2006;58(3):331-5.
35. Canu GL, Cappellacci F, Noordzij JP, Piras S, Erdas E, Calò PG, et al. A mini-invasive approach is feasible in patients with primary hyperparathyroidism and discordant or negative localisation studies. *Updates Surg*. 2022;74(2):747-55.
36. Cappello ZJ, Bumpous JM. Is bilateral exploration still the standard of care for primary hyperparathyroidism?: outcomes of focused radio-guided parathyroidectomy and bilateral explorations. *Laryngoscope*. 2013;123(11):2587-8.
37. Castellana M, Virilli C, Palermo A, Giorgino F, Giovanella L, Trimboli P. Primary hyperparathyroidism with surgical indication and negative or equivocal scintigraphy: safety and reliability of PTH washout. A systematic review and meta-analysis. *Eur J Endocrinol*. 2019;181(3):245-53.
38. Chandler NR, Chidambaram S, Van Den Heede K, DiMarco AN, Tolley NS, Palazzo FF. Correlation of Preoperative Imaging Findings and Parathyroidectomy Outcomes Support NICE 2019 Guidance. *J Clin Endocrinol Metab*. 2022;107(3):e1242-e8.
39. Cheren'ko SM, Tovkaï OA, Sheptukha SA. [Minimally invasive and endoscopic methods in surgical treatment of patients with primary hyperparathyroidism: advantages and limitations]. *Klin Khir*. 2012(7):16-9.
40. Choy KW. Primary Hyperparathyroidism. *N Engl J Med*. 2018;379(25):e43.
41. Dahiya D, Abuji K, Kumari P, Gautam A, Bhadada S, Sood A, et al. Surgical outcome after focused parathyroidectomy: experience from a tertiary care centre in North India. *Pol Przegl Chir*. 2021;93(5):1-5.
42. das Neves MC, Santos RO, Ohe MN. Surgery for primary hyperparathyroidism. *Arch Endocrinol Metab*. 2022;66(5):678-88.
43. Day KM, Elsayed M, Monchik JM. No Need to Abandon Focused Unilateral Exploration for Primary Hyperparathyroidism with Intraoperative Monitoring of Intact Parathyroid Hormone. *J Am Coll Surg*. 2015;221(2):518-23.
44. de la Plaza Llamas R, Ramia Àngel JM, Arteaga Peralta V. Letter to the Editor regarding "Does it have a negative economic impact the intraoperative parathyroid hormone assay in primary hyperparathyroidism?". *Head Neck*. 2018;40(2):444-5.
45. Del Rio P, Tosi G, Loderer T, Bonati E, Cozzani F, Ruffini L. Preoperative imaging evaluation in primary hyperparathyroidism and associated thyroid disease. *Ann Ital Chir*. 2021;92:471-8.
46. Di Marco A, Mechera R, Glover A, Papachristos A, Clifton-Bligh R, Delbridge L, et al. Focused parathyroidectomy without intraoperative parathyroid hormone measurement in primary hyperparathyroidism: Still a valid approach? *Surgery*. 2021;170(5):1383-8.
47. diMarco AN, Sywak MS, Sidhu SB. Primary Hyperparathyroidism. *N Engl J Med*. 2018;379(25):e43.
48. Dowthwaite SA, Young JE, Pasternak JD, Yoo J. Surgical management of primary hyperparathyroidism. *J Clin Densitom*. 2013;16(1):48-53.
49. Dralle H, Hein J. [Parathyroidectomy for unclear localization: Minimally invasive or open]. *Chirurg*. 2013;84(7):598.
50. Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *Surg Clin North Am*. 2014;94(3):607-23.
51. España M, Sastre I, Ceballos RJ, Bustos MEF. VATS parathyroidectomy for primary hyperparathyroidism. *Multimed Man Cardiothorac Surg*. 2020;2020.
52. Flynn MB, Civelek AC. Current status of surgical techniques for parathyroidectomy for untreated primary hyperparathyroidism: is the technology worth it? *Am Surg*. 2010;76(7):663-71.
53. Fraker DL, Harsono H, Lewis R. Minimally invasive parathyroidectomy: benefits and requirements of localization, diagnosis, and intraoperative PTH monitoring, long-term results. *World J Surg*. 2009;33(11):2256-65.
54. Francucci CM, Ceccoli L, Caudarella R, Rilli S, Vescini F, Boscaro M. Asymptomatic primary hyperparathyroidism: surgical and medical management. *J Endocrinol Invest*. 2011;34(7 Suppl):50-4.
55. Frank E, Watson W, Fujimoto S, De Andrade Filho P, Inman J, Simental A. Surgery versus Imaging in Non-Localizing Primary Hyperparathyroidism: A Cost-Effectiveness Model. *Laryngoscope*. 2020;130(12):E963-e9.
56. Garas G, Holsinger FC, Grant DG, Athanasiou T, Arora A, Tolley N. Is robotic parathyroidectomy a feasible and safe alternative to targeted open parathyroidectomy for the treatment of primary hyperparathyroidism? *Int J Surg*. 2015;15:55-60.
57. Gasparri G. Updates in primary hyperparathyroidism. *Updates Surg*. 2017;69(2):217-23.
58. Gencoglu EA, Aras M, Moray G, Aktas A. The effectiveness of low-dose versus high-dose 99mTc MIBI protocols for radioguided surgery in patients with primary hyperparathyroidism. *Nucl Med Commun*. 2014;35(4):398-404.
59. Giraldez-Rodríguez LA, Giraldez-Casasnovas LJ. Minimally invasive parathyroidectomy as treatment for primary hyperparathyroidism. *Bol Asoc Med P R*. 2008;100(1):27-32.
60. Gittoes NJ, Cooper MS. Primary hyperparathyroidism--is mild disease worth treating? *Clin Med (Lond)*. 2010;10(1):45-9.
61. Gollisch K, Siggelkow H. [Asymptomatic primary hyperparathyroidism : Operation or observation?]. *Internist (Berl)*. 2021;62(5):496-504.
62. Gracie D, Hussain SS. Use of minimally invasive parathyroidectomy techniques in sporadic primary hyperparathyroidism: systematic review. *J Laryngol Otol*. 2012;126(3):221-7.
63. Grady JA, Bumpous JM, Fleming MM, Flynn MB, Turbiner E, Lentsch EJ, et al. Advantages of a targeted approach in minimally invasive radioguided parathyroidectomy surgery for primary hyperparathyroidism. *Laryngoscope*. 2006;116(3):431-5.
64. Habib Z, Camacho P. Primary hyperparathyroidism: an update. *Curr Opin Endocrinol Diabetes Obes*. 2010;17(6):554-60.
65. Hamidi S, Koksai H, Kurukahvecioglu O, Taneri F. Primary hyperparathyroidism due to parathyroid adenoma. *Saudi Med J*. 2007;28(2):305; author reply -6.
66. Harari A, Allendorf J, Shifrin A, DiGorgi M, Inabnet WB. Negative preoperative localization leads to greater resource use in the era of minimally invasive parathyroidectomy. *Am J Surg*. 2009;197(6):769-73.
67. Hinnie J. The management of primary hyperparathyroidism. *Scott Med J*. 2013;58(4):251-3.
68. Hughes DT, Miller BS, Park PB, Cohen MS, Doherty GM, Gauger PG. Factors in conversion from minimally invasive parathyroidectomy to bilateral parathyroid exploration for primary hyperparathyroidism. *Surgery*. 2013;154(6):1428-34; discussion 34-5.
69. Iglesias P, Díez JJ. Current treatments in the management of patients with primary hyperparathyroidism. *Postgrad Med J*. 2009;85(999):15-23.
70. Insogna KL. Primary Hyperparathyroidism. *N Engl J Med*. 2018;379(25):e43.
71. Irkorucu O, Değer KC, Reyhan E, Arslan E. Open mini-incision parathyroidectomy for solitary parathyroid adenoma: surgical limitations. *Eur Arch Otorhinolaryngol*. 2014;271(3):625.
72. Ishii H, Stechman MJ, Watkinson JC, Aspinall S, Kim DS. A Review of Parathyroid Surgery for Primary Hyperparathyroidism from the United Kingdom Registry of Endocrine and Thyroid Surgery (UKRETS). *World J Surg*. 2021;45(3):782-9.
73. Johnson NA, Carty SE, Tublin ME. Parathyroid imaging. *Radiol Clin North Am*. 2011;49(3):489-509, vi.
74. Johnson NA, Tublin ME, Ogilvie JB. Parathyroid imaging: technique and role in the preoperative evaluation of primary hyperparathyroidism. *AJR Am J Roentgenol*. 2007;188(6):1706-15.
75. Kandil E, Malazai AJ, Alrashdeedi S, Tufano RP. Minimally invasive/focused parathyroidectomy in patients with negative sestamibi scan results. *Arch Otolaryngol Head Neck Surg*. 2012;138(3):223-5.
76. Karakas E, Schneider R, Rothmund M, Bartsch DK, Schlosser K. Initial surgery for benign primary hyperparathyroidism: an analysis of 1,300 patients in a teaching hospital. *World J Surg*. 2014;38(8):2011-8.

77. Kelly KJ, Chen H, Sippel RS. Primary hyperparathyroidism. *Cancer Treat Res.* 2010;153:87-103.
78. Khan ZF, Lew JI. Intraoperative Parathyroid Hormone Monitoring in the Surgical Management of Sporadic Primary Hyperparathyroidism. *Endocrinol Metab (Seoul).* 2019;34(4):327-39.
79. Khokar AM, Kuchta KM, Moo-Young TA, Winchester DJ, Prinz RA. Increasing trend of bilateral neck exploration in primary hyperparathyroidism. *Am J Surg.* 2020;219(3):466-70.
80. Kiernan CM, Wang T, Perrier ND, Grubbs EG, Solórzano CC. Bilateral Neck Exploration for Sporadic Primary Hyperparathyroidism: Use Patterns in 5,597 Patients Undergoing Parathyroidectomy in the Collaborative Endocrine Surgery Quality Improvement Program. *J Am Coll Surg.* 2019;228(4):652-9.
81. Kim SM, Shu AD, Long J, Montez-Rath ME, Leonard MB, Norton JA, et al. Declining Rates of Inpatient Parathyroidectomy for Primary Hyperparathyroidism in the US. *PLoS One.* 2016;11(8):e0161192.
82. Köberle R, Bendik CF. [Primary hyperparathyroidism]. *Ther Umsch.* 2020;77(9):433-40.
83. Krausz MM, Younis O, Mahamid A, Bidder M, Alfici R. [MINIMALLY INVASIVE PARATHYROIDECTOMY (MIP) UNDER LOCAL ANESTHESIA FOR TREATMENT OF PRIMARY HYPERPARATHYROIDISM (PHPT) CAUSED BY A SINGLE ADENOMA]. *Harefuah.* 2017;156(1):14-8.
84. Kunstman JW, Udelsman R. Superiority of minimally invasive parathyroidectomy. *Adv Surg.* 2012;46:171-89.
85. Laird AM, Libutti SK. Minimally Invasive Parathyroidectomy Versus Bilateral Neck Exploration for Primary Hyperparathyroidism. *Surg Oncol Clin N Am.* 2016;25(1):103-18.
86. Langusch CC, Norlen O, Titmuss A, Donoghue K, Holland AJ, Shun A, et al. Focused image-guided parathyroidectomy in the current management of primary hyperparathyroidism. *Arch Dis Child.* 2015;100(10):924-7.
87. Laxague F, Angeramo CA, Armella ED, Valinoti AC, Mezzadri NA, Fernández Vila JM. Preoperative matching studies in the diagnosis of parathyroid adenoma for primary hyperparathyroidism: Can we avoid intraoperative PTH monitoring? *Cir Esp (Engl Ed).* 2021;99(8):572-7.
88. Lecourt A, Creff G, Coudert P, De Crouy Chanel O, Guggenbuhl P, Jegoux F. Surgical management of MILD hyperparathyroidism. *Eur Arch Otorhinolaryngol.* 2021;278(10):3901-10.
89. Lew JI, Solorzano CC. Surgical management of primary hyperparathyroidism: state of the art. *Surg Clin North Am.* 2009;89(5):1205-25.
90. Lim MS, Jinih M, Ngai CH, Foley NM, Redmond HP. The utility of the radionuclide probe in parathyroidectomy for primary hyperparathyroidism. *Ann R Coll Surg Engl.* 2017;99(5):369-72.
91. Liu H, Zhang C. A commentary on "The role of rapid intraoperative parathyroid hormone (ioPTH) assay in determining outcome of parathyroidectomy in primary hyperparathyroidism: A systematic review and meta-analysis" (*Int J Surg* 2021; 92:106042). *Int J Surg.* 2021;94:106129.
92. Lundstroem AK, Trolle W, Soerensen CH, Myschetzky PS. Preoperative localization of hyperfunctioning parathyroid glands with 4D-CT. *Eur Arch Otorhinolaryngol.* 2016;273(5):1253-9.
93. Machado NN, Wilhelm SM. Diagnosis and Evaluation of Primary Hyperparathyroidism. *Surg Clin North Am.* 2019;99(4):649-66.
94. Majcen M, Hocevar M. Surgical options in treating patients with primary hyperparathyroidism. *Radiol Oncol.* 2020;54(1):22-32.
95. Marrocci C, Pinchera A. Is parathyroidectomy beneficial in patients with mild, asymptomatic primary hyperparathyroidism? *Nat Clin Pract Endocrinol Metab.* 2007;3(11):727.
96. Meier C. Benefits and place of calcimimetics in the management of primary hyperparathyroidism. *Ann Endocrinol (Paris).* 2015;76(2):163-4.
97. Mekel M, Levit B, Kluger Y, Ish-Shalom S, Segal E, Bishara B. [SHOULD PATIENTS WITH PRIMARY HYPERPARATHYROIDISM AND NEGATIVE IMAGING STUDIES BE REFERRED TO SURGERY?]. *Harefuah.* 2017;156(9):578-81.
98. Melfa GI, Raspanti C, Attard M, Cocorullo G, Attard A, Mazzola S, et al. Comparison of minimally invasive parathyroidectomy under local anaesthesia and minimally invasive video-assisted parathyroidectomy for primary hyperparathyroidism: a cost analysis. *G Chir.* 2016;37(2):61-7.
99. Miccoli P, Berti P, Materazzi G, Ambrosini CE, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTHa) during endoscopic parathyroidectomy: A prospective randomized trial. *Surg Endosc.* 2008;22(2):398-400.
100. Mihai R, Barczynski M, Iacobone M, Sitges-Serra A. Surgical strategy for sporadic primary hyperparathyroidism an evidence-based approach to surgical strategy, patient selection, surgical access, and reoperations. *Langenbecks Arch Surg.* 2009;394(5):785-98.
101. Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol.* 2016;174(1):D1-8.
102. Mohebbati A, Shaha AR. Imaging techniques in parathyroid surgery for primary hyperparathyroidism. *Am J Otolaryngol.* 2012;33(4):457-68.
103. Morris LG, Myssiorek D. When is surgery indicated for asymptomatic primary hyperparathyroidism? *Laryngoscope.* 2009;119(12):2291-2.
104. Mourad M, Buemi A, Darius T, Maiter D. Surgical options for primary hyperparathyroidism. *Ann Endocrinol (Paris).* 2015;76(5):638-42.
105. Nagar S, Reid D, Czako P, Long G, Shanley C. Outcomes analysis of intraoperative adjuncts during minimally invasive parathyroidectomy for primary hyperparathyroidism. *Am J Surg.* 2012;203(2):177-81.
106. Najafian A, Kahan S, Olson MT, Tufano RP, Zeiger MA. Intraoperative PTH May Not Be Necessary in the Management of Primary Hyperparathyroidism Even with Only One Positive or Only Indeterminate Preoperative Localization Studies. *World J Surg.* 2017;41(6):1500-5.
107. Nilsson IL. Primary hyperparathyroidism: should surgery be performed on all patients? Current evidence and residual uncertainties. *J Intern Med.* 2019;285(2):149-64.
108. Norlén O, Wang KC, Tay YK, Johnson WR, Grodski S, Yeung M, et al. No need to abandon focused parathyroidectomy: a multicenter study of long-term outcome after surgery for primary hyperparathyroidism. *Ann Surg.* 2015;261(5):991-6.
109. Norman J, Lopez J, Politz D. Abandoning unilateral parathyroidectomy: why we reversed our position after 15,000 parathyroid operations. *J Am Coll Surg.* 2012;214(3):260-9.
110. Olatoke SA, Agodirin OS, Rahman GA, Habeeb OG, Jimoh RO, Ahmed BA, et al. Serial pathologic fractures of five long bones on four separate occasions in a patient with primary hyperparathyroidism, challenges of management in a developing country: a case report. *Pan Afr Med J.* 2013;15:45.
111. Pallan S, Rahman MO, Khan AA. Diagnosis and management of primary hyperparathyroidism. *Bmj.* 2012;344:e1013.
112. Papier A, Kenig J, Barczynski M. [Evaluation of different intraoperative iPTH assay criteria in monitoring of minimally invasive parathyroidectomy for primary hyperparathyroidism]. *Przegl Lek.* 2014;71(1):14-8.
113. Paravastu SC, Chadwick DR. Parathyroidectomy in a district general hospital: outcomes and evolution in the era of minimally invasive surgery. *Int J Surg.* 2012;10(7):373-7.
114. Pasieka JL, Parsons L, Jones J. The long-term benefit of parathyroidectomy in primary hyperparathyroidism: a 10-year prospective surgical outcome study. *Surgery.* 2009;146(6):1006-13.
115. Paspala A, Spartalis E, Nastos C, Tsourouflis G, Dimitroulis D, Pikoulis E, et al. Robotic-assisted parathyroidectomy and short-term outcomes: a systematic review of the literature. *J Robot Surg.* 2020;14(6):821-7.
116. Patel KN, Caso R. Intraoperative Parathyroid Hormone Monitoring: Optimal Utilization. *Surg Oncol Clin N Am.* 2016;25(1):91-101.
117. Patel N, Mihai R. Long-term Cure of Primary Hyperparathyroidism After Scan-Directed Parathyroidectomy: Outcomes From A UK Endocrine Surgery Unit. *World J Surg.* 2022;46(9):2189-94.
118. Pelizzo MR, Pagetta C, Piotto A, Sorgato N, Merante Boschini I, Toniato A, et al. Surgical treatment of primary hyperparathyroidism: from bilateral neck exploration to minimally invasive surgery. *Minerva Endocrinol.* 2008;33(2):85-93.
119. Perrier N, Lang BH, Farias LCB, Poch LL, Sywak M, Almuqit M, et al. Surgical Aspects of Primary Hyperparathyroidism. *J Bone Miner Res.* 2022;37(11):2373-90.
120. Philippon M, Guerin C, Taieb D, Vaillant J, Morange I, Brue T, et al. Bilateral neck exploration in patients with primary hyperparathyroidism and discordant imaging results: a single-centre study. *Eur J Endocrinol.* 2014;170(5):719-25.
121. Prasannan S, Davies G, Bochner M, Kollias J, Malycha P. Minimally invasive parathyroidectomy using surgeon-performed ultrasound and sestamibi. *ANZ J Surg.* 2007;77(9):774-7.
122. Quinn AJ, Ryan É J, Garry S, James DL, Boland MR, Young O, et al. Use of Intraoperative Parathyroid Hormone in Minimally Invasive Parathyroidectomy for Primary Hyperparathyroidism: A Systematic Review and Meta-analysis. *JAMA Otolaryngol Head Neck Surg.* 2021;147(2):135-43.
123. Rajan S, Ravindhran B, George B, Bantwal G, Ayyar V, Manjuaanth S. Simplified intraoperative parathormone assay for primary hyperparathyroidism in a resource-limited setting. *Biomark Med.* 2021;15(5):331-6.
124. Richards ML, Thompson GB, Farley DR, Grant CS. An optimal algorithm for intraoperative parathyroid hormone monitoring. *Arch Surg.* 2011;146(3):280-5.
125. Rosenthal AA, Solomon RJ, Capasso T, Eyerly-Webb SA. Sestamibi Scanning and Intraoperative Parathyroid Hormone Results for Parathyroid Resection in Primary Hyperparathyroidism. *Am Surg.* 2018;84(8):e325-e7.
126. Rubello D, Kapse N, Grassetto G, Massaro A, Al-Nahhas A. Minimally invasive radio-guided surgery for primary hyperparathyroidism: From preoperative to intraoperative localization imaging. *Ann Endocrinol (Paris).* 2010;71(6):511-8.
127. Sadeghi N, Akin E, Lee JY, Roland J, Knoll S. Targeted parathyroidectomy: effectiveness and intraoperative rapid-parathormone dynamics. *Laryngoscope.* 2008;118(11):1997-2002.
128. Sadik KW, Kell M, Gorey T. Minimally invasive parathyroidectomy using surgical sonography. *Int J Med Sci.* 2011;8(4):283-6.
129. Salhi H, Bouziane T, Maaroufi M, Alaoui NI, El Ouahabi H. Primary hyperparathyroidism: Correlation between cervical ultrasound and MIBI scan. *Ann Afr Med.* 2022;21(2):161-4.
130. Seeliger B, Alesina PF, Koch JA, Hinrichs J, Meier B, Walz MK. Diagnostic value and clinical impact of complementary CT scan prior to surgery for non-localized primary hyperparathyroidism. *Langenbecks Arch Surg.* 2015;400(3):307-12.

- 131 Seib CD, Meng T, Suh I, Cisco RM, Lin DT, Morris AM, et al. Undertreatment of primary hyperparathyroidism in a privately insured US population: Decreasing utilization of parathyroidectomy despite expanding surgical guidelines. *Surgery*. 2021;169(1):87-93.
- 132 Shindo ML, Rosenthal JM. Minimal access parathyroidectomy using the focused lateral approach: technique, indication, and results. *Arch Otolaryngol Head Neck Surg*. 2007;133(12):1227-34.
- 133 Singer MC, Pucar D, Mathew M, Terris DJ. Improved localization of sestamibi imaging at high-volume centers. *Laryngoscope*. 2013;123(1):298-301.
- 134 Smith N, Magnuson JS, Vidrine DM, Kulbersh B, Peters GE. Minimally invasive parathyroidectomy: use of intraoperative parathyroid hormone assays after 2 preoperative localization studies. *Arch Otolaryngol Head Neck Surg*. 2009;135(11):1108-11.
- 135 Smith RB. Minimally invasive radioguided parathyroidectomy performed for primary hyperparathyroidism. *Otolaryngol Head Neck Surg*. 2010;142(3):462; author reply -3.
- 136 Solorzano CC, Carneiro-Pla D. Minimizing cost and maximizing success in the preoperative localization strategy for primary hyperparathyroidism. *Surg Clin North Am*. 2014;94(3):587-605.
- 137 Soon PS, Yeh MW, Sywak MS, Roach P, Delbridge LW, Sidhu SB. Minimally invasive parathyroidectomy using the lateral focused miniincision approach: Is there a learning curve for surgeons experienced in the open procedure? *J Am Coll Surg*. 2007;204(1):91-5.
- 138 Sosa JA. How Best to Approach Surgery for Primary Hyperparathyroidism-Can We All Agree? *JAMA Surg*. 2016;151(10):969.
- 139 Sugino K, Ito K, Nagahama M, Kitagawa W, Shibuya H, Ohkuwa K, et al. Minimally invasive surgery for primary hyperparathyroidism with or without intraoperative parathyroid hormone monitoring. *Endocr J*. 2010;57(11):953-8.
- 140 Suliburk JW, Perrier ND. Primary hyperparathyroidism. *Oncologist*. 2007;12(6):644-53.
- 141 Taieb A, Seman M, Menegaux F, Trésallet C. Surgical technique parathyroidectomy through a minimally invasive gland-centered localized approach for primary hyperparathyroidism. *J Visc Surg*. 2013;150(6):403-6.
- 142 Tunca F, Akici M, Işcan Y, Cem Sormaz I, Giles Senyurek Y, Terzioğlu T. The impact of combined interpretation of localization studies on image-guided surgical approaches for primary hyperparathyroidism. *Minerva Endocrinol*. 2017;42(3):213-22.
- 143 Udelsman R, Lin Z, Donovan P. The superiority of minimally invasive parathyroidectomy based on 1650 consecutive patients with primary hyperparathyroidism. *Ann Surg*. 2011;253(3):585-91.
- 144 Udelsman R, Pasieka JL, Sturgeon C, Young JE, Clark OH. Surgery for asymptomatic primary hyperparathyroidism: proceedings of the third international workshop. *J Clin Endocrinol Metab*. 2009;94(2):366-72.
- 145 Ullmann TM, Adam MA, Sosa JA. Surgeon Volume and Outcomes in Primary Hyperparathyroidism-What Is Old Is New Again. *JAMA Surg*. 2022;157(7):589.
- 146 Vaid S, Pandelidis S. Minimally invasive parathyroidectomy: a community hospital experience. *Arch Surg*. 2011;146(7):876-8.
- 147 Van Den Heede K, Bonheure A, Brusselsaers N, Van Slycke S. Long-term outcome of surgical techniques for sporadic primary hyperparathyroidism in a tertiary referral center in Belgium. *Langenbecks Arch Surg*. 2022;407(7):3045-55.
- 148 van der Plas WY, Noltes ME, Schaeffers A, Brouwers AH, van der Horst-Schrivers ANA, Kruijff S. [Diagnostic approach and treatment of primary hyperparathyroidism]. *Ned Tijdschr Geneesk*. 2017;161:D1870.
- 149 Van Udelsman B, Udelsman R. Surgery in primary hyperparathyroidism: extensive personal experience. *J Clin Densitom*. 2013;16(1):54-9.
- 150 Vetshev PS, Drozhzhin AY, Zhivotov VA, Yankin PL, Poddubny EI, Krastyn EA. [Current approach to the diagnosis and treatment of primary hyperparathyroidism]. *Khirurgiia (Mosk)*. 2019(6):26-34.
- 151 Walker MD, Bilezikian JP. Primary hyperparathyroidism: recent advances. *Curr Opin Rheumatol*. 2018;30(4):427-39.
- 152 Williams BA, Trites JR, Taylor SM, Bullock MJ, Hart RD. Surgical management of primary hyperparathyroidism in Canada. *J Otolaryngol Head Neck Surg*. 2014;43(1):44.
- 153 Wu S, Hwang SS, Haigh PI. Influence of a negative sestamibi scan on the decision for parathyroid operation by the endocrinologist and surgeon. *Surgery*. 2017;161(1):35-43.
- 154 Yeh MW, Wiseman JE, Ituarte PH, Pasternak JD, Hwang RS, Wu B, et al. Surgery for primary hyperparathyroidism: are the consensus guidelines being followed? *Ann Surg*. 2012;255(6):1179-83.
- 155 Yew MK, Thompson IJ. Minimally invasive parathyroidectomy: an audit of a change in clinical practice. *ANZ J Surg*. 2007;77(1-2):24-6.
- 156 Zammit M, Pierce K, Bailey L, Rowland M, Waghorn A, Shore S. Challenging NICE guidelines on parathyroid surgery. *Surgeon*. 2022;20(4):e105-e11.
- 157 Zanocco K, Heller M, Sturgeon C. Cost-effectiveness of parathyroidectomy for primary hyperparathyroidism. *Endocr Pract*. 2011;17 Suppl 1:69-74.
- 158 Zanocco K, Sturgeon C. How should age at diagnosis impact treatment strategy in asymptomatic primary hyperparathyroidism? A cost-effectiveness analysis. *Surgery*. 2008;144(2):290-8.
- 159 Zanocco KA, Wu JX, Yeh MW. Parathyroidectomy for asymptomatic primary hyperparathyroidism: A revised cost-effectiveness analysis incorporating fracture risk reduction. *Surgery*. 2017;161(1):16-24.
- 160 Zhu CY, Nguyen DT, Yeh MW. Who Benefits from Treatment of Primary Hyperparathyroidism? *Surg Clin North Am*. 2019;99(4):667-79.
- 161 Aarum S, Nordenström J, Reinhér E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for primary hyperparathyroidism: the new versus the old order. A randomised controlled trial of preoperative localisation. *Scand J Surg*. 2007;96(1):26-30.
- 162 Abbott DE, Cantor SB, Grubbs EG, Santora R, Gomez HF, Evans DB, et al. Outcomes and economic analysis of routine preoperative 4-dimensional CT for surgical intervention in de novo primary hyperparathyroidism: does clinical benefit justify the cost? *J Am Coll Surg*. 2012;214(4):629-37; discussion 37-9.
- 163 Adler JT, Sippel RS, Chen H. New trends in parathyroid surgery. *Curr Probl Surg*. 2010;47(12):958-1017.
- 164 Al-Qurayshi Z, Hauch A, Srivastav S, Kandil E. Ethnic and economic disparities effect on management of hyperparathyroidism. *Am J Surg*. 2017;213(6):1134-42.
- 165 Aliabadi-Wahle S, Kelly TL, Rozenfeld Y, Carlisle JR, Naeole LK, Negreanu FA, et al. Treatment strategies for primary hyperparathyroidism: what is the cost? *Am Surg*. 2014;80(11):1146-51.
- 166 Alobuia WM, Meng T, Cisco RM, Lin DT, Suh I, Tamura MK, et al. Racial disparities in the utilization of parathyroidectomy among patients with primary hyperparathyroidism: Evidence from a nationwide analysis of Medicare claims. *Surgery*. 2022;171(1):8-16.
- 167 Badii B, Staderini F, Foppa C, Tofani L, Skalamera I, Fiorenza G, et al. Cost-benefit analysis of the intraoperative parathyroid hormone assay in primary hyperparathyroidism. *Head Neck*. 2017;39(2):241-6.
- 168 Baliski C, Nosyk B, Melck A, Bugis S, Rosenberg F, A HA. The cost-effectiveness of three strategies for the surgical treatment of symptomatic primary hyperparathyroidism. *Ann Surg Oncol*. 2008;15(10):2653-60.
- 169 Barczynski M. Minimally invasive parathyroidectomy without intraoperative parathyroid hormone monitoring: when and why? *J Postgrad Med*. 2009;55(4):239-40.
- 170 Barczyński M, Gólkowski F, Nawrot I. The current status of intraoperative iPTH assay in surgery for primary hyperparathyroidism. *Gland Surg*. 2015;4(1):36-43.
- 171 Belozeroff V, Cooper K, Hess G, Chang CL. Healthcare use and costs before and after parathyroidectomy in patients on dialysis. *BMC Health Serv Res*. 2013;13:248.
- 172 Bunch PM, Kelly HR. Preoperative Imaging Techniques in Primary Hyperparathyroidism: A Review. *JAMA Otolaryngol Head Neck Surg*. 2018;144(10):929-37.
- 173 Burke E, Waris A, O'Donoghue G. Surgical Parathyroidectomy Services. *Ir Med J*. 2019;112(4):916.
- 174 Chen H. Parathyroid Disease: Often Forgotten and Undertreated. *Am Surg*. 2020;86(7):730-3.
- 175 Cohen O, Tufano RP, Anuwong A, Shaha AR, Olsen KD, Zafereo M, et al. Transoral endoscopic vestibular approach for thyroidectomy and parathyroidectomy - From promise to practice. *Am J Otolaryngol*. 2021;42(5):103022.
- 176 Contreras K, Baquero R, Buitrago G. Clinical and Economical Outcomes Associated with Parathyroidectomy: A 5-Year Population-Based Study in a Middle-Income Country with Universal Health Coverage. *Int J Nephrol*. 2020;2020:7250250.
- 177 Danese MD, Fox KM, Duryea JL, Desai P, Rubin RJ. The rate, cost and outcomes of parathyroidectomy in the united states dialysis population from 2016-2018. *BMC Nephrol*. 2022;23(1):220.
- 178 Dimas S, Michas S, Christakis I, Augoustis C, Alevizaki M. Minimally invasive parathyroidectomy in patients with previous neck surgery. *Hormones (Athens)*. 2012;11(2):160-5.
- 179 Dombrowsky A, Weiss D, Bushman N, Chen H, Balentine CJ. Can imaging studies be omitted in patients with sporadic primary hyperparathyroidism? *J Surg Res*. 2018;231:257-62.
- 180 Dream S, Chen H, Lindeman B. Tertiary Hyperparathyroidism: Why the Delay? *Ann Surg*. 2021;273(3):e120-e2.
- 181 Dyas AR, Lovell KM, Balentine CJ, Wang TN, Porterfield JR, Jr., Chen H, et al. Reducing cost and improving operating room efficiency: examination of surgical instrument processing. *J Surg Res*. 2018;229:15-9.
- 182 Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *Surg Clin North Am*. 2014;94(3):607-23.
- 183 Frank E, Watson W, Fujimoto S, De Andrade Filho P, Inman J, Simental A. Surgery versus Imaging in Non-Localizing Primary Hyperparathyroidism: A Cost-Effectiveness Model. *Laryngoscope*. 2020;130(12):E963-e9.
- 184 Gracie D, Hussain SS. Use of minimally invasive parathyroidectomy techniques in sporadic primary hyperparathyroidism: systematic review. *J Laryngol Otol*. 2012;126(3):221-7.
- 185 Hassan-Smith ZK, Criseno S, Gittoes NJL. Mild primary hyperparathyroidism-to treat or not to treat? *Br Med Bull*. 2019;129(1):53-67.
- 186 Hollenbeak CS, Lendel I, Beus KS, Ruda JM, Stack BC, Jr. The cost of screening for synchronous thyroid disease in patients presenting with primary hyperparathyroidism. *Arch Otolaryngol Head Neck Surg*. 2007;133(10):1013-21.

- 187 Ikeda Y, Takayama J, Takami H. Minimally invasive radioguided parathyroidectomy for hyperparathyroidism. *Ann Nucl Med*. 2010;24(4):233-40.
- 188 Jang S, Mandabach M, Aburjania Z, Balentine CJ, Chen H. Racial disparities in the cost of surgical care for parathyroidectomy. *J Surg Res*. 2018;221:216-21.
- 189 Jang S, Mandabach M, Aburjania Z, Balentine CJ, Chen H. Wide variation in cost of surgical care for parathyroidectomy: is there a need for standardization of practice? *Surgery*. 2018;163(3):638-42.
- 190 Karahan Ö, Okus A, Sevinç B, Eryılmaz MA, Ay S, Çaycı M, et al. Minimally invasive parathyroidectomy under local anesthesia. *J Postgrad Med*. 2013;59(1):21-4.
- 191 Kim SM, Long J, Montez-Rath ME, Leonard MB, Norton JA, Chertow GM. Rates and Outcomes of Parathyroidectomy for Secondary Hyperparathyroidism in the United States. *Clin J Am Soc Nephrol*. 2016;11(7):1260-7.
- 192 Kim SM, Shu AD, Long J, Montez-Rath ME, Leonard MB, Norton JA, et al. Declining Rates of Inpatient Parathyroidectomy for Primary Hyperparathyroidism in the US. *PLoS One*. 2016;11(8):e0161192.
- 193 Kunstman JW, Kirsch JD, Mahajan A, Udelsman R. Clinical review: Parathyroid localization and implications for clinical management. *J Clin Endocrinol Metab*. 2013;98(3):902-12.
- 194 Kunstman JW, Udelsman R. Superiority of minimally invasive parathyroidectomy. *Adv Surg*. 2012;46:171-89.
- 195 Lanitis S, Chortis P, Sourtse G, Gkanis V, Lainas S, Tournis S, et al. Shifting from open to video-assisted parathyroidectomy: effect of the adjustment period on safety, clinical outcomes and cost. *Ann R Coll Surg Engl*. 2022;104(4):295-301.
- 196 Lau WL, Obi Y, Kalantar-Zadeh K. Parathyroidectomy in the Management of Secondary Hyperparathyroidism. *Clin J Am Soc Nephrol*. 2018;13(6):952-61.
- 197 Lee L, Steward DL. Techniques for parathyroid localization with ultrasound. *Otolaryngol Clin North Am*. 2010;43(6):1229-39, vi.
- 198 Lorenz K, Dralle H. [Intraoperative parathyroid hormone determination for primary hyperparathyroidism]. *Chirurg*. 2010;81(7):636, 8-42.
- 199 McManus C, Oh A, Lee JA, Hur C, Kuo JH. Timing of parathyroidectomy for tertiary hyperparathyroidism with end-stage renal disease: A cost-effectiveness analysis. *Surgery*. 2021;169(1):94-101.
- 200 Melfa GI, Raspani C, Attard M, Cocorullo G, Attard A, Mazzola S, et al. Comparison of minimally invasive parathyroidectomy under local anaesthesia and minimally invasive video-assisted parathyroidectomy for primary hyperparathyroidism: a cost analysis. *G Chir*. 2016;37(2):61-7.
- 201 Miccoli P, Berti P, Materazzi G, Ambrosini CE, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTHa) during endoscopic parathyroidectomy: A prospective randomized trial. *Surg Endosc*. 2008;22(2):398-400.
- 202 Mihai R, Weisters M, Stechman MJ, Gleeson F, Sadler G. Cost-effectiveness of scan-directed parathyroidectomy. *Langenbecks Arch Surg*. 2008;393(5):739-43.
- 203 Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol*. 2016;174(1):D1-8.
- 204 Mishra A, Kapoor L, Mishra SK. Post-operative care through tele-follow up visits in patients undergoing thyroidectomy and parathyroidectomy in a resource-constrained environment. *J Telemed Telecare*. 2009;15(2):73-6.
- 205 Morris LF, Zanocco K, Ituarte PH, Ro K, Duh QY, Sturgeon C, et al. The value of intraoperative parathyroid hormone monitoring in localized primary hyperparathyroidism: a cost analysis. *Ann Surg Oncol*. 2010;17(3):679-85.
- 206 National Guideline C. NICE Evidence Reviews Collection. Evidence review for surgical localisation: Hyperparathyroidism (primary): diagnosis, assessment and initial management: Evidence review D. London: National Institute for Health and Care Excellence (NICE) Copyright © NICE 2019.; 2019.
- 207 Neves MCD, Abrahão AR, Abrahão M, Rosano M, Rocha LAD, Machado H, et al. Sestamibi scan in renal parathyroidectomy: a worthwhile preoperative exam? *Braz J Otorhinolaryngol*. 2022;88(5):740-4.
- 208 Niederle B, Wémeau JL. Is surgery necessary for 'mild' or 'asymptomatic' hyperparathyroidism? *Eur J Endocrinol*. 2015;173(3):D13-20.
- 209 Nilsson IL. Primary hyperparathyroidism: should surgery be performed on all patients? Current evidence and residual uncertainties. *J Intern Med*. 2019;285(2):149-64.
- 210 O'Connell DA, Seikaly H, Harris JR. Central laboratory versus point of care testing in intraoperative monitoring of parathyroid hormone levels: cost comparison. *J Otolaryngol Head Neck Surg*. 2008;37(11):91-7.
- 211 Paker M, Fisher S, Mazzawi S, Colodner R, Ashkenazi D. Intraoperative Parathyroid Aspiration and Parathyroid Hormone Assay During Parathyroidectomy for Primary Hyperparathyroidism. *Isr Med Assoc J*. 2017;19(12):731-5.
- 212 Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *Am J Otolaryngol*. 2021;42(1):102819.
- 213 Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian J Endocrinol Metab*. 2019;23(3):347-52.
- 214 Rroji M, Spasovski G. Calcimimetics versus parathyroidectomy: What is preferable? *Int Urol Nephrol*. 2018;50(7):1271-5.
- 215 Saravana-Bawan B, Auguste BL, Zahirieh A, Devon K. Ambulatory Parathyroidectomy for Secondary Hyperparathyroidism at a Large Dialysis Program in Toronto: A Program Report. *Can J Kidney Health Dis*. 2022;9:20543581221127937.
- 216 Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally Invasive Parathyroidectomy without Intraoperative PTH Performed after Positive Ultrasonography as the only Diagnostic Method in Patients with Primary Hyperparathyroidism. *World J Surg*. 2019;43(6):1525-31.
- 217 Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia - A Feasibility Study. *Indian J Endocrinol Metab*. 2019;23(1):67-71.
- 218 Setabutr D, Vakharia K, Nogan SJ, Kamel GN, Allen T, Saunders BD, et al. Comparison of SPECT/CT and planar MIBI in terms of operating time and cost in the surgical management of primary hyperparathyroidism. *Ear Nose Throat J*. 2015;94(10-11):448-52.
- 219 Snopok I, Viebahn R, Walz M, Zgoura P, Alesina PF. Minimally invasive video-assisted parathyroidectomy (MIVAP) versus conventional parathyroidectomy for renal hyperparathyroidism: a retrospective multicenter study. *Updates Surg*. 2022;74(4):1419-28.
- 220 Solorzano CC, Carneiro-Pla D. Minimizing cost and maximizing success in the preoperative localization strategy for primary hyperparathyroidism. *Surg Clin North Am*. 2014;94(3):587-605.
- 221 Spiros D, Nikolaos R, Ioannis C. Minimally invasive parathyroidectomy in patients with previous endocrine surgery. *Jsis*. 2011;15(4):499-503.
- 222 Stalberg P, Delbridge L, van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy--current concepts. *Surgeon*. 2007;5(5):301-8.
- 223 To H, Otto G. Selective intra-operative parathyroid hormone in re-do neck exploration in parathyroidectomy: A case report. *Int J Surg Case Rep*. 2019;57:1-4.
- 224 Tolley N, Arora A, Palazzo F, Garas G, Dhawan R, Cox J, et al. Robotic-assisted parathyroidectomy: a feasibility study. *Otolaryngol Head Neck Surg*. 2011;144(6):859-66.
- 225 Tublin ME, Pryma DA, Yim JH, Ogilvie JB, Mountz JM, Bencherif B, et al. Localization of parathyroid adenomas by sonography and technetium tc 99m sestamibi single-photon emission computed tomography before minimally invasive parathyroidectomy: are both studies really needed? *J Ultrasound Med*. 2009;28(2):183-90.
- 226 Udelsman R, Lin Z, Donovan P. The superiority of minimally invasive parathyroidectomy based on 1650 consecutive patients with primary hyperparathyroidism. *Ann Surg*. 2011;253(3):585-91.
- 227 Wang TS, Cheung K, Farrokhyar F, Roman SA, Sosa JA. Would scan, but which scan? A cost-utility analysis to optimize preoperative imaging for primary hyperparathyroidism. *Surgery*. 2011;150(6):1286-94.
- 228 Wilhelm SM, Wang TS, Ruan DT, Lee JA, Asa SL, Duh QY, et al. The American Association of Endocrine Surgeons Guidelines for Definitive Management of Primary Hyperparathyroidism. *JAMA Surg*. 2016;151(10):959-68.
- 229 Xie S, Kuriakose J, Beninato T, Carayannopoulos M, Trooskin SZ, Libutti SK, et al. Association of Implementation of Operating Room-Based Parathyroid Hormone Testing with Operative Time and Cost. *J Am Coll Surg*. 2022;235(6):906-12.
- 230 Yap A, Hope TA, Graves CE, Kluijfhout W, Shen WT, Gosnell JE, et al. A cost-utility analysis of 18F-fluorocholine-positron emission tomography imaging for localizing primary hyperparathyroidism in the United States. *Surgery*. 2022;171(1):55-62.
- 231 Yuan CM, Nee R, Narayan R, Abbott KC. Treatment of secondary hyperparathyroidism with parathyroidectomy instead of cinacalcet: time to pick the low-hanging fruit? *Am J Kidney Dis*. 2012;60(2):179-81.
- 232 Zammit M, Pierce K, Bailey L, Rowland M, Waghorn A, Shore S. Challenging NICE guidelines on parathyroid surgery. *Surgeon*. 2022;20(4):e105-e11.
- 233 Zanocco K, Heller M, Sturgeon C. Cost-effectiveness of parathyroidectomy for primary hyperparathyroidism. *Endocr Pract*. 2011;17 Suppl 1:69-74.
- 234 Zanocco K, Sturgeon C. How should age at diagnosis impact treatment strategy in asymptomatic primary hyperparathyroidism? A cost-effectiveness analysis. *Surgery*. 2008;144(2):290-8.
- 235 Zanocco KA, Wu JX, Yeh MW. Parathyroidectomy for asymptomatic primary hyperparathyroidism: A revised cost-effectiveness analysis incorporating fracture risk reduction. *Surgery*. 2017;161(1):16-24.
- 236 Zhu CY, Nguyen DT, Yeh MW. Who Benefits from Treatment of Primary Hyperparathyroidism? *Surg Clin North Am*. 2019;99(4):667-79.
- 237 Aarum S, Nordenström J, Reinhér E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for primary hyperparathyroidism: the new versus the old order. A randomised controlled trial of preoperative localisation. *Scand J Surg*. 2007;96(1):26-30.
- 238 Abbott DE, Cantor SB, Grubbs EG, Santora R, Gomez HF, Evans DB, et al. Outcomes and economic analysis of routine preoperative 4-dimensional CT for surgical intervention in de novo primary hyperparathyroidism: does clinical benefit justify the cost? *J Am Coll Surg*. 2012;214(4):629-37; discussion 37-9.
- 239 Agha A, Hornung M, Rennert J, Uller W, Lighvani H, Schlitt HJ, et al. Contrast-enhanced ultrasonography for localization of pathologic glands in patients with primary hyperparathyroidism. *Surgery*. 2012;151(4):580-6.
- 240 Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: Feasibility and outcome. *J Visc Surg*. 2018;155(4):253-8.

- 241 Bunch PM, Kelly HR. Preoperative Imaging Techniques in Primary Hyperparathyroidism: A Review. *JAMA Otolaryngol Head Neck Surg.* 2018;144(10):929-37.
- 242 Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *Surg Clin North Am.* 2014;94(3):607-23.
- 243 Frank E, Watson W, Fujimoto S, De Andrade Filho P, Inman J, Simental A. Surgery versus Imaging in Non-Localizing Primary Hyperparathyroidism: A Cost-Effectiveness Model. *Laryngoscope.* 2020;130(12):E963-e9.
- 244 Gracie D, Hussain SS. Use of minimally invasive parathyroidectomy techniques in sporadic primary hyperparathyroidism: systematic review. *J Laryngol Otol.* 2012;126(3):221-7.
- 245 Hassan-Smith ZK, Criseno S, Gittoes NJL. Mild primary hyperparathyroidism-to treat or not to treat? *Br Med Bull.* 2019;129(1):53-67.
- 246 Karahan Ö, Okus A, Sevinç B, Eryılmaz MA, Ay S, Çaycı M, et al. Minimally invasive parathyroidectomy under local anesthesia. *J Postgrad Med.* 2013;59(1):21-4.
- 247 Kunstman JW, Kirsch JD, Mahajan A, Udelsman R. Clinical review: Parathyroid localization and implications for clinical management. *J Clin Endocrinol Metab.* 2013;98(3):902-12.
- 248 Kunstman JW, Udelsman R. Superiority of minimally invasive parathyroidectomy. *Adv Surg.* 2012;46:171-89.
- 249 Lanitis S, Chortis P, Sourtse G, Gkanis V, Lainas S, Tournis S, et al. Shifting from open to video-assisted parathyroidectomy: effect of the adjustment period on safety, clinical outcomes and cost. *Ann R Coll Surg Engl.* 2022;104(4):295-301.
- 250 Lee GS, McKenzie TJ, Mullan BP, Farley DR, Thompson GB, Richards ML. A Multimodal Imaging Protocol, (123I)/(99)Tc-Sestamibi, SPECT, and SPECT/CT, in Primary Hyperparathyroidism Adds Limited Benefit for Preoperative Localization. *World J Surg.* 2016;40(3):589-94.
- 251 Lee L, Steward DL. Techniques for parathyroid localization with ultrasound. *Otolaryngol Clin North Am.* 2010;43(6):1229-39, vi.
- 252 Lubitz CC, Stephen AE, Hodin RA, Pandharipande P. Preoperative localization strategies for primary hyperparathyroidism: an economic analysis. *Ann Surg Oncol.* 2012;19(13):4202-9.
- 253 Madorin CA, Owen R, Coakley B, Lowe H, Nam KH, Weber K, et al. Comparison of radiation exposure and cost between dynamic computed tomography and sestamibi scintigraphy for preoperative localization of parathyroid lesions. *JAMA Surg.* 2013;148(6):500-3.
- 254 Melfa GI, Raspanti C, Attard M, Cocorullo G, Attard A, Mazzola S, et al. Comparison of minimally invasive parathyroidectomy under local anaesthesia and minimally invasive video-assisted parathyroidectomy for primary hyperparathyroidism: a cost analysis. *G Chir.* 2016;37(2):61-7.
- 255 Miccoli P, Berti P, Materazzi G, Ambrosini CE, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTHa) during endoscopic parathyroidectomy: A prospective randomized trial. *Surg Endosc.* 2008;22(2):398-400.
- 256 Mihai R, Weisters M, Stechman MJ, Gleeson F, Sadler G. Cost-effectiveness of scan-directed parathyroidectomy. *Langenbecks Arch Surg.* 2008;393(5):739-43.
- 257 Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol.* 2016;174(1):D1-8.
- 258 Morris LF, Zanocco K, Ituarte PH, Ro K, Duh QY, Sturgeon C, et al. The value of intraoperative parathyroid hormone monitoring in localized primary hyperparathyroidism: a cost analysis. *Ann Surg Oncol.* 2010;17(3):679-85.
- 259 National Guideline C. NICE Evidence Reviews Collection. Evidence review for surgical localisation: Hyperparathyroidism (primary): diagnosis, assessment and initial management: Evidence review D. London: National Institute for Health and Care Excellence (NICE) Copyright © NICE 2019.; 2019.
- 260 Pata G, Casella C, Magri GC, Lucchini S, Panarotto MB, Crea N, et al. Financial and clinical implications of low-energy CT combined with 99m Technetium-sestamibi SPECT for primary hyperparathyroidism. *Ann Surg Oncol.* 2011;18(9):2555-63.
- 261 Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *Am J Otolaryngol.* 2021;42(1):102819.
- 262 Powell DK, Nwofe F, Goldfarb RC, Ongseng F. Tc-99m sestamibi parathyroid gland scintigraphy: added value of Tc-99m pertechnetate thyroid imaging for increasing interpretation confidence and avoiding additional testing. *Clin Imaging.* 2013;37(3):475-9.
- 263 Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally Invasive Parathyroidectomy without Intraoperative PTH Performed after Positive Ultrasonography as the only Diagnostic Method in Patients with Primary Hyperparathyroidism. *World J Surg.* 2019;43(6):1525-31.
- 264 Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia - A Feasibility Study. *Indian J Endocrinol Metab.* 2019;23(1):67-71.
- 265 Setabutr D, Vakharia K, Nogan SJ, Kamel GN, Allen T, Saunders BD, et al. Comparison of SPECT/CT and planar MIBI in terms of operating time and cost in the surgical management of primary hyperparathyroidism. *Ear Nose Throat J.* 2015;94(10-11):448-52.
- 266 Seving AI, Derici ZS, Bekiş R, Canda T, Saydam MA, Harmancıoğlu O. Success of minimally invasive single-gland exploration using the quick intraoperative parathyroid assay. *Acta Chir Belg.* 2010;110(4):463-6.
- 267 Solorzano CC, Carneiro-Pla D. Minimizing cost and maximizing success in the preoperative localization strategy for primary hyperparathyroidism. *Surg Clin North Am.* 2014;94(3):587-605.
- 268 Stalberg P, Delbridge L, van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy--current concepts. *Surgeon.* 2007;5(5):301-8.
- 269 Tublin ME, Pryma DA, Yim JH, Ogilvie JB, Mountz JM, Bencherif B, et al. Localization of parathyroid adenomas by sonography and technetium tc 99m sestamibi single-photon emission computed tomography before minimally invasive parathyroidectomy: are both studies really needed? *J Ultrasound Med.* 2009;28(2):183-90.
- 270 Vitetta GM, Ravera A, Mensa G, Fuso L, Neri P, Carriero A, et al. Actual role of color-doppler high-resolution neck ultrasonography in primary hyperparathyroidism: a clinical review and an observational study with a comparison of (99m)Tc-sestamibi parathyroid scintigraphy. *J Ultrasound.* 2019;22(3):291-308.
- 271 von Breitenbuch P, Iesalnieks I, Piso P, Schlitt HJ, Agha A. [Primary hyperparathyroidism: clinical symptoms, diagnostic significance and localisation--a retrospective analysis]. *Zentralbl Chir.* 2007;132(6):497-503.
- 272 Yap A, Hope TA, Graves CE, Kluijfhout W, Shen WT, Gosnell JE, et al. A cost-utility analysis of 18F-fluorocholine-positron emission tomography imaging for localizing primary hyperparathyroidism in the United States. *Surgery.* 2022;171(1):55-62.
- 273 Zafereo M, Yu J, Angelos P, Brumund K, Chuang HH, Goldenberg D, et al. American Head and Neck Society Endocrine Surgery Section update on parathyroid imaging for surgical candidates with primary hyperparathyroidism. *Head Neck.* 2019;41(7):2398-409.
- 274 Zanocco K, Heller M, Sturgeon C. Cost-effectiveness of parathyroidectomy for primary hyperparathyroidism. *Endocr Pract.* 2011;17 Suppl 1:69-74.
- 275 Aarum S, Nordenström J, Reihné E, Zedenius J, Jacobsson H, Danielsson R, et al. Operation for primary hyperparathyroidism: the new versus the old order. A randomised controlled trial of preoperative localisation. *Scand J Surg.* 2007;96(1):26-30.
- 276 Acar N, Hacıyanlı M, Coskun M, Erdogan NK, Celik SC, Hacıyanlı SG, et al. Diagnostic value of four-dimensional computed tomography and four-dimensional magnetic resonance imaging in primary hyperparathyroidism when first-line imaging was inadequate. *Ann R Coll Surg Engl.* 2020;102(4):294-9.
- 277 Barczyński M, Gólkowski F, Nawrot I. The current status of intraoperative iPTH assay in surgery for primary hyperparathyroidism. *Gland Surg.* 2015;4(1):36-43.
- 278 Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: Feasibility and outcome. *J Visc Surg.* 2018;155(4):253-8.
- 279 Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *Surg Clin North Am.* 2014;94(3):607-23.
- 280 Farley DR. Technetium-99m 2-methoxyisobutyl isonitrile-scintigraphy: preoperative and intraoperative guidance for primary hyperparathyroidism. *World J Surg.* 2004;28(12):1207-11.
- 281 Kunstman JW, Udelsman R. Superiority of minimally invasive parathyroidectomy. *Adv Surg.* 2012;46:171-89.
- 282 Lee GS, McKenzie TJ, Mullan BP, Farley DR, Thompson GB, Richards ML. A Multimodal Imaging Protocol, (123I)/(99)Tc-Sestamibi, SPECT, and SPECT/CT, in Primary Hyperparathyroidism Adds Limited Benefit for Preoperative Localization. *World J Surg.* 2016;40(3):589-94.
- 283 Miccoli P, Berti P, Materazzi G, Ambrosini CE, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTHa) during endoscopic parathyroidectomy: A prospective randomized trial. *Surg Endosc.* 2008;22(2):398-400.
- 284 Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol.* 2016;174(1):D1-8.
- 285 National Guideline C. NICE Evidence Reviews Collection. Evidence review for surgical localisation: Hyperparathyroidism (primary): diagnosis, assessment and initial management: Evidence review D. London: National Institute for Health and Care Excellence (NICE) Copyright © NICE 2019.; 2019.
- 286 Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *Am J Otolaryngol.* 2021;42(1):102819.
- 287 Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian J Endocrinol Metab.* 2019;23(3):347-52.
- 288 Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally Invasive Parathyroidectomy without Intraoperative PTH Performed after Positive Ultrasonography as the only Diagnostic Method in Patients with Primary Hyperparathyroidism. *World J Surg.* 2019;43(6):1525-31.
- 289 Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia - A Feasibility Study. *Indian J Endocrinol Metab.* 2019;23(1):67-71.
- 290 Stalberg P, Delbridge L, van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy--current concepts. *Surgeon.* 2007;5(5):301-8.
- 291 Tolley N, Garas G, Palazzo F, Prichard A, Chaidas K, Cox J, et al. Long-term prospective evaluation comparing robotic parathyroidectomy with minimally invasive open parathyroidectomy for primary hyperparathyroidism. *Head Neck.* 2016;38 Suppl 1:E300-6.
- 292 Udelsman R, Lin Z, Donovan P. The superiority of minimally invasive parathyroidectomy based on 1650 consecutive patients with primary hyperparathyroidism. *Ann Surg.* 2011;253(3):585-91.

293 Wilhelm SM, Wang TS, Ruan DT, Lee JA, Asa SL, Duh QY, et al. The American Association of Endocrine Surgeons Guidelines for Definitive Management of Primary Hyperparathyroidism. *JAMA Surg.* 2016;151(10):959-68.

294 Xie S, Kuriakose J, Beninato T, Carayannopoulos M, Trooskin SZ, Libutti SK, et al. Association of Implementation of Operating Room-Based Parathyroid Hormone Testing with Operative Time and Cost. *J Am Coll Surg.* 2022;235(6):906-12.

295 Yap A, Hope TA, Graves CE, Kluijfhout W, Shen WT, Gosnell JE, et al. A cost-utility analysis of 18F-fluorocholine-positron emission tomography imaging for localizing primary hyperparathyroidism in the United States. *Surgery.* 2022;171(1):55-62.

296 Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: Feasibility and outcome. *J Visc Surg.* 2018;155(4):253-8.

297 Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *Surg Clin North Am.* 2014;94(3):607-23.

298 Kunstman JW, Kirsch JD, Mahajan A, Udelsman R. Clinical review: Parathyroid localization and implications for clinical management. *J Clin Endocrinol Metab.* 2013;98(3):902-12.

299 Kunstman JW, Udelsman R. Superiority of minimally invasive parathyroidectomy. *Adv Surg.* 2012;46:171-89.

300 Lee L, Steward DL. Techniques for parathyroid localization with ultrasound. *Otolaryngol Clin North Am.* 2010;43(6):1229-39, vi.

301 Miccoli P, Berti P, Materazzi G, Ambrosini CE, Fregoli L, Donatini G. Endoscopic bilateral neck exploration versus quick intraoperative parathormone assay (qPTHa) during endoscopic parathyroidectomy: A prospective randomized trial. *Surg Endosc.* 2008;22(2):398-400.

302 Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol.* 2016;174(1):D1-8.

303 Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *Am J Otolaryngol.* 2021;42(1):102819.

304 Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian J Endocrinol Metab.* 2019;23(3):347-52.

305 Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia - A Feasibility Study. *Indian J Endocrinol Metab.* 2019;23(1):67-71.

306 Stalberg P, Delbridge L, van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy--current concepts. *Surgeon.* 2007;5(5):301-8.

307 Tolley N, Garas G, Palazzo F, Prichard A, Chaidas K, Cox J, et al. Long-term prospective evaluation comparing robotic parathyroidectomy with minimally invasive open parathyroidectomy for primary hyperparathyroidism. *Head Neck.* 2016;38 Suppl 1:E300-6.

308 Wilhelm SM, Wang TS, Ruan DT, Lee JA, Asa SL, Duh QY, et al. The American Association of Endocrine Surgeons Guidelines for Definitive Management of Primary Hyperparathyroidism. *JAMA Surg.* 2016;151(10):959-68.

309 Xie S, Kuriakose J, Beninato T, Carayannopoulos M, Trooskin SZ, Libutti SK, et al. Association of Implementation of Operating Room-Based Parathyroid Hormone Testing with Operative Time and Cost. *J Am Coll Surg.* 2022;235(6):906-12.

310 Aspinall SR, Boase S, Malycha P. Long-term symptom relief from primary hyperparathyroidism following minimally invasive parathyroidectomy. *World J Surg.* 2010;34(9):2223-7.

311 Henry JF, Sebag F, Cherenko M, Ippolito G, Taieb D, Vaillant J. Endoscopic parathyroidectomy: why and when? *World J Surg.* 2008;32(11):2509-15.

312 McLean T, Delbridge L. Comparison of consumer information on the internet to the current evidence base for minimally invasive parathyroidectomy. *World J Surg.* 2010;34(6):1304-11.

313 Miccoli P, Materazzi G, Baggiani A, Miccoli M. Mini-invasive video-assisted surgery of the thyroid and parathyroid glands: a 2011 update. *J Endocrinol Invest.* 2011;34(6):473-80.

314 Sharanappa V, Mishra A, Bhatia V, Mayilvagnan S, Chand G, Agarwal G, et al. Pediatric Primary Hyperparathyroidism: Experience in a Tertiary Care Referral Center in a Developing Country Over Three Decades. *World J Surg.* 2021;45(2):488-95.

315 Venkat R, Kouniavsky G, Tufano RP, Schneider EB, Dackiw AP, Zeiger MA. Long-term outcome in patients with primary hyperparathyroidism who underwent minimally invasive parathyroidectomy. *World J Surg.* 2012;36(1):55-60.

316 Wojtczak B, Strycka J, Kaliszewski K, Rudnicki J, Bolanowski M, Barczyński M. Surgical implications of recent modalities for parathyroid imaging. *Gland Surg.* 2020;9(Suppl 2):S86-s94.

317 Bombil I, Louw L, Mitchell C, Mahlobo F, Muganza RA, Madima NR. Sonar guided focused parathyroidectomy under cervical block. *S Afr J Surg.* 2018;56(2):30-3.

318 Mallikarjuna VJ, Mathew V, Ayyar V, Bantwal G, Ganesh V, George B, et al. Five-year Retrospective Study on Primary Hyperparathyroidism in South India: Emerging Roles of Minimally Invasive Parathyroidectomy and Preoperative Localization with Methionine Positron Emission Tomography-Computed Tomography Scan. *Indian J Endocrinol Metab.* 2018;22(3):355-61.

319 Nouikes Zitouni S. Monocentric experience of primary hyperparathyroidism surgery in Algeria. *Surg Open Sci.* 2021;4:32-6.

320 Sharanappa V, Mishra A, Bhatia V, Mayilvagnan S, Chand G, Agarwal G, et al. Pediatric Primary Hyperparathyroidism: Experience in a Tertiary Care Referral Center in a Developing Country Over Three Decades. *World J Surg.* 2021;45(2):488-95.

321 Sebikali MJ, Warwick JM, Doruyter AG. Combined versus subtraction-only technique in parathyroid scintigraphy: effect on scan interpretation. *Nucl Med Commun.* 2020;41(9):883-7.

322 Aliabadi-Wahle S, Kelly TL, Rozenfeld Y, Carlisle JR, Naeole LK, Negreanu FA, et al. Treatment strategies for primary hyperparathyroidism: what is the cost? *Am Surg.* 2014;80(11):1146-51.

323 Chen LS, Singh RJ. Niche point-of-care endocrine testing - Reviews of intraoperative parathyroid hormone and cortisol monitoring. *Crit Rev Clin Lab Sci.* 2018;55(2):115-28.

324 Clark N, Schneider DF, Vrabec S, Bauer PS, Chen H, Sippel RS. Increased efficiency of endocrine procedures performed in an ambulatory operating room. *J Surg Res.* 2013;184(1):200-3.

325 Kulkarni RP, Ituarte PH, Gunderson D, Yeh MW. Clinical pathways improve hospital resource use in endocrine surgery. *J Am Coll Surg.* 2011;212(1):35-41.

326 Mishra A, Kapoor L, Mishra SK. Post-operative care through tele-follow up visits in patients undergoing thyroidectomy and parathyroidectomy in a resource-constrained environment. *J Telemed Telecare.* 2009;15(2):73-6.

327 National Guideline C. NICE Evidence Reviews Collection. Evidence review for surgical localisation: Hyperparathyroidism (primary): diagnosis, assessment and initial management: Evidence review D. London: National Institute for Health and Care Excellence (NICE) Copyright © NICE 2019.; 2019.

328 Peel JK, Melck AL. Same-day discharge after unilateral parathyroidectomy is safe. *Can J Surg.* 2016;59(4):242-6.

329 Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *Am J Otolaryngol.* 2021;42(1):102819.

330 Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian J Endocrinol Metab.* 2019;23(3):347-52.

331 Schneider R, Kolios G, Koch BM, Fernández ED, Bartsch DK, Schlosser K. An economic comparison of surgical and medical therapy in patients with secondary hyperparathyroidism--the German perspective. *Surgery.* 2010;148(6):1091-9.

332 National Guideline C. NICE Evidence Reviews Collection. Evidence review for surgical localisation: Hyperparathyroidism (primary): diagnosis, assessment and initial management: Evidence review D. London: National Institute for Health and Care Excellence (NICE) Copyright © NICE 2019.; 2019.

333 Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *Am J Otolaryngol.* 2021;42(1):102819.

334 Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian J Endocrinol Metab.* 2019;23(3):347-52.

335 Siddiqui MI, Pasha HA, Asad R, Talati JJ. Changing paradigms in the surgical management of hyperparathyroidism at a tertiary care hospital in a developing country. *J Pak Med Assoc.* 2019;69(9):1360-4.

336 Usta A, Alhan E, Cinel A, Türkyılmaz S, Erem C. A 20-year study on 190 patients with primary hyperparathyroidism in a developing country: Turkey experience. *Int Surg.* 2015;100(4):648-55.

337 Abbott DE, Cantor SB, Grubbs EG, Santora R, Gomez HF, Evans DB, et al. Outcomes and economic analysis of routine preoperative 4-dimensional CT for surgical intervention in de novo primary hyperparathyroidism: does clinical benefit justify the cost? *J Am Coll Surg.* 2012;214(4):629-37; discussion 37-9.

338 Adler JT, Sippel RS, Chen H. New trends in parathyroid surgery. *Curr Probl Surg.* 2010;47(12):958-1017.

339 Badii B, Staderini F, Foppa C, Tofani L, Skalamera I, Fiorenza G, et al. Cost-benefit analysis of the intraoperative parathyroid hormone assay in primary hyperparathyroidism. *Head Neck.* 2017;39(2):241-6.

340 Baliski C, Nosyk B, Melck A, Bugis S, Rosenberg F, A HA. The cost-effectiveness of three strategies for the surgical treatment of symptomatic primary hyperparathyroidism. *Ann Surg Oncol.* 2008;15(10):2653-60.

341 Barczyński M. Minimally invasive parathyroidectomy without intraoperative parathyroid hormone monitoring: when and why? *J Postgrad Med.* 2009;55(4):239-40.

342 Barczyński M, Gołkowski F, Nawrot I. The current status of intraoperative iPTH assay in surgery for primary hyperparathyroidism. *Gland Surg.* 2015;4(1):36-43.

343 Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: Feasibility and outcome. *J Visc Surg.* 2018;155(4):253-8.

344 Bunch PM, Kelly HR. Preoperative Imaging Techniques in Primary Hyperparathyroidism: A Review. *JAMA Otolaryngol Head Neck Surg.* 2018;144(10):929-37.

345 Denmeade KA, Constable C, Reed WM. Use of (99m)Tc 2-methoxyisobutyl isonitrite in minimally invasive radioguided surgery in patients with primary hyperparathyroidism: A narrative review of the current literature. *J Med Radiat Sci.* 2013;60(2):58-66.

- 346 Dimas S, Michas S, Christakis I, Augoustis C, Alevizaki M. Minimally invasive parathyroidectomy in patients with previous neck surgery. *Hormones (Athens)*. 2012;11(2):160-5.
- 347 Dobrinja C, Santandrea G, Giacca M, Stenner E, Ruscio M, de Manzini N. Effectiveness of Intraoperative Parathyroid Monitoring (ioPTH) in predicting a multiglandular or malignant parathyroid disease. *Int J Surg*. 2017;41 Suppl 1:S26-s33.
- 348 Frank E, Watson W, Fujimoto S, De Andrade Filho P, Inman J, Simental A. Surgery versus Imaging in Non-Localizing Primary Hyperparathyroidism: A Cost-Effectiveness Model. *Laryngoscope*. 2020;130(12):E963-e9.
- 349 Gracie D, Hussain SS. Use of minimally invasive parathyroidectomy techniques in sporadic primary hyperparathyroidism: systematic review. *J Laryngol Otol*. 2012;126(3):221-7.
- 350 Ikeda Y, Takayama J, Takami H. Minimally invasive radioguided parathyroidectomy for hyperparathyroidism. *Ann Nucl Med*. 2010;24(4):233-40.
- 351 Karahan Ö, Okus A, Sevinç B, Eryilmaz MA, Ay S, Çaycı M, et al. Minimally invasive parathyroidectomy under local anesthesia. *J Postgrad Med*. 2013;59(1):21-4.
- 352 Kunstman JW, Kirsch JD, Mahajan A, Udelsman R. Clinical review: Parathyroid localization and implications for clinical management. *J Clin Endocrinol Metab*. 2013;98(3):902-12.
- 353 Kunstman JW, Udelsman R. Superiority of minimally invasive parathyroidectomy. *Adv Surg*. 2012;46:171-89.
- 354 Lalonde MN, Correia RD, Syktiotis GP, Schaefer N, Matter M, Prior JO. Parathyroid Imaging. *Semin Nucl Med*. 2023.
- 355 Lee L, Steward DL. Techniques for parathyroid localization with ultrasound. *Otolaryngol Clin North Am*. 2010;43(6):1229-39, vi.
- 356 Lorenz K, Dralle H. [Intraoperative parathyroid hormone determination for primary hyperparathyroidism]. *Chirurg*. 2010;81(7):636, 8-42.
- 357 Lubitz CC, Stephen AE, Hodin RA, Pandharipande P. Preoperative localization strategies for primary hyperparathyroidism: an economic analysis. *Ann Surg Oncol*. 2012;19(13):4202-9.
- 358 Melfa GI, Raspanti C, Attard M, Cocorullo G, Attard A, Mazzola S, et al. Comparison of minimally invasive parathyroidectomy under local anaesthesia and minimally invasive video-assisted parathyroidectomy for primary hyperparathyroidism: a cost analysis. *G Chir*. 2016;37(2):61-7.
- 359 Mihai R, Weisters M, Stechman MJ, Gleeson F, Sadler G. Cost-effectiveness of scan-directed parathyroidectomy. *Langenbecks Arch Surg*. 2008;393(5):739-43.
- 360 Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol*. 2016;174(1):D1-8.
- 361 Morris LF, Zanocco K, Ituarte PH, Ro K, Duh QY, Sturgeon C, et al. The value of intraoperative parathyroid hormone monitoring in localized primary hyperparathyroidism: a cost analysis. *Ann Surg Oncol*. 2010;17(3):679-85.
- 362 Opoku-Boateng A, Bolton JS, Corsetti R, Brown RE, Oxner C, Fuhrman GM. Use of a sestamibi-only approach to routine minimally invasive parathyroidectomy. *Am Surg*. 2013;79(8):797-801.
- 363 Peel JK, Melck AL. Same-day discharge after unilateral parathyroidectomy is safe. *Can J Surg*. 2016;59(4):242-6.
- 364 Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian J Endocrinol Metab*. 2019;23(3):347-52.
- 365 Schneider R, Hinrichs J, Meier B, Walz MK, Alesina PF. Minimally Invasive Parathyroidectomy without Intraoperative PTH Performed after Positive Ultrasonography as the only Diagnostic Method in Patients with Primary Hyperparathyroidism. *World J Surg*. 2019;43(6):1525-31.
- 366 Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia - A Feasibility Study. *Indian J Endocrinol Metab*. 2019;23(1):67-71.
- 367 Spiros D, Nikolaos R, Ioannis C. Minimally invasive parathyroidectomy in patients with previous endocrine surgery. *Jcls*. 2011;15(4):499-503.
- 368 Stalberg P, Delbridge L, van Heerden J, Barraclough B. Minimally invasive parathyroidectomy and thyroidectomy--current concepts. *Surgeon*. 2007;5(5):301-8.
- 369 Tolley N, Garas G, Palazzo F, Prichard A, Chaidas K, Cox J, et al. Long-term prospective evaluation comparing robotic parathyroidectomy with minimally invasive open parathyroidectomy for primary hyperparathyroidism. *Head Neck*. 2016;38 Suppl 1:E300-6.
- 370 Tublin ME, Pryma DA, Yim JH, Ogilvie JB, Mountz JM, Bencherif B, et al. Localization of parathyroid adenomas by sonography and technetium tc 99m sestamibi single-photon emission computed tomography before minimally invasive parathyroidectomy: are both studies really needed? *J Ultrasound Med*. 2009;28(2):183-90.
- 371 Udelsman R, Lin Z, Donovan P. The superiority of minimally invasive parathyroidectomy based on 1650 consecutive patients with primary hyperparathyroidism. *Ann Surg*. 2011;253(3):585-91.
- 372 Udelsman R, Pasieka JL, Sturgeon C, Young JE, Clark OH. Surgery for asymptomatic primary hyperparathyroidism: proceedings of the third international workshop. *J Clin Endocrinol Metab*. 2009;94(2):366-72.
- 373 Wang TS, Cheung K, Farrokhyar F, Roman SA, Sosa JA. Would scan, but which scan? A cost-utility analysis to optimize preoperative imaging for primary hyperparathyroidism. *Surgery*. 2011;150(6):1286-94.
- 374 Wilhelm SM, Wang TS, Ruan DT, Lee JA, Asa SL, Duh QY, et al. The American Association of Endocrine Surgeons Guidelines for Definitive Management of Primary Hyperparathyroidism. *JAMA Surg*. 2016;151(10):959-68.
- 375 Ypsilantis E, Charfare H, Wassif WS. Intraoperative PTH Assay during Minimally Invasive Parathyroidectomy May Be Helpful in the Detection of Double Adenomas and May Minimise the Risk of Recurrent Surgery. *Int J Endocrinol*. 2010;2010:178671.
- 376 Zafereo M, Yu J, Angelos P, Brumund K, Chuang HH, Goldenberg D, et al. American Head and Neck Society Endocrine Surgery Section update on parathyroid imaging for surgical candidates with primary hyperparathyroidism. *Head Neck*. 2019;41(7):2398-409.
- 377 Zammit M, Pierce K, Bailey L, Rowland M, Waghorn A, Shore S. Challenging NICE guidelines on parathyroid surgery. *Surgeon*. 2022;20(4):e105-e11.
- 378 Zanocco K, Heller M, Sturgeon C. Cost-effectiveness of parathyroidectomy for primary hyperparathyroidism. *Endocr Pract*. 2011;17 Suppl 1:69-74.
- 379 Faye M, Keita N, Lemrabott AT, Algouzmani I, Faye M, Mbengue M, et al. Surgical management of secondary hyperparathyroidism in dialysis patients in Senegal. *Saudi J Kidney Dis Transpl*. 2021;32(5):1424-30.
- 380 Mallikarjuna VJ, Mathew V, Ayyar V, Bantwal G, Ganesh V, George B, et al. Five-year Retrospective Study on Primary Hyperparathyroidism in South India: Emerging Roles of Minimally Invasive Parathyroidectomy and Preoperative Localization with Methionine Positron Emission Tomography-Computed Tomography Scan. *Indian J Endocrinol Metab*. 2018;22(3):355-61.
- 381 Miller JL, Agarwal AA, Singh KR, Sonkar AA, Kushwaha JK, Shrivastav A. Day Case Thyroid and Parathyroid Surgery: Time to Replicate Same in Developing Countries. *World J Surg*. 2015;39(8):2102-3.
- 382 Nouikes Zitouni S. Monocentric experience of primary hyperparathyroidism surgery in Algeria. *Surg Open Sci*. 2021;4:32-6.
- 383 Olatoke SA, Agodirin OS, Rahman GA, Habeeb OG, Jimoh RO, Ahmed BA, et al. Serial pathologic fractures of five long bones on four separate occasions in a patient with primary hyperparathyroidism, challenges of management in a developing country: a case report. *Pan Afr Med J*. 2013;15:45.
- 384 Paruk IM, Esterhuizen TM, Maharaj S, Pirie FJ, Motala AA. Characteristics, management and outcome of primary hyperparathyroidism in South Africa: a single-centre experience. *Postgrad Med J*. 2013;89(1057):626-31.
- 385 Rajan S, Ravindhran B, George B, Bantwal G, Ayyar V, Manjanth S. Simplified intraoperative parathormone assay for primary hyperparathyroidism in a resource-limited setting. *Biomark Med*. 2021;15(5):331-6.
- 386 Sharanappa V, Mishra A, Bhatia V, Mayilvagnan S, Chand G, Agarwal G, et al. Pediatric Primary Hyperparathyroidism: Experience in a Tertiary Care Referral Center in a Developing Country Over Three Decades. *World J Surg*. 2021;45(2):488-95.
- 387 Siddiqui MI, Pasha HA, Asad R, Talati JJ. Changing paradigms in the surgical management of hyperparathyroidism at a tertiary care hospital in a developing country. *J Pak Med Assoc*. 2019;69(9):1360-4.
- 388 Usta A, Alhan E, Cinel A, Türkyılmaz S, Erem C. A 20-year study on 190 patients with primary hyperparathyroidism in a developing country: Turkey experience. *Int Surg*. 2015;100(4):648-55.
- 389 Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: Feasibility and outcome. *J Visc Surg*. 2018;155(4):253-8.
- 390 Calò PG, Medas F, Loi G, Erdas E, Pisano G, Nicolosi A. Feasibility of unilateral parathyroidectomy in patients with primary hyperparathyroidism and negative or discordant localization studies. *Updates Surg*. 2016;68(2):155-61.
- 391 El-Hady HA, Radwan HS. Focused parathyroidectomy for single parathyroid adenoma: a clinical account of 20 patients. *Electron Physician*. 2018;10(6):6974-80.
- 392 Kandil E, Malazai AJ, Alrasheedi S, Tufano RP. Minimally invasive/focused parathyroidectomy in patients with negative sestamibi scan results. *Arch Otolaryngol Head Neck Surg*. 2012;138(3):223-5.
- 393 Kim WW, Lee YM, Sung TY, Chung KW, Hong SJ. Selection of parathyroidectomy methods for primary hyperparathyroidism according to concordance between ultrasonography and MIBI scan results. *Gland Surg*. 2021;10(1):298-306.
- 394 Kovatcheva RD, Vlahov JD, Shinkov AD, Borissova AM, Hwang JH, Arnaud F, et al. High-intensity focused ultrasound to treat primary hyperparathyroidism: a feasibility study in four patients. *AJR Am J Roentgenol*. 2010;195(4):830-5.
- 395 Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia - A Feasibility Study. *Indian J Endocrinol Metab*. 2019;23(1):67-71.
- 396 Bombil I, Louw L, Mitchell C, Mahlobo F, Mugaanza RA, Madima NR. Sonar guided focused parathyroidectomy under cervical block. *S Afr J Surg*. 2018;56(2):30-3.
- 397 Benhami A, Chuffart E, Christou N, Liva-Yonnet S, Mathonnet M. Ambulatory surgery under local anesthesia for parathyroid adenoma: Feasibility and outcome. *J Visc Surg*. 2018;155(4):253-8.

- 398 Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *Surg Clin North Am.* 2014;94(3):607-23.
- 399 Lorenz K, Dralle H. [Intraoperative parathyroid hormone determination for primary hyperparathyroidism]. *Chirurg.* 2010;81(7):636, 8-42.
- 400 Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol.* 2016;174(1):D1-8.
- 401 National Guideline C. NICE Evidence Reviews Collection. Evidence review for surgical localisation: Hyperparathyroidism (primary): diagnosis, assessment and initial management: Evidence review D. London: National Institute for Health and Care Excellence (NICE) Copyright © NICE 2019.; 2019.
- 402 Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *Am J Otolaryngol.* 2021;42(1):102819.
- 403 Slepavicius A, Beisa V, Janusonis V, Strupas K. Focused versus conventional parathyroidectomy for primary hyperparathyroidism: a prospective, randomized, blinded trial. *Langenbecks Arch Surg.* 2008;393(5):659-66.
- 404 Udelsman R, Pasieka JL, Sturgeon C, Young JE, Clark OH. Surgery for asymptomatic primary hyperparathyroidism: proceedings of the third international workshop. *J Clin Endocrinol Metab.* 2009;94(2):366-72.
- 405 Badii B, Staderini F, Foppa C, Tofani L, Skalamera I, Fiorenza G, et al. Cost-benefit analysis of the intraoperative parathyroid hormone assay in primary hyperparathyroidism. *Head Neck.* 2017;39(2):241-6.
- 406 Cheung K, Wang TS, Farrokhyar F, Roman SA, Sosa JA. A meta-analysis of preoperative localization techniques for patients with primary hyperparathyroidism. *Ann Surg Oncol.* 2012;19(2):577-83.
- 407 Frank E, Watson W, Fujimoto S, De Andrade Filho P, Inman J, Simental A. Surgery versus Imaging in Non-Localizing Primary Hyperparathyroidism: A Cost-Effectiveness Model. *Laryngoscope.* 2020;130(12):E963-e9.
- 408 Lubitz CC, Stephen AE, Hodin RA, Pandharipande P. Preoperative localization strategies for primary hyperparathyroidism: an economic analysis. *Ann Surg Oncol.* 2012;19(13):4202-9.
- 409 Mihai R, Weisters M, Stechman MJ, Gleeson F, Sadler G. Cost-effectiveness of scan-directed parathyroidectomy. *Langenbecks Arch Surg.* 2008;393(5):739-43.
- 410 Minisola S, Cipriani C, Diacinti D, Tartaglia F, Scillitani A, Pepe J, et al. Imaging of the parathyroid glands in primary hyperparathyroidism. *Eur J Endocrinol.* 2016;174(1):D1-8.
- 411 Solorzano CC, Carneiro-Pla D. Minimizing cost and maximizing success in the preoperative localization strategy for primary hyperparathyroidism. *Surg Clin North Am.* 2014;94(3):587-605.
- 412 Udelsman R, Pasieka JL, Sturgeon C, Young JE, Clark OH. Surgery for asymptomatic primary hyperparathyroidism: proceedings of the third international workshop. *J Clin Endocrinol Metab.* 2009;94(2):366-72.
- 413 Wang TS, Cheung K, Farrokhyar F, Roman SA, Sosa JA. Would scan, but which scan? A cost-utility analysis to optimize preoperative imaging for primary hyperparathyroidism. *Surgery.* 2011;150(6):1286-94.
- 414 Zammit M, Pierce K, Bailey L, Rowland M, Waghorn A, Shore S. Challenging NICE guidelines on parathyroid surgery. *Surgeon.* 2022;20(4):e105-e11.
- 415 Ahsan T, Erum U, Inam Pal KM, Jabeen R, Qureshi SG, Rehman UL, et al. The many guises of primary hyperparathyroidism: An unchanged scenario. *J Pak Med Assoc.* 2017;67(4):580-5.
- 416 Butt WT, Azim A, Abbas A, Gauhar TM, Afzal A, Azim KM. Parathyroid carcinoma. *J Coll Physicians Surg Pak.* 2012;22(9):588-90.
- 417 Fatima T, Das B, Sattar S, Jabeen S, Khan AA, Islam N. The Utility of Ultrasound in the Preoperative Localization of Primary Hyperparathyroidism: Insights from Pakistan. *Cureus.* 2020;12(8):e9835.
- 418 Junaid M, Sobani ZA, Kazi M, Khan MJ. Minimally invasive endoscopic selective parathyroidectomy. *J Pak Med Assoc.* 2012;62(4):402-5.
- 419 Latif A, Gastelum AA, Farhan K, Jagadesh S, Mutnuri S. Treatment approach for primary hyperparathyroidism in pregnancy. *Proc (Bayl Univ Med Cent).* 2020;34(1):191-3.
- 420 Siddiqui MI, Pasha HA, Asad R, Talati JJ. Changing paradigms in the surgical management of hyperparathyroidism at a tertiary care hospital in a developing country. *J Pak Med Assoc.* 2019;69(9):1360-4.
- 421 Contreras K, Baquero R, Buitrago G. Clinical and Economical Outcomes Associated with Parathyroidectomy: A 5-Year Population-Based Study in a Middle-Income Country with Universal Health Coverage. *Int J Nephrol.* 2020;2020:7250250.
- 422 Sosa JA, Tuggle CT, Wang TS, Thomas DC, Boudourakis L, Rivkees S, et al. Clinical and economic outcomes of thyroid and parathyroid surgery in children. *J Clin Endocrinol Metab.* 2008;93(8):3058-65.
- 423 Dahiya D, Abuji K, Kumari P, Gautam A, Bhadada S, Sood A, et al. Surgical outcome after focused parathyroidectomy: experience from a tertiary care centre in North India. *Pol Przegl Chir.* 2021;93(5):1-5.
- 424 Jha CK, Bichoo RA, Yadav SK. Letter to editor in response to article entitled "Ultrasound based focused neck exploration for primary hyperparathyroidism". *Am J Surg.* 2017;214(5):981-2.
- 425 Lala M. Management of Primary Hyperparathyroidism. *Indian J Surg Oncol.* 2022;13(1):143-51.
- 426 Naik AH, Wani MA, Wani KA, Laway BA, Malik AA, Shah ZA. Intraoperative Parathyroid Hormone Monitoring in Guiding Adequate Parathyroidectomy. *Indian J Endocrinol Metab.* 2018;22(3):410-6.
- 427 Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian J Endocrinol Metab.* 2019;23(3):347-52.
- 428 Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia - A Feasibility Study. *Indian J Endocrinol Metab.* 2019;23(1):67-71.
- 429 Sharma A, Patil V, Sarathi V, Purandare N, Hira P, Memon S, et al. Dual-phase computed tomography for localization of parathyroid lesions in children and adolescents with primary hyperparathyroidism. *Ann Endocrinol (Paris).* 2023.
- 430 Yadav SK, Mishra SK, Mishra A, Mayilvagnan S, Chand G, Agarwal G, et al. Surgical Management of Primary Hyperparathyroidism in the Era of Focused Parathyroidectomy: A Study in Tertiary Referral Centre of North India. *Indian J Endocrinol Metab.* 2019;23(4):468-72.
- 431 Aphale R, Damle N, Chumber S, Khan M, Khadgawat R, Dharmashaktu Y, et al. Impact of Fluoro-Choline PET/CT in Reduction in Failed Parathyroid Localization in Primary Hyperparathyroidism. *World J Surg.* 2023.
- 432 Kaur P, Gattani R, Singhal AA, Sarin D, Arora SK, Mithal A. Impact of preoperative imaging on surgical approach for primary hyperparathyroidism: Data from single institution in India. *Indian J Endocrinol Metab.* 2016;20(5):625-30.
- 433 Mallikarjuna VJ, Mathew V, Ayyar V, Bantwal G, Ganesh V, George B, et al. Five-year Retrospective Study on Primary Hyperparathyroidism in South India: Emerging Roles of Minimally Invasive Parathyroidectomy and Preoperative Localization with Methionine Positron Emission Tomography-Computed Tomography Scan. *Indian J Endocrinol Metab.* 2018;22(3):355-61.
- 434 Murugan N, Kandasamy D, Sharma R, Goyal A, Gupta AK, Tandon N, et al. Comparison of 4DMRI and 4DCT for the preoperative evaluation of patients with primary hyperparathyroidism. *Eur J Radiol.* 2021;138:109625.
- 435 Padma KS, Lakshman K, Srikanta SS. Feasibility of rapid parathormone assay for enabling minimally invasive parathyroid excision. *Indian J Surg.* 2013;75(3):210-5.
- 436 Padma S, Sundaram PS. Parathyroid scintigraphy, histopathology correlation in patients with tropical pancreatitis and coexisting primary hyperparathyroidism. *Indian J Nucl Med.* 2013;28(1):5-10.
- 437 Pradhan R, Gupta S, Agarwal A. Focused Parathyroidectomy Using Accurate Preoperative Imaging and Intraoperative PTH: Tertiary Care Experience. *Indian J Endocrinol Metab.* 2019;23(3):347-52.
- 438 Sen S, Cherian AJ, Ramakant P, Reka K, Paul MJ, Abraham DT. Focused Parathyroidectomy Under Local Anesthesia - A Feasibility Study. *Indian J Endocrinol Metab.* 2019;23(1):67-71.
- 439 Sharanappa V, Mishra A, Bhatia V, Mayilvagnan S, Chand G, Agarwal G, et al. Pediatric Primary Hyperparathyroidism: Experience in a Tertiary Care Referral Center in a Developing Country Over Three Decades. *World J Surg.* 2021;45(2):488-95.
- 440 Sreevathsa MR, Melanta K. Unilateral Exploration for Parathyroid Adenoma. *Indian J Surg Oncol.* 2017;8(2):142-5.
- 441 Badii B, Staderini F, Foppa C, Tofani L, Skalamera I, Fiorenza G, et al. Cost-benefit analysis of the intraoperative parathyroid hormone assay in primary hyperparathyroidism. *Head Neck.* 2017;39(2):241-6.
- 442 Barczyński M, Gofkowski F, Nawrot I. The current status of intraoperative iPTH assay in surgery for primary hyperparathyroidism. *Gland Surg.* 2015;4(1):36-43.
- 443 Dombrowsky A, Weiss D, Bushman N, Chen H, Balentine CJ. Can imaging studies be omitted in patients with sporadic primary hyperparathyroidism? *J Surg Res.* 2018;231:257-62.
- 444 Lorenz K, Dralle H. [Intraoperative parathyroid hormone determination for primary hyperparathyroidism]. *Chirurg.* 2010;81(7):636, 8-42.
- 445 Madorin CA, Owen R, Coakley B, Lowe H, Nam KH, Weber K, et al. Comparison of radiation exposure and cost between dynamic computed tomography and sestamibi scintigraphy for preoperative localization of parathyroid lesions. *JAMA Surg.* 2013;148(6):500-3.
- 446 Baloch MN, Aslam T, Maher M. Surgical management of hyperparathyroidism. *J Coll Physicians Surg Pak.* 2007 Nov;17(11):683-5.

**Full Text Studies Excluded With Reasons**

	<b>Author</b>	<b>Title</b>	<b>Year</b>	<b>Country &amp; (UNDP/WB Classification)</b>	<b>Study Type</b>	<b>No. of Patients</b>	<b>Reason</b>	<b>Reference</b>
1	Puthenveetil et al	Asymptomatic vs Symptomatic Primary Hyperparathyroidism: Comparison of Clinico-investigative Profile and Surgical Outcomes in Resource-Limited Setting	2022	India (Developing/Lower-Middle Income Country)	Multicentric retrospective study	213	No discussion on FP	(1)
2	Budge et al	Bone health in patients undergoing surgery for primary hyperparathyroidism at Tygerberg Hospital, Cape Town, South Africa	2020	South Africa (Developing/Upper-Middle Income Country)	Retrospective study	56	No discussion on FP	(2)
3	Badii et al	Cost–benefit analysis of the intraoperative parathyroid hormone assay in primary hyperparathyroidism	2017	Italy (Developed/High Income Country)	Retrospective study	264	Study from a developed country	(3)
4	Zanocco et al	Cost-effectiveness of parathyroidectomy for primary hyperparathyroidism	2011	USA (Developed/High Income Country)	Literature Review	-	Literature from developed countries	(4)
5	Mihai et al	Cost-effectiveness of scan-directed parathyroidectomy	2008	UK (Developed/High Income Country)	Prospective Analysis	200	Study from a developed country	(5)
6	Baliski et al	The cost-effectiveness of three strategies for the surgical treatment of symptomatic primary hyperparathyroidism	2008	Canada (Developed/High Income Country)	Retrospective study	94	Study from a developed country	(6)
7	Flynn et al	Current status of surgical techniques for parathyroidectomy for untreated primary hyperparathyroidism: Is the technology worth it?	2010	USA (Developed/High Income Country)	Literature Review	-	Literature from developed countries	(7)
8	Karakas et al	Initial Surgery for Benign Primary Hyperparathyroidism: An Analysis of 1,300 Patients in a Teaching Hospital	2014	Germany (Developed/High Income Country)	Retrospective study	1300	Study from a developed country	(8)
9	Vaid et al	Minimally invasive parathyroidectomy: a community hospital experience	2011	USA (Developed/High Income Country)	Retrospective study	188	Study from a developed country	(9)
10	Elaraj et al.	Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes	2014	USA (Developed/High Income Country)	Literature Review	-	Literature from developed countries	(10)
11	Paravastu et al.	Parathyroidectomy in a district general hospital: Outcomes and evolution in the era of minimally invasive surgery	2012	UK (Developed/High Income Country)	Prospective Analysis	368	Study from a developed country	(11)
12	Piccin et al.	Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience	2021	Italy (Developed/High Income Country)	Retrospective study	336	Study from a developed country	(12)

13	Lubitz et al	Preoperative Localization Strategies for Primary Hyperparathyroidism: An Economic Analysis	2012	USA (Developed/High Income Country)	Literature Review	-	Literature from developed countries	(13)
14	Bhadada et al.	Primary hyperparathyroidism: insights from the Indian PHPT registry	2017	India (Developing/Lower-Middle Income Country)	Retrospective study	464	No discussion on FP	(14)
15	Dumzela et al	Profile of patients operated on for primary hyperparathyroidism at Chris Hani Baragwanath Academic Hospital	2020	South Africa (Developing/Upper-Middle Income Country)	Retrospective study	77	No discussion on FP	(15)
16	Jang et al	Racial disparities in the cost of surgical care for parathyroidectomy	2018	USA (Developed/High Income Country)	Retrospective study	899	Study from a developed country	(16)
17	Anderson et al	Rural General Surgery: A Review of the Current Situation and Realities from a Rural Community Practice in Central Nebraska	2012	USA (Developed/High Income Country)	Literature Review	-	Literature from developed countries	(17)
18	Frank et al	Surgery versus Imaging in Non-Localizing Primary Hyperparathyroidism: A Cost-Effectiveness Model	2020	USA (Developed/High Income Country)	Retrospective Cost Analysis	347	Study from a developed country	(18)
19	Pradeep et al.	Systematic Review of Primary Hyperparathyroidism in India: The Past, Present, and the Future Trends	2011	India (Developing/Lower-Middle Income Country)	Systematic Review	858	No discussion on FP	(19)
20	Aliabadi-Wahle et al	Treatment strategies for primary hyperparathyroidism: what is the cost?	2014	USA (Developed/High Income Country)	Retrospective Cost Analysis	740	Study from a developed country	(20)
21	Ahmed et al	Ultrasound - first imaging modality in the detection of parathyroid adenomas	2008	South Africa (Developing/Upper-Middle Income Country)	Prospective Analysis	12	No discussion on FP	(21)
22	Fatima et al	The Utility of Ultrasound in the Preoperative Localization of Primary Hyperparathyroidism: Insights from Pakistan	2020	Pakistan (Developing/Lower-Middle Income Country)	Retrospective study	79	No discussion on FP	(22)
23	Morris et al	The Value of Intraoperative Parathyroid Hormone Monitoring in Localized Primary Hyperparathyroidism: A Cost Analysis	2010	USA (Developed/High Income Country)	Literature Review	4280	Literature from developed countries	(23)
24	Wang et al	Would scan, but which scan? A cost-utility analysis to optimize preoperative imaging for primary hyperparathyroidism	2011	USA (Developed/High Income Country)	Literature Review	-	Literature from developed countries	(24)

\*FP = Focused Parathyroidectomy

1. Puthenveetil P, Panchangam RB. Asymptomatic vs Symptomatic Primary Hyperparathyroidism: Comparison of Clinico-investigative Profile and Surgical Outcomes in Resource-Limited Setting. *Indian Journal of Surgery*. 2022;84(1):100-3.
2. Budge M, Conradie W, Beviss-Challinor K, Martin L, Conradie M, Coetzee A. Bone health in patients undergoing surgery for primary hyperparathyroidism at Tygerberg Hospital, Cape Town, South Africa. *Journal of Endocrinology, Metabolism and Diabetes in South Africa*. 2021;26(1):16-23.
3. Badii B, Staderini F, Foppa C, Tofani L, Skalamera I, Fiorenza G, et al. Cost-benefit analysis of the intraoperative parathyroid hormone assay in primary hyperparathyroidism. *Head Neck*. 2017;39(2):241-6.
4. Zanicco K, Angelos P, Sturgeon C. Cost-effectiveness analysis of parathyroidectomy for asymptomatic primary hyperparathyroidism. *Surgery*. 2006;140(6):874-81; discussion 81-2.
5. Mihai R, Weisters M, Stechman MJ, Gleeson F, Sadler G. Cost-effectiveness of scan-directed parathyroidectomy. *Langenbecks Arch Surg*. 2008;393(5):739-43.
6. Baliski C, Nosyk B, Melck A, Bugis S, Rosenberg F, H. Anis A. The cost-effectiveness of three strategies for the surgical treatment of symptomatic primary hyperparathyroidism. *Annals of Surgical Oncology*. 2008;15:2653-60.
7. Flynn MB, Civelek AC. Current status of surgical techniques for parathyroidectomy for untreated primary hyperparathyroidism: is the technology worth it? *Am Surg*. 2010;76(7):663-71.
8. Karakas E, Schneider R, Rothmund M, Bartsch D, Schlosser K. Initial Surgery for Benign Primary Hyperparathyroidism: An Analysis of 1,300 Patients in a Teaching Hospital. *World Journal of Surgery*. 2014;38(8):2011-8.
9. Vaid S, Pandelidis S. Minimally invasive parathyroidectomy: a community hospital experience. *Arch Surg*. 2011;146(7):876-8.
10. Elaraj D, Sturgeon C. Operative treatment of primary hyperparathyroidism: balancing cost-effectiveness with successful outcomes. *Surgical Clinics*. 2014;94(3):607-23.
11. Paravastu SCV, Chadwick DR. Parathyroidectomy in a district general hospital: Outcomes and evolution in the era of minimally invasive surgery. *International Journal of Surgery*. 2012;10(7):373-7.
12. Piccin O, D'Alessio P, Cioccoloni E, Burgio L, Poggi C, Altieri P, et al. Pre-operative imaging workup for surgical intervention in primary hyperparathyroidism: A tertiary referral center experience. *Am J Otolaryngol*. 2021;42(1):102819.
13. Lubitz CC, Stephen AE, Hodin RA, Pandharipande P. Preoperative localization strategies for primary hyperparathyroidism: an economic analysis. *Ann Surg Oncol*. 2012;19(13):4202-9.
14. Bhadada SK, Arya AK, Mukhopadhyay S, Khadgawat R, Sukumar S, Lodha S, et al. Primary hyperparathyroidism: insights from the Indian PHPT registry. *Journal of bone and mineral metabolism*. 2018;36:238-45.
15. Dumzela A, Bombil I. PROFILE OF PATIENTS OPERATED ON FOR PRIMARY HYPERPARATHYROIDISM AT CHRIS HANI BARAGWANATH ACADEMIC HOSPITAL. *Rheumatology*. 13:17.1.
16. Jang S, Mandabach M, Aburjania Z, Balentine CJ, Chen H. Racial disparities in the cost of surgical care for parathyroidectomy. *J Surg Res*. 2018;221:216-21.
17. Anderson RL, Anderson MA. Rural General Surgery: A Review of the Current Situation and Realities from a Rural Community Practice in Central Nebraska. *Online Journal of Rural Research & Policy [Internet]*. 2012; 7(2).
18. Frank E, Watson W, Fujimoto S, De Andrade Filho P, Inman J, Simental A. Surgery versus Imaging in Non-Localizing Primary Hyperparathyroidism: A Cost-Effectiveness Model. *Laryngoscope*. 2020;130(12):E963-e9.
19. Pradeep P, Jayashree B, Mishra A, Mishra S. Systematic review of primary hyperparathyroidism in India: the past, present, and the future trends. *International journal of Endocrinology*. 2011;2011.
20. Aliabadi-Wahle S, Kelly TL, Rozenfeld Y, Carlisle JR, Naeole LK, Negreanu FA, et al. Treatment strategies for primary hyperparathyroidism: what is the cost? *Am Surg*. 2014;80(11):1146-51.
21. Ahmed R, Khan N, Ellemidin S, Gayaparsad K. Ultrasound-first imaging modality in the detection of parathyroid adenomas. *SA Journal of Radiology*. 2008;12(3):64-8.
22. Fatima T, Das B, Sattar S, Jabeen S, Khan AA, Islam N. The Utility of Ultrasound in the Preoperative Localization of Primary Hyperparathyroidism: Insights from Pakistan. *Cureus*. 2020;12(8):e9835.
23. Morris LFMD, Zanicco KMD, Ituarte PH, G P, Ro K, Duh Q-yMDF, et al. The Value of Intraoperative Parathyroid Hormone Monitoring in Localized Primary Hyperparathyroidism: A Cost Analysis. *Annals of Surgical Oncology*. 2010;17(3):679-85.
24. Wang TS, Cheung K, Farrokhyar F, Roman SA, Sosa JA. Would scan, but which scan? A cost-utility analysis to optimize preoperative imaging for primary hyperparathyroidism. *Surgery*. 2011;150(6):1286-94.